



Scholars Research Library

Der Pharmacia Lettre, 2012, 4 (2):436-442
(<http://scholarsresearchlibrary.com/archive.html>)



Anticataract and antioxidant activities of *Achyranthes aspera* Linn. against calcium-induced cataractogenesis using goat lenses

Muthuswamy Umamaheswari*, Sundaram Dhinesh, Kuppusamy Asokkumar, Thirumalaiswamy, Sivashanmugam, Varadharajan Subhadradevi, Puliyath Jagannath and Arumugam Madeswaran

Department of Pharmacology, College of Pharmacy, Sri Ramakrishna Institute of Paramedical Sciences, Coimbatore, Tamilnadu, India

ABSTRACT

The present study evaluated the *in vitro* anticataract and antioxidant activities of aqueous leaves extract of *Achyranthes aspera* Linn. against calcium-induced cataractogenesis using goat lenses. Transparent isolated goat lenses were incubated in artificial aqueous humor and divided into seven experimental groups. The extracts at a dose of 100 µg/ml and 200 µg/ml were incubated simultaneously with calcium (10 mM) for a period of 16 h. Vitamin E (100 µg/ml) was used as the standard drug. At the end of the incubation, levels of various biochemical parameters such as protein content, malondialdehyde (MDA), lipid hydroperoxides (LH), Cu²⁺-induced lipoprotein diene formation, Ca²⁺ATPase and enzymatic antioxidants like catalase (CAT), glutathione peroxidase (GPx), superoxide dismutase (SOD), glutathione reductase (GSSH) and non-enzymatic antioxidant like reduced glutathione (GSH) were measured in the lens homogenate. Incubation with calcium chloride produced a mature cataract and there was a significant increase in LH and MDA and a decrease in protein content, Ca²⁺ATPase, Cu²⁺-induced lipoprotein diene formation and enzymatic and non-enzymatic antioxidants when compared to normal control. Our results indicated that simultaneous incubation of the plant extracts prevented the preoxidative damage caused by calcium chloride, which is evidenced from the improved antioxidant potential. The aqueous leaves extract of *Achyranthes aspera* protected the lens against calcium-induced oxidative damage which might be helpful in delaying the progression of cataract.

Key words: antioxidant; cataract; calcium chloride; *Achyranthes aspera*.

INTRODUCTION

Cataract is a major cause of blindness worldwide and more so in developing countries. Surgery is the only effective treatment for cataract and the exact mechanism is not clear. Although the surgery is recognized as being one of the safest procedure, there are significant rate of complications, leading to irreversibly blind eyes [1]. Studies are being conducted to explore the mechanism of cataractogenesis using various models of cataract and to target crucial steps to stop this process. Limitations in accessibility, acceptability, and affordability of cataract surgical services make it more relevant and essential to look into alternative pharmacological measures for treatment of this disease. Thus, much eagerness is being laid on identification of natural compounds that will help to inhibit cataractogenesis [2].

Increased levels of lenticular calcium activate calcium dependent proteases. The activated proteases hydrolyze cytoskeletal proteins and lens crystalline. Crystalline cleavage would effect from lower molecular weight peptides that could, in turn, aggregate to produce higher molecular weight proteins [3].

In addition oxidative stress implicated in the cataract induced by glucose and age related process due to the development of superoxide ($O_2^{\cdot-}$) radicals and H_2O_2 because of these free radicals are eagerly react with the biomolecules [4]. The toxic effects of the reactive oxygen species are neutralized in the lens by antioxidants such as vitamin E, ascorbic acid, the glutathione system (GSH peroxidase, GSH reductase), superoxide dismutase and catalase. The enzymatic (glutathione peroxidase, catalase, superoxide dismutase) and non-enzymatic (glutathione, ascorbate, cysteine) antioxidant system activities are decreased in the lens and aqueous humor during aging and in the development of cataract [5].

Historically plants have been used in folk medicine to treat various diseases and are rich natural sources of antioxidants. Researchers have examined the effect of plants used by indigenous people to treat disorders of the eye [6].

Acyranthus aspera Linn. belonging to the family Amaranthaceae is widely scattered in southern region of India. The leaves are reported to possess alkaloids, flavanoids, saponins, tannins and phenolic compounds. The leaves of this species are used in Indian traditional medicine for the treatment of ophthalmic and other eye infections, it also has antifertility, nephroprotective, post-coital antifertility, antioxidant, reduce inflammatory swellings, nephroprotective and immunomodulatory activities [6]. Various studies have been reported for its hypoglycaemic effect [7] anti cancer [8] anti fungal [9] and increase the thyroid hormone levels [10].

The main objective of the study was to evaluate the aqueous extract of leaves of *Acyranthus aspera* Linn. for its *in vitro* anticataract and antioxidant activities against calcium-induced cataractogenesis using goat lenses.

MATERIALS AND METHODS

Drugs and chemicals

Calcium chloride and vitamin E were obtained from SD fine chemicals, Mumbai. 1-amino 2-naphthol-4-sulfonic acid, 5, 5-dithiobis-(2-nitrobenzoic acid) (DTNB), Nitro blue tetrazolium chloride (NBT), Nicotinamide adenine dinucleotide reduced salt (NADH), Nicotinamide adenine dinucleotide phosphate reduced tetra sodium salt (NADPH), Oxidised glutathione, Reduced glutathione, Adenosine-5'-triphosphate (ATP) were obtained from Himedia Laboratories Ltd., Mumbai. Fresh goat lenses were obtained from the slaughterhouse, Coimbatore. All other chemicals used in the study were obtained commercially and were of analytical grade.

Plant collection and authentication

The leaves of *Acyranthus aspera* Linn. were collected from Pachai hills, Salem, Tamilnadu, India during the month of June 2010. The plant was identified and authenticated by Mr. G.V.S. Murthy, Joint Director, C-I/C, Botanical survey of India, Tamil Nadu Agricultural University Campus, Coimbatore bearing the reference number BSI/SRC/5/23/10-11/Tech-595.

Plant extract

Achyranthes aspera aqueous leaf extract

The fresh leaves were shade dried, powdered and 25 g of the powder was soaked separately with 250 ml of sterile distilled water and kept at room temperature for 24 hrs and macerated using a mechanical shaker for 4 h. The extract was filtered through muslin cloth and the marc was again soaked with the same volume of water for 12 h and then further extracted using a mechanical shaker for 4h and filtered. The filtrates were then combined concentrated at 40°C [10]. The *Achyranthes aspera* Linn. leaves extract was abbreviated as ALE.

Calcium-induced cataract

Fresh goat eyeballs were obtained from a local slaughterhouse within two hours after killing of the animals and the lenses were isolated. Fresh goat lenses were incubated for 16 hours at 37°C in 10 mM $CaCl_2$ solution made in Tris-HCL buffer (0.01 mM) pH-7.4 [11].

A total of 30 goat lenses were used and divided into five experimental groups consisting of 6 in each group:

Group I : Tris-HCL (solvent control)

Group II: CaCl₂ 10 mM alone (negative control)

Group III: ALE (100 µg/ml) + CaCl₂ 10 mM

Group IV: ALE (200 µg/ml) + CaCl₂ 10 mM

Group V: Vitamin E (100 µg/ml) + CaCl₂ 10 mM

Examination of lens opacity

To study the opacity of the lens, the lenses from the control and experimental groups were placed on a wire mesh and photographed [12].

Preparation of lens homogenate

After incubation, lenses were homogenized in 10 volumes of 0.1M potassium phosphate buffer, pH 7.0. The homogenate was centrifuged at 10,000 rpm for 1 h and the supernatant was used for estimation of biochemical parameters [13].

Biochemical parameters

Estimation of total protein content

To 0.1 ml of lens homogenate, 4.0ml of alkaline copper solution was added and allowed to stand for 10min. Then, 0.4 ml of phenol reagent was added very rapidly and mixed quickly and incubated in room temperature for 30 mins for colour development. Reading was taken against blank prepared with distilled water at 610 nm in UV-visible spectrophotometer. The protein content was calculated from standard curve prepared with bovine serum albumin and expressed as µg/mg lens tissue [14].

Estimation of lipid hydroperoxides (LH)

About 0.1ml of lens homogenate was treated with 0.9 ml of Fox reagent (188 mg butylated hydroxytoluene (BHT), 7.6 mg xylenol orange and 9.8 ammonium ion sulphate were added to 90 ml of methanol and 10ml 250 mM sulphuric acid) and incubated for 3 min. The colour developed was read at 560 nm using a colorimeter. The values are expressed as nmoles/ mg lens protein [15].

Estimation of malondialdehyde (MDA)

Lenses were homogenized in 10% (w/v) 0.1 M Tris-HCl buffer (pH 7.5). One milliliter of the homogenate was combined with 2 ml of TCA-TBA-HCl reagent 15% trichloroacetic acid (TCA) and 0.375% thiobarbituric acid (TBA) in 0.25 N HCl and boiled for 15 min. Precipitate was removed after cooling by centrifugation at 1000g for 10 min and absorbance of the sample was read at 535 nm against a blank without tissue homogenate. The values are expressed as nmoles of MDA/ min/ mg lens protein [16].

Inhibition of Cu²⁺ induced lipoprotein diene formation

Lens homogenate was diluted to 0.67% in phosphate buffered saline. Control experiments consisted of identical assay conditions but without the sample. Oxidation was initiated immediately after addition of sample by the addition of 12 µM final concentration of Cu²⁺ added as CuSO₄·5H₂O dissolved in deionized distilled water. Oxidation was determined by measuring the absorbance at 234 nm using a UV-Visible Spectrophotometer. Absorbance was taken after 120 mins at 37°C. The lipoprotein diene formation was measured from the absorbance at a time. The absorbance provides an indication of protection of tissue lipoprotein against oxidation [16].

Assay of Ca²⁺-ATPase activity

To added 0.1 ml of the lens homogenate (10% (w/ v) in 0.25 M sucrose) and add 0.2 ml of the substrate, ATP. The tubes were incubated for 30 min in a water bath at 37°C. The enzyme activity is stopped by adding 2 ml of 10% TCA. Then 0.2 ml of ATP is added and the same kept in ice for 20 min. All the tubes were then centrifuged at 2500 rpm for 10 min and the supernatant collected. The protein free supernatant was analyzed for inorganic phosphate. For that 3 ml of the supernatant was treated with 1 ml of ammonium molybdate and 0.4 ml 1-amino-2-naphthol-4-sulfonic acid (ANSA). The colour developed was read at 680 nm after 20 min and the inorganic phosphate value expressed as nm three parallel experiments were conducted [17].

Determination of enzymatic antioxidants*Assay of superoxide dismutase (SOD)*

The assay mixture contained 1.2 ml sodium pyrophosphate buffer (0.052 M, pH 8.3), 0.1 ml of 186 μ M Phenazonium methosulphate (PMS), 0.3 ml of 300 μ M NBT, 0.2 ml of 780 μ M NADH, 1.0 ml homogenate (lens were homogenized in 10% (w/v) 0.25 M sucrose buffer) and distilled water to a final volume of 3.0 ml. Reaction was started by the addition of NADH and incubated at 30°C for 1 min. The reaction was stopped by the addition of 1.0 ml glacial acetic acid and the mixture stirred vigorously. 4.0 ml of n-butanol was added to the mixture and shaken well. The mixture was allowed to stand for 10 min, centrifuged, the butanol layer taken out and the absorbance was measured at 560 nm against a butanol blank. A system devoid of enzyme served as the control and three parallel experiments were conducted [18].

Assay of catalase (CAT)

The reaction mixture contained 2.0 ml of homogenate (lens were homogenized in 10% (w/v) 50 mM phosphate buffer, pH 7.0) and 1.0 ml of 30 mM hydrogen peroxide (in 50 mM phosphate buffer, pH 7.0). A system devoid of the substrate (hydrogen peroxide) served as the control. Reaction was started by the addition of the substrate and decrease in absorbance monitored at 240 nm for 30 s at 25°C. The difference in absorbance per unit time was expressed as the activity and three parallel experiments were conducted. One unit is defined as the amount of enzyme required to decompose 1.0 M of hydrogen peroxide per minute at pH 7.0 and 25°C [19].

Estimation of glutathione reductase (GSSH)

The enzyme activity was determined spectrophotometrically by the decrease in absorbance of NADPH at 340 nm. The reaction mixture contained 2.1 ml of 0.25 mM, potassium phosphate buffer pH 7.6, 0.1 ml of 0.001 M NADPH, 0.2 ml of 0.0165 M oxidised glutathione and 0.1 ml (10 mg/ml) of bovine serum albumin (BSA). The reaction was started by the addition of 0.02 ml of lens homogenate with mixing and the decrease in absorbance at 340 nm was measured for 3 min against a blank. Glutathione activity was expressed as nmoles NADPH oxidized/min/ mg lens protein at 30°C [20].

Estimation of glutathione peroxidase (GPx)

The reaction mixture consists of 0.2 ml of 0.4 M Tris buffer, 0.1 ml of sodium azide, 0.1 ml of hydrogen peroxide, 0.2 ml of glutathione and 0.2 ml of lens homogenate supernatant incubated at 37°C for 10 min. The reaction was arrested by the addition of 10% TCA and the absorbance was taken at 340 nm. Activity was expressed as nmoles/min/mg lens protein [21].

Determination of non enzymatic antioxidant:*Estimation of glutathione (GSH)*

Lenses were homogenized in 10% (w/v) cold 20 mM EDTA solution on ice. After deproteinization with 5% TCA, an aliquot of the supernatant was allowed to react with 150 μ M DTNB [5, 5-dithiobis-(2-nitrobenzoic acid)]. The product was detected and quantified spectrophotometrically at 416 nm. Pure GSH was used as standard for establishing the calibration curve and three parallel experiments were conducted [22].

Statistical analysis

Statistical analysis was carried out by using one-way analysis of variance (ANOVA) followed by Dunnett's test. Results are expressed as mean \pm SEM of six lenses in each group. P values < 0.05 were considered significant.

Table 1. Phytochemical screening


Phytochemicals	<i>Achyranthes aspera L.</i>
Alkaloids	+
Flavonoids	+
Saponins	+
Tannins and phenolics	+
Terpenoids	+

RESULTS

Phytochemical screening

The aqueous leaf extract of *Achyranthes aspera* Linn. was subjected to the phytochemical screening and for the presence of various phytochemical constituents.

Fig 1. Photographs of lenses in normal and experimental groups incubated with calcium



Photographs of lenses in normal and experimental groups incubated with calcium chloride are shown in Fig 1(a-e).

Fig 1a shows the normal lens Fig 1b is the lens incubated with calcium chloride (10 mM) for a period of 16 hrs showing complete opacification of the lens fibres compared to normal control. Fig 1c are the lenses incubated simultaneously with calcium chloride (10 mM) and ASE at a concentration of 100 µg/ml showing a decrease in opacity compared to cataractous lenses. Fig 1d are the lenses incubated simultaneously with calcium chloride (10 mM) and ALE at a concentration of 200 µg/ml showing a decrease in opacity compared to cataractous lenses. Fig 1e

is the lens incubated with calcium chloride (10 mM) and vitamin E (100 µg/ml) showing almost normal transparency when compared to cataractous lenses.

Effect of the aqueous leaf extract of *Achyranthes aspera* Linn. on lens protein and lipid peroxidation in control and experimental groups

There was a significant ($P < 0.01$) decrease in the level of total protein, Ca^{2+} ATPase, Cu^{2+} -induced lipoprotein diene formation and an increase in the level of malondialdehyde and lipid hydroperoxides in glucose-induced cataractous lenses when compared to normal control. Incubation with the aqueous leaf extract of *Achyranthes aspera* at dose of 100 & 200 µg/ml and Vitamin E (100 µg/ml) simultaneously with calcium chloride for 16 h caused a significant ($P < 0.01$) increase in the total protein, Ca^{2+} ATPase, Cu^{2+} -induced lipoprotein diene formation and a decrease in the level of malondialdehyde and lipid hydroperoxides (Table 2).

Table 2. Effect of the aqueous leaf extract of *Achyranthes aspera* Linn. on lens protein, MDA, LH, Ca^{2+} ATPase and Cu^{2+} induced lipoprotein diene in control and experimental groups

GROUP	Protein (mmoles/min/ mg lens tissue)	MDA (µmoles/min/mg protein)	LH (nmoles/min/mg protein)	Ca^{2+} ATPase (µmoles/min/mg inorganic phosphate)	Cu^{2+} induced Lipoprotein diene absorbance
Normal control	116.29±4.96	0.117±0.003	2.75±0.22	4.65±0.18	0.8522±0.07
Calcium control	64.23±4.28 ^a	0.66±0.05 ^a	8.675±1.17 ^a	1.19±0.11 ^a	0.1983±0.03 ^a
ALE (100 µg/ml)	90.95±3.94 ^b	0.36±0.008 ^b	4.57±0.27 ^b	3.61±0.39 ^b	0.5836±0.04 ^b
ALE (200 µg/ml)	96.31±3.07 ^b	0.29±0.01 ^b	3.96±0.38 ^b	3.89±0.38 ^b	0.6369±0.07 ^b
Vitamin-E (100 µg/ml)	108.52±4.77 ^b	0.19 ^b	3.08±0.2 ^b	4.14±0.18 ^b	0.7738±0.07 ^b

Values are mean ± SEM; n=6 in each; ^a $P < 0.01$ when compared to normal control;
^b $P < 0.01$ when compared to calcium control (one way ANOVA followed by Dunnett's test).

Effect of the aqueous leaf extract of *Achyranthes aspera* Linn. on lens enzymatic and non enzymatic antioxidants in control and experimental groups

Incubation with calcium chloride 10 mM for 16 h produced a significant ($P < 0.01$) decrease in the enzymatic antioxidants like catalase, superoxide dismutase, peroxidase, glutathione peroxidase and glutathione reductase and the non-enzymatic antioxidant reduced glutathione in the lens homogenate when compared to normal control. Incubation with the aqueous leaf extract of *Achyranthes aspera* at doses of (100 & 200 µg/ml) and Vitamin E simultaneously with glucose significantly ($P < 0.01$) restored the levels of both enzymatic and non enzymatic antioxidant enzymes which is almost similar to the control group (Table 3).

Table 3. Effect of the aqueous leaf extract of *Achyranthes aspera* Linn. on lens enzymatic and non enzymatic antioxidants in control and experimental groups

GROUP	Catalase (mmoles/min/ mg protein)	GPx (µmoles/min/ mg protein)	SOD (µmoles/min/ mg protein)	GSSH (µmoles/min/ mg protein)	GSH (µmoles/min/ mg protein)
Normal control	1.61±0.05	3.18±0.11	5.37±0.3	2.42±0.27	3.81±0.25
Calcium control (10 mM)	0.302±0.02 ^a	0.64±0.08 ^a	1.87±0.24 ^a	0.42±0.04 ^a	1.07±0.07 ^a
ALE (100 µg/ml)	1.09±0.116 ^b	2.5±0.24 ^b	3.64±0.27 ^b	1.85±0.19 ^b	2.95±0.24 ^b
ALE (200 µg/ml)	1.15±0.11 ^b	2.96±0.17 ^b	4.45±0.48 ^b	2.01±0.19 ^b	3.04±0.2 ^b
Vitamin-E (100 µg/ml)	1.45±0.15 ^b	3.07±0.19 ^b	5.07±0.26 ^b	2.18±0.29 ^b	3.45±0.21 ^b

Values are mean ± SEM; n=6 in each; ^a $P < 0.01$ when compared to normal control;
^b $P < 0.01$ when compared to calcium control (one way ANOVA followed by Dunnett's test).

The aqueous leaves extract of *Achyranthes aspera* protected the lens against calcium-induced oxidative damage which might be helpful in delaying the progression of cataract.

CONCLUSION

To conclude, the study suggested that the aqueous extract of *Achyranthes aspera* L. leaves possess anticataract and antioxidant activities, which might be helpful in preventing or slowing the progress of cataract. Further *in vivo* studies and investigations on the isolation and identification of active components in the leaves may lead to chemical entities with potential for clinical use in the prevention and treatment of cataract.

REFERENCES

- [1] M.S. Moghaddam, A.P. Kumar, G.B. Reddy, V.S. Ghole, *J. Ethnopharmacol.*, **2005**, 97, 397.
- [2] R. Manikandan, R. Thiagarajan, S. Beulaja, S. Chindhu, K. Mariammal, G. Sudhandiran, M. Arumugam, *Chem. Biol. Inter.* **2009**, 181, 202.
- [3] J. Wang, A. Bhatnagar, H. Naseem, P. Dhir, K. Satish, S. K. Srivastava, *Invest. Ophthalm. Visual science*, **1996**, 37, 915.
- [4] Z. Kyselova, M. Stefek, V. Bauer, *J Diab. & its Compl.*, **2004**, 18, 129.
- [5] D. Ozmen, E. Erkin, I. Guner, S. Habil, O. Bayindir, *Turkish J. Endocrin. & Metab.*, **2000**, 1, 1.
- [6] M. Umamaheshwari, T. K. Chatterjee, *Inter. J. Biomed. & Pharm. Sci.*, **2008**, 2, 70.
- [7] M. S. Akhtar, J. Iqbal, *J. Ethnopharmacol.*, **1991**, 3, 49.
- [8] A. Chakraborty, A. Brantner, T. Mukainaka, Y. Nobukuni, H. Tokuda, H. Nishino, *Cancer letters*, **2002**, 177, 1.
- [9] E.K. Elumalai, N. Chandrasekaran, T. Thirumalai, C. Sivakumar, V.S. Therasa, E. David, *Int. J. Pharmtech Res.*, **2009**, 1, 1576.
- [10] P. Tahiliani, K. Anand, *J. Ethnopharmacol.*, **2000**, 71, 527.
- [11] D. V. Rai, K. S. Kohli, *Polish J. Med. & Phys. Eng.*, **2006**, 12, 1.
- [12] J. Sanderson, J. M. Marcantonio, G. A. Duncan, *Invest. Ophthalm. & Vis. Sci.*, **2000**, 41, 2255.
- [13] H. Son, H. Kim, Y. H. Kwon, *J. Nut. Sci. Vit.*, **2007**, 53, 324.
- [14] O. H. Lowry, N. J. Rosenbourgh, A. L. Farr, R. J. Randall, *J. Biol. Chem.*, **1951**, 193, 265.
- [15] W. G. Niehaus, B. Samuelsson, *Eur. J. Biochem.*, **1968**, 6, 126.
- [16] J. M. Hodyson, J. M. P. Foot, K. D. Croft, I. B. Puddy, T. A. Mori, L. J. Beilin, *J. Sci. of Food & Agri.*, **1999**, 79, 561.
- [17] G. Rorive, A. Kleinzeller, *Methods Enzymol.*, **1974**, 32, 303.
- [18] P. Kakkar, B. Das, P. N. Viswanathan, *Ind. J. Biochem. & Biophy.* 1984, 21, 130.
- [19] H. Aebi, *Methods Enzymol.*, **1984**, 105, 121.
- [20] I. Carlberg, B. Mannervik, *Methods Enzymol.*, **1985**, 113, 484.
- [21] A. L. Tappel, *Methods Enzymol.*, **1978**, 52, 506.
- [22] G. L. Ellman, *Arch bio-chem biophys* **1959**, 82, 70.