

Effect of salicylic acid and salinity on some morphological characteristics of Aloe Vera

Nasrin Miri*, Forough Mortezaee Nejad, Hossein Zeinali

¹Department of Horticulture, Faculty of Agriculture, Khorasgan Branch, Islamic Azad University, Isfahan, Iran

*Corresponding email: yas.rahimiyan.yr@gmail.com

(Received on:21-7-14)

(Accepted on:2-8-14)

ABSTRACT

Aloe Vera plant belongs to Liliaceous family and it is one of the important herbs. Importance of this plant is for its leaves and gel. Since salinity stress can be effective on the quality of some characteristics of aloe vera, this study performed to investigate the effect of salicylic acid and salinity on some morphological characteristics of Aloe vera plant. Completely randomized factorial design with 3 replicates was used and data were analyzed by using the General Linear Model Procedure of SAS, 2001. Data from this study showed that salicylic acid and salinity had significant effect on biochemical and morphological characteristics for Aloe Vera. Data showed that leave diameter was affected by salinity and use salicylic acid (0.05 mm) could significant effect on leave diameter, Also Increase of salinity could reduce wet and dried gel weight, but Interaction between salinity and Salicylic acid (SA) could reduce salinity effects. In conclusion data showed that salicylic could damp some effect of salinity on Aloe Vera plant.

Key words: Aloe Vera, Salicylic acid, Salinity, Morphological characteristics.

INTRODUCTION

Aloe vera L. is a perennial liliaceous plant with succulent green leaves joined at the stem in a whorled pattern. It is highly appreciated due to its short growth period and high economic value among all the aloe species, and is used in pharmaceuticals, folk medicine, healthcare, cosmetic products and food products [1]. There is, however, little scientific evidence of the effectiveness or safety of Aloe Vera extracts for either cosmetic or medicinal purposes, and what positive evidence is available is frequently contradicted by other studies [2,4,6]. Aloe vera, an anti-oxidant rich plant, contains vitamins such as A, C, and E plus the minerals, zinc, and selenium. Anti-oxidants help boost the immune system and combat free radicals in the body. Another component of Aloe Vera consists of the lignins, a major structural material of cellulose content that allows for penetrative properties. [10, 13]. Aloe Vera contains salicylic acid which is an aspirin-like compound with anti-inflammatory, analgesic, and anti-bacterial properties. It has anti-pyretic properties for reducing fevers. Other constituents of Aloe vera would include prostaglandins, tannins, magnesium lactate, resins, mannins and proteins such as lectins, monosulfonic acid and gibberlin [22,24]. Salt stress is a limiting factor of plant growth and yield and becoming a serious problem in the world [3,8,11]. Salinity stress is one of the most important abiotic factors that it can reduce the growth on dry and semi-dry plants [14,15,20,23,26,27]. Since, there are many dry lands available in Iran study about salinity for growth factor is important. Salicylic acid (SA) is a monohydroxybenzoic acid, a type of phenolic acid and a beta hydroxy acid [6,7]. This colorless crystalline organic acid is widely used in organic synthesis and functions as a plant hormone. SA is a phenolic phytohormone and is found in plants with roles in plant growth and development, photosynthesis, transpiration, ion uptake and transport. SA also induces specific changes in leaf anatomy and

chloroplast structure. SA is involved in endogenous signaling, mediating in plant defense against pathogens. [16] It plays a role in the resistance to pathogens by inducing the production of pathogenesis-related proteins. [18] It is involved in the systemic acquired resistance in which a pathogenic attack on one part of the plant induces resistance in other parts. The signal can also move to nearby plants by salicylic acid being converted to the volatile ester, methyl salicylate [12]. SA which is produced by root cells plays a prominent role in various physiological processes such as plant's growth, ions uptake, photosynthesis and germination [7,18]. Since SA is one of the important compounds to resistance from plant stress this study is about effect of salicylic acid and salinity on some morphological characteristics of Aloe Vera plant.

MATERIALS AND METHODS

The experiment was conducted in a greenhouse in (2013 - 2014) in the agricultural faculty, university of Khorasgan, Isfahan, Iran. This experiment was conducted in bi-factorial design in completely randomized design with three replications. At the end of experiment data were collected and analyzed by using the General Linear Model Procedure of (SAS, 2001). Differences between means were analyzed by Duncan's multiple ranges test and P value less than 0.05 was considered as significant [19].

Plant material:

Sprouts of Aloe. Vera with 12 cm length obtained in vitro culture used in this work.

Culture conditions and treatments:

Aloe sprouts were cultured in greenhouse in pots with 20 cm diameter containing cocopeat and perlite and were irrigated with nutrient solutions containing different levels of NaCl (0, 24, 12, and 36 ds/m) and spraying Salicylic acid (0,0.5,1 and 1.5 mM) on leaves. After 9 month from growth period some characteristics such as leaves diameter and depth of them, wet gel weight, dry gel weight, gel Ph and EC were evaluated.

RESULTS

Data showed that leaves depth in E0. S0 and E0. S0.5 was higher than other and leaves diameter was higher for E0. S1.5, E0. S1 and E12.S0.5 respectively ($p < 0.05$). Also EC gel were higher when E24 and E36 were used. The pH was affected by of experimental treatments ($p < 0.05$). Wet gel and dry gel weight were higher when E0. S0 and E0.S0.5 were applied and the minimum weight of wet and dried gel was for E36. S1.5 ($p < 0.05$).

Table -1 Mean comparison of morphological characteristics of Aloe Vera

Treatment	leave depth	leave diameter	EC gel	pH	Wet gel weight	Dried gel weight
E0.S0 ^r	0.40 ^{de}	14.26 ^c	3.34 ^j	4.76 ^c	11.58 ^b	1.51 ^a
E0.S0.5	0.40 ^{de}	15.76 ^a	2.74 ^l	4.64 ^d	13.06 ^a	1.14 ^b
E0. S1	0.13 ^e	14.41 ^b	2.76 ^l	4.74 ^c	8.70 ^{de}	0.60 ^h
E0. S1.5	0.17 ^e	15.76 ^a	3.22 ^k	4.94 ^a	7.45 ^f	1.02 ^c
E12.S0	0.63 ^{cd}	11.44 ^d	6.52 ^h	4.74 ^c	8.19 ^{ef}	0.93 ^e
E12.S0.5	0.63 ^{cd}	14.00 ^c	5.93 ⁱ	4.90 ^a	10.72 ^c	0.97 ^d
E12.S1	0.73 ^{cd}	11.31 ^d	5.92 ^j	4.75 ^c	4.26 ^g	0.79 ^f
E12. S1.5	0.80 ^c	9.68 ^e	6.52 ⁱ	4.82 ^b	8.77 ^d	0.78 ^g
E24. S0	1.80 ^b	8.57 ^g	7.94 ^d	4.61 ^d	1.72 ^j	0.20 ⁿ
E24. S0.5	2.20 ^a	6.81 ^h	7.32 ^g	4.84 ^b	4.31 ^g	0.27 ^j
E24. S1	1.83 ^{ab}	9.06 ^f	7.57 ^e	4.76 ^c	2.54 ⁱ	0.24 ^l
E24. S1.5	1.73 ^b	5.30 ⁱ	7.45 ^f	4.64 ^d	3.44 ^h	0.27 ⁱ
E36. S0	2.23 ^a	5.58 ⁱ	10.33 ^a	4.65 ^d	2.56 ^l	0.19 ^o
E36. S0.5	1.97 ^a	5.37 ⁱ	9.44 ^c	4.74 ^c	2.64 ⁱ	0.22 ^m
E36. S 1	2.23 ^a	4.55 ^j	9.74 ^b	4.65 ^d	2.79 ⁱ	0.25 ^k
E36.S1.5	1.77 ^b	4.89 ^j	9.45 ^c	4.64 ^f	1.38 ^j	0.16 ^p

* E = Salinity, S = Salicylic acid

DISCUSSION

Salinity is one of the main problems which can negatively affect soil fertility and productivity. Salinity of soil can lead to osmotic stress, decrease in available water, ionic stress and cells' ionic imbalance. Salinity decreases germination, fresh and dry weight of radical and plumule in germination phase [20,23,26,27].

Some researchers showed that the impact of salt stress has been correlated with some morphological and physiological traits like reduction in fresh and dry weight[1]. In fact, salinity affects plant metabolism by disturbing physiological and biochemical processes of plants due to ionic and osmotic imbalances which results in the reduction of plant growth and productivity[3]. The deleterious effects of salinity on plant growth are associated with low osmotic potential of soil solution, nutritional imbalance, specific ion effect, or a combination of these factors[6,9]. Zan et al (2007), studying the physiological and ecological characters on Aloe Vera under seawater irrigation (EC= 23.4 ds m⁻¹) reported that salinity stress caused a decrease in tissue water, total soluble sugars and glucose[25]. Moghbeli et al(2012) ,showed that salinity affected the number of leave, plant height, number of sprout, root weight, plant weight, leave weight, total gel weight, and root dry weight[11].

Moghbeli et al, (2012) had shown , salinity decreased plant leaves and roots growth and their dry matter all measured characteristics showed differences between control and 2,4,6 and 8 ds/m salinity[11]. Zan et al , (2007) indicated that salt stress led to decreasing TSS in both cultivars of Aloe Vera and it is a limiting factor of plant growth and yield[11]. Fuentes et al, (1988) and (Fuentes and Rodriguez, 1988) reported similar report and revealed that the height of leaves and sprouts reduced with increasing salinity in different Aloe spp[5]. Senaratna et al. (2000) reported that the effective dose of SA for alleviating the injury of tomato and bean plants under water stress, ranged from 0.1 to 0.5 mM[21].

Abdullahi et al,(2011) showed that the comparison of means showed that the effect of various levels of salicylic acid on all characteristics was separate at different statistical levels[1]. Salicylic acid is a phenolic substance. Phenols are defined as compounds with a hydroxyl group with active derivatives. Salicylic acid is found in many plants [7,12,16]. Talebi et al. (2012) had shown treating the borage seeds with salicylic acid has increased root's fresh weight[23]. Abdullahi et al,(2011) shown that plant growth and plant gel TSS levels increased as a result of salicylic acid treatment[1]. The axial role of salicylic acid in translating messages as a defense response against pathogenic factors is well known. Salicylic acid also plays an active role in controlling transpiration, stomata closure, seed germination, fruit yield, glycolysis, flowering, heat generation and heat tolerance[18]. The role of salicylic acid in plant leaves and generative systems is well known and its highest level is found in tropical plants infected by necrotizing pathogens. The type of effect that salicylic acid had on plants provides a good indication that this substance increased the chlorophyll level, and as a result the degree of photosynthesis, carbohydrate reserve, cellular division and leaf surface[1,12,21,23].

CONCLUSION

Our findings demonstrate that salinity and salicylic acid had detrimental effect on Aloe vera characteristics as there were differences in biochemical characteristics and morphological characteristics on treats. Salicylic acids by synthesis some compounds maybe can induce effects by salinity stress.

Acknowledgments

The authors sincerely acknowledge for the partially helps provided by the Isfahan agriculture and natural resource researches center.

REFERENCES

- [1]. Abdollahi M, Jafarpour M, Zeinali H (2011) Effect of various Salicylic Acid concentrations on growth of Aloe vera L. *International Journal of Agri science*, 1(5): 311-313.
- [2]. Ashraf M, Harris PJC (2004). Potential biochemical indicators of salinity tolerance in., 166: 3-16.
- [3]. Cha-um S, Kirdmanee C(2009) Effect of salt stress on praline accumulation, photosynthetic ability and growth characters in two maize cultivars. *Journal of Botany*. 42: 78-98.
- [4]. El-Tayeb, M. A. 2005. Response of barley grains to the interactive effect of salinity and salicylic acid. *Plant Growth Regulation*. 45: 215-225.
- [5]. Fuentes V, Rodriguez N, Rodriguez C, Ramos R (1988). Salinity tolerance including Aloe arborescens and other species. *Agrotecnia Cuba*, 20: 1-6.
- [6]. Gunes A, Inal A, Alpaslan M, Eraslan F, Guneri E, Cicek N(2007) Salicylic acid induced changes on some physiological parameters symptomatic for oxidative stress and mineral nutrition in maize (*Zea mays* L.) grown under salinity. *Journal of Plant Physiology*. 164: 728-736.
- [7]. Hayat S, Ahmad A (2007) Salicylic acid: a plant hormone. 1st edition, Springer, Netherlands.

- [8].Horvath E, Szalai G, Janda, T (2007) Induction of abiotic stress tolerance by salicylic acid signaling. *Journal of Plant Growth Regulation* 26: 290-300.
- [9].Hu t. 2013. Stomatal and Metabolic Limitations to Photosynthesis Resulting from NaCl Stress in Perennial Ryegrass Genotypes Differing in Salt Tolerance .vol. 138 no. 5 350-357.
- [10].Joseph B, Justin R S (2010) Pharmacognostic and phytochemical properties of Aloe vera Linn. - An overview. *Int. J. of Pharma. Sci. Review and Res.* 2:106-110.
- [11].Moghbeli.E, Fathollahi,S, Salari,H,Ahmadi,G. Saliqehdar.F, Safari,A and Hosseini Grouh,M(2012). Effects of salinity stress on growth and yield of Aloe vera L. *Journal of Medicinal Plants Research* Vol. 6(16).3272-3277.
- [12].Misra N, Saxena P(2009) Effect of salicylic acid on proline metabolism in lentil grown under salinity stress. *Plant Science*, 177: 181-189.
- [13].Mukherjee A, Roychowdhury B (2008) the in vitro propagation of Aloe Vera sp. *TIG Research Journal* 1 : (2) 116-119.
- [14].Nobel PS, Berry WL(1985). Element and salinity responses of Agave species. California Univ., Los Angeles. U.S.A., *Am. J. Bot.*, 72: 686-694.
- [15].Olfati J-A, Moqbeli E, Fathollahi S, Estaji A (2012) Salinity stress effects changed during Aloe vera L. vegetative growth. *Journal of Stress Physiology & Biochemistry* .Vol. 8 Issue 2, p152-158. 7p.
- [16].Popova L, Ananieva V, Hristova V, Christov K, Georgieva K, Alexieva V. stoinova Zh(2003) Salicylic acid- and Methyl jasmonate- induced protection on photosynthesis to paraquat oxidative stress. *Bulg. J. plant physiol.* special issue. 133-152.
- [17].Rajeswari RM, Umadevi C, Sharmila R, Pushpa S, Selvavenkadesh K. P, Sampath K, Debjit B. (2012) Aloe vera: The Miracle Plant Its Medicinal and Traditional Uses in India. *Journal of Pharmacognosy and Photochemistry*. 1(4)185-221.
- [18].Raskin I (1992) Role of salicylic acid in plants. *Annu. Rev. Plant Physiology. Plant Mol. Biol.* 43: 439-463.
- [19].SAS Institute. (2001). SAS/STAT User's Guide for Personal Computer .Release 6.12 SAS Institute, Inc., Cary,N.C., USA.
- [20].Sahu P, Kumar N J , Shrivastava A(2011)Comparatives performance of Aloe vera and Aloe ferox spECies under pH along with desiccation stresses *International Journal of Drug Discovery and Herbal Research*, 1(1): 14-17.
- [21].Senaratna T., Touchell D., Bunn E., Dixon K., 2000. Acetyl salicylic acid (Aspirin) and Salicylic acid induces multiple stress tolerance in bean and tomato plant. *Plant Growth Regulation*, 30, 157–161.
- [22].Surjushe AR, Vasani R, Suple DG (2008) Aloe vera: A short review. *Indian J Dermatol* 53: 163-166.
- [23].Talebi.S, M. Jafarpour, A. Mohammadkhani, A. sadeghi (2012).The effect of different concentrations of salicylic acid and sodium chloride on Iranian Borage. *Intl J Agri Crop Sci.* Vol., 4 (18).
- [24].Tyler V (1993) *The honest herbal: A sensible guide to the use of herbs and related remedies.* 3rd ed. Binghamton, New York: Pharmaceutical Products Press.
- [25].Zan MJ, Chang HW, Zhao PL, Wei JG (2007). Physiological and ecological characters studies on Aloe vera under soil salinity and seawater irrigation. *Process Biochem.*, 42: 710–714.
- [26].Zhang S, Jie S, Wang H, Feng G(2010) Effect of salinity on seed germination, ion content and photosynthesis of cotyledons in halophytes or xerophyte growing in Central Asia. *Journal of Plant Ecology* Volume 3, Number 4, Pages 259–267
- [27].Zheng QS, Zhao-Pu LIU, You-Liang LIU, xingMing_EN (2004) Effects of iso-osmotic salt and water stresses on growth and ionic distribution in aloe seedlings. *Chinese Journal of Plant.* 28(6): 823-827.