



Scholars Research Library

Der Pharmacia Lettre, 2013, 5 (3):99-139
(<http://scholarsresearchlibrary.com/archive.html>)



Fullerene nanoparticles operating the apoptosis and cell proliferation processes in normal and malignant cells

¹Marina A. Orlova*, ¹Tatiana P. Trofimova and ²Oleg A. Shatalov

¹Department of Radiochemistry, Faculty of Chemistry, M. V. Lomonosov Moscow State University, Moscow, Russia

²Department of Pharmacology, School of Pharmacy, I. M. Sechenov Moscow State Medical University, Moscow, Russia

ABSTRACT

Fullerene derivatives superfamily attracts a serious attention of pharmacologists since some of these variable agents were proven to be not only drug delivery carriers but anti - cancer and immunomodulators as well. Most specifically, photodynamic therapies of malignant tumors are known for the fullerenes engagement. However, there is an obvious deficit of information on the cellular and molecular mechanisms of the fullerenes pharmacological effects which is a true obstacle on the way leading to practical medicinal use of the latters. Particularly, the mode of both direct (immediate) and distant side effects origin along with a fullerenes impact on necrosis, apoptosis and cell proliferation processes are no doubt needed to get far more clarified. It is hardly possible to exaggerate a significance of the fullerene nanoparticles functionalization type, their sizes and surface nanotopology for further promoting of either cytoprotective or cytotoxic effects. Noteworthy, the antioxidant properties of some water soluble fullerene derivatives were revealed while the fullerenes induced ROS formation might be also occurred. One of the most intriguing peculiarity of the fullerenes as pharmacophores consists in capabilities of some of them to intervene into the structure domains of functional proteins including enzymes and organelles linked receptors as well as to play a role of intercalators interacting with DNA double helix which, in turn, leads to a number of crucial consequences such as the biopolymer conformational flexibility shifts, catalytic activity changes, ligand docking affinity impacts in cell signaling pathways. Last not least, the fullerenes are about to compete with several natural metabolites and effectots which is itself a valid platform for pharmacological outreach. This Review deals with an Authors's original attempt to analyse the above mentioned points with an aim to elucidate those properties, methodological and structural, of numerous fullerene adducts that determine their apoptosis- and cell proliferation – modulating effects with a special respect to a target cell / tumor type.

Key words: C₆₀-based nanoparticles: cytostatics, antioxidants, immunomodulators, radioprotectors, free radical traps

INTRODUCTION

Nanotechnologies, that study particles of about 1-100 nm in size, open new possibilities for different fields of research and industry [1] with involvement of fields of pharmacy and medicine. Fullerenes as a structurally distinguished class of chemical compounds were discovered relatively recently (1985) and are of a particular interest among nanomaterials. Nanoparticles of 1-100 nm in size have various physical and chemical parameters, mechanisms and biomedical application. This is due to location of majority of their atoms at interphase of outer surface of the particles that, in turn, provides physicians and biologists with new quantum and mechanical mechanisms of their action. To imagine the size of nanoparticles let's take a height of a child of 4 years old to be equal to 1, than a diameter of the red blood cell will be in the range of 10⁻⁵-10⁻⁶, diameter of hepatitic C virus is ~10⁻⁷, carbon nanotube - ~10⁻⁸-10⁻⁹, and water molecule – 10⁻¹⁰ [2].

Fullerenes possess properties that make them attractive for various applied investigations. In particular, they may enter reactions of addition at double bonds (exoderivatives), introduction of atoms and clusters into the carbon backbone (endoderivatives) and are able to form heterofullerenes (metallofullerenes) and supramolecules. Due to this, they possess a whole spectrum of various effector properties useful for the human organism.

Buckminster fullerene (C_{60}) and its water-soluble derivatives, prepared by addition of various functional groups to the fullerene core, are suitable under different conditions for cytoprotection as well as antitumor therapy [3]. For example, some hydroxylated C_{60} and C_{60} -malonates behave as absorbers (acceptors, sponges) of radicals and protect cells from damages, caused by the oxidative stress [3-8]. However, they are also capable under higher concentrations and also, due to impact of UV-light, to cause a cell death through ROS (reactive oxygen species)-dependent and ROS-independent mechanisms [8-11]. In the case of some fullerene derivatives a cytotoxic effect regarding certain (in particular, cancer) cells has been described [12]. This may be purposefully used for tumor therapy without significant damage of a healthy tissue [13,14]. All fullerene derivatives bearing S-triazine fragment demonstrated a moderate to good level of inhibition of bacteria [15]. It is interesting, that the stable colloid solution of hydrated fullerenes forms a structured water layer around its carbone backbone that represents a special spheric cluster. Finally, a broad set of spheric clusters ($C_{60}@H_2O_n)_m$ [16] may be formed that leads to occurrence of particular properties mediated by water structure (Fig.1b).

Colloid solutions of C_{60} during interaction with water are able to form aggregates of different size and, respectively, of different toxicity. So, a like increasing of attractability of fullerenes for their usage in medical purposes, there is an uncertainty due to their toxicity and remote consequences of their application. Especially, it is pricked up when nanoparticles are used everywhere, for example, in cosmetics and other social branches. Derivatization of C_{60} decreasing of apparent toxicity may significantly influence on the mode of interaction of fullerenes with biological systems [17]. This increases the uncertainty in relation to the consequences for health and sometimes establishes an effect of "Trojan horse". So, it is of great importance to assess correctly risks and consequences of the fullerene-based nanotechnologies development. Debates regarding toxicity of nano C_{60} and hydroxylated fullerenes are described in detail in a review [18].

Over past several years the field of functional drug component delivery to definite organs and tissues in the organism (drug delivery) has acquired a special interest in the medical research. The fullerene derivatives, especially, porphyrin and supramolecular ones, are capable to act as a transporter as well as a drug. The latter is provided by the fact that fullerenes possess their own influence on the oxidative stress, proliferation, apoptosis, gene expression and other important functions of the organism.

A number of books that comprehensively describe an organic chemistry of fullerenes [19,20] and their application in biochemistry and medicine [21] were issued in 2005. Many reviews were also published [22-27] on the investigation of synthesis of new compounds, belonging to this class, and their possible medical usage [28-34]. Antimicrobial action of fullerenes and their derivatives was considered in the review [35]. Water-soluble fullerenes are of particular significance for medicine and their porphyrin derivatives [36,37] and fullerols (hydroxyfullerenes, fullerenols) [38] attract great attention.

Since main research fields on fullerenes are devoted to synthesis of new constructions, biochemical results have, in many cases, some stochastic character. In the present work we tried to concentrate our attention on the important aspects of fullerenes impact on the organism, including those linked with apoptosis and proliferation. These phenomena are especially important for cancer and neurodegenerative diseases prevention and therapeutics. We have also considered trends in intensively developing field of usage of fullerene nanoparticles on drug delivery and targeting. The question if fullerenes are useful for medical purposes or they are one more risk factor is a subject for debates in the scientific literature [39], and there is no certain answer for it. Also, the risks and success of nanomaterials application, as a whole, are discussed in detail [40].

It is necessary to take into account ecological and technogenic components of fullerenes impact, because fullerenes emerge in the environment from natural and anthropogenic sources such as volcano eruptions, forest fires and burning of carbon materials [41]. Moreover, in the last time production and usage of industrial fullerenes in different branches of industry are greatly increased. All these facts increase importance of comprehensive assessment of their toxicity for the environment and human health. Recent data [42] on risk assessment demonstrate a possible non-toxic concentration of fullerenes with average pore size of 96 nm that equal to 0.39 mg/m^3 . Calculations were made on the basis of trials results on rat lungs. However, this is not the last result in this research field.

I. FULLERENES AND OXIDATIVE STRESS.

One of the main factors of the impact of environment and effectors on apoptosis is an oxidative stress that is expressed in emergence of ROS. The latter includes singlet oxygen, superoxide radical, hydrogen peroxide and hydroxyl radicals, which arise in catalyzed by metal ions Fenton and Haber-Weiss reactions. ROS participate in initiation of receptor as well as non-receptor mechanisms of apoptosis, autophagy, lipid peroxidation and indirectly influence on many signaling pathways [43,44].

It is known, that the main impact of nanoparticles, as a whole, and fullerenes, in particular, is concluded in the oxidative stress that emerges in the organism due to their appearance in it [45]. However, definite consequences greatly depend on physicochemical properties of nanoparticles: size, shape, degree of dispersion in the solution, solubility in water, side chain composition etc. Their action is in the range from toxic to protective one. Concentration dependence is one of the most important indicators that determine consequences of the fullerene introduction. Thus, the most obvious effector impacts of fullerenes on proliferative processes are linked with their "relationships" with ROS, and the results of which may be directly opposite ones.

Due to its unique spheric structure, C₆₀ has a possibility to accept up to 6 electrons [46]. These electrons move quickly around the structure (fullerene grid) due to dipole moments. When C₆₀ is subjected to action of light, then an electron moves to more high energetic level, thereby producing an excited singlet C₆₀ that reacts with O₂ to form a singlet oxygen (¹O₂) [20].

Fullerenes are exclusively effective generators of the singlet oxygen with quantum yield of ^SO₂ close to one. They absorb greatly in UV- and slightly – in visible spectrum [24] that allows their usage in photodynamic therapy (PDT, chapter VIII). Originally formed the singlet excited state (^SC₆₀^{*}) undergoes intercombination transition into a triplet state (^TC₆₀^{*}). The triplet excited state is an excellent acceptor of electrons and potential producer of the superoxide anion radical O₂^{•-} (Fig.2). This pathway, in contrast to singlet oxygen generation, usually is observed in organic solvents, predominantly, in polar ones, especially, in the presence of reducers such as NADH [47].

Increasing number of functional groups, added to fullerene, leads to decreasing of quantum yield of the singlet oxygen. Toxicity is significantly decreased or even is fully absent in hydrophilic fullerenes [48]. Fullerene derivatives are able to control exogenic as well as endogenic ROS [49].

It is necessary to distinguish a pure fullerene C₆₀ and so-called nanoC₆₀ or n-C₆₀ that is an aggregate with indefinite (but, sometimes, stable) composition and poorly predicted size due to aggregation. Size of aggregates depends on preparation conditions. Nano-C₆₀ is formed often in the presence of a solubilizing agent, and a part of the latter, as suggested, is introduced into such nanoparticles [50,51]. However, in some cases, a spontaneous transition of C₆₀ into n-C₆₀ was described [7,52]. It is considered that "pure" C₆₀ does not exhibit visible toxicity that, however, depends on its concentration. In the same time, n-C₆₀ is, as a rule, a cytotoxic agent [24], which power depends on the size of aggregates and type of solution.

It is known [53] that the ROS-produced cytotoxicity is decreased under increasing of fullerene framework functionalization. However, there are suggestions that cytotoxicity of n-C₆₀ is not an internal property of C₆₀, but a property of a residual solvent, for example, tetrahydrofuran (THF) [7], intercalating into the grid. Suspensions, prepared during long stirring in water only, in some cases, did not demonstrate acute toxicity [54]. Suggested cytotoxic action of the residual THF [55,56] have not been confirmed yet. Authors [57] consider that water suspensions, prepared in the absence of organic solvents, not only have no acute or slightly acute toxicity, but must be protectors. Thus, the discussion about the reasons for the aggregates toxicity: contribution of solubilizers remainders or newly acquired properties of C₆₀ due to changes of the size and morphology because of the aggregation (in particular, due to a possibility or impossibility of a direct interaction with proteins), is still actual. Most likely, both variants are probable however, the second one is more sufficient under correct execution of an experiment.

C₆₀ toxicity was evaluated on different cells [5,58]. No cytotoxicity was observed in alveolar macrophages at 1.41-226 µg/cm² of C₆₀ [59]. In rat lungs there was no effect under inhalation of C₆₀ in concentration up to 3 µg/kg. [60] and particles of nC₆₀ of 55 nm in size at concentration 2.22 mg/m³ induced minimum alterations [61]. All this evidence that C₆₀-cytotoxicity depends, in addition to other factors (procedures for preparation and concentration), on the cell type. For example, thin layers of pure C₆₀ may even act as a suitable substrate for formation of cell colonies that was demonstrated [25] on human osteoblast-like MG-63 cells.

Stable and homogeneous C₆₀-dispersion medium did not influence on survival and did not cause apoptosis nor necrosis in human keratinocytes (HaCaT). The same picture was observed for A549 human lung cancer cells [62]. However, in both cases the cell proliferation was hampered.

Antioxidant properties of fullerene derivatives are determined by affinity of C₆₀ to electrons capture. This indicator is high for tris-malonyl-C₆₀ (C3) that removes superoxide radical in various cell lines at the level of active metal-containing mimetic of superoxide dismutase (SOD). DFT (density functional theory)-calculations showed [63] that, after superoxide radical contacting with C3 surface, its unpaired electron moves onto fullerene. In the result the aggressive O₂⁻ is transformed into neutral O₂. After that another superoxide radical reacts with C3-radical to form hydrogen peroxide, involving the electron transferred at the first stage. This process is an exothermic one with low energy of activation.

To modulate nC₆₀ toxicity, mainly, caused by ROS-production, and improve possibilities for its medical usage, fullerenes undergo different treatments to become soluble or pseudo-soluble in water. This process, besides other properties, influences on the ability to aggregation and depends on the type of side radicals. There are different structures for derivatives of fullerene: rods, bubbles, balls, membranes and linear constructions [64,65]. Morphology of C₆₀ derivatives aggregation, mainly, linked with hydrophobic interactions and hydrogen bonds. Few hydrophilic components are sufficient to prevent strong hydrophobic three-dimensional interactions between fullerene fragments and decrease their tendency for formation of aggregates. Differences in the affinity to electron and physical properties (degree of aggregation) influence greatly on biological and biomedical activity of derivatized fullerenes [66]. Supramolecular properties may promote affinity to radicals.

To prepare soluble samples of C₆₀, methods that lead to different results in geno- a cyto-toxicity are used. At present time, 4 approaches are offered to provide fullerenes with functions of solubility: hydro- and amphi-phility [24]:

1 – Chemical modification of fullerene framework by addition of different hydrophilic functional groups (derivatization or functionalization of fullerene) [12,67,68].

2 – Introduction of fullerene into water-soluble supramolecular structures with help of SAS (for example, polyvinylpyrrolidone - PVP [69]), calixarenes or cyclodextrins (CD) [70-72], when the fullerene nucleus is fully coated by modifier and has no contact with water. It is necessary to consider that cyclodextrins themselves tend to form aggregates in solution, linked with each other by hydrogen bonds network. CD-fullerene-conjugates demonstrated independent assembling of amphiphilic conjugates into spheric micelles in water [73] that, however, does not interfere capture of OH-radical by C₆₀-fragments.

3. – Method of exchange of solvents (MES) [74,75] uses volatile mixing with water solvents that dissolves fullerene. After water addition the solvent is evaporated and nC₆₀ suspension is formed [76]. The suspension is also prepared with use of ultrasound to promote fullerene transfer from non-polar solvents into water [77], by dispersing in water with Tween-80 [57,78], sugar syrup polyoxyethylene-hydrogenized castor bean oil [79]. Initial concentrations of C₆₀, in these cases, were less than 100 µg/ml [74,75]. Suspensions of nanosize dimension with large concentrations were prepared [78] by combination of SAS and disruption by ball grinder.

4. – Long (more than 2 weeks) stirring of pure C₆₀ with water. However, large aggregates may be formed and concentration of fullerene may be low [54,77].

Effector action of C₆₀, suspended in the solution, was determined on *E.coli*. [80] in dependence on degree of dispersing. It was considered that solubilizing solvents promote more high toxicity. 1% concentrations of solvents and 0.04% Tween-80 were used. According to these data, the most safe solvent in necessity of biocompatibility is N,N-dimethylformamide, but the solubility is not directly linked with cytotoxicity. Moreover, ozonation of nC₆₀ obviously and greatly increases degree of *E.coli* inactivation [81].

Modification of C₆₀ by hydrophilic component produces amphiphilic molecules that dissolve C₆₀ in H₂O. In that, a question is under discussion, is it possible to consider the amphiphilic molecules, prepared during modification of C₆₀, as really water-soluble ones? High level of solubility in water – 34 mg/ml at pH 7.4 – was obtained [82,83] with dendrite derivatives of fullerene with 18 carboxyl groups.

ROS-mediated cytotoxicity of different nC₆₀ suspensions (prepared by MES) was studied [84] with use of THF (THF/nC₆₀), ethanol (EtOH/nC₆₀) and stirring in water (aqu/nC₆₀). According to the ability to generate ROS, they were arranged in the next sequence: THF/nC₆₀ > EtOH/nC₆₀ > aqu/nC₆₀. Mathematical modeling of the singlet oxygen generation showed that the power of its quenching (THF/nC₆₀ < EtOH/nC₆₀ < aqu/nC₆₀) by solvents, intercalated into fullerene crystals, influences on their ability to produce ROS and causes damage of cells.

The oxidative stress induction, leading to mediated by extracellular signal-regulated kinase (ERK) cells death from necrosis, accompanying by damages of cell membranes without degradation of DNA [85], was observed at high concentration of nC_{60} (1 $\mu\text{g}/\text{ml}$), while studying the anti-glioma action of nC_{60} (MES in DMSO) on human cell line U251 and rat C6 [3]. A similar picture was observed in murine sarcoma cells. Autophagy and proliferation blocking in G2/M phase took place at low concentration (0.25 $\mu\text{g}/\text{ml}$ nC_{60}). C_{60} in concentration-dependent manner influences on differentiation of murine embryonic stem cells [86].

In case of endotracheal instillation and inhalation, C_{60} fine dispersed in water with 0.1% Tween-80 (33 nm, agglomerates up to 96 nm), in a dose of 0.1-1 mg and at different exposing duration [87] do not cause inflammation reaction (there was no significant elevation of neutrophils number). Clearance-kinetics of C_{60} after intratracheal instillation and inhalation showed that concentration of C_{60} in the liver and brain was lower than 8,9 ng/g tissue and in lungs the loading was reduced with time and depended on the initial concentration of C_{60} [88].

Nano- C_{60} (THF) are able to cause a lysis of human erythrocytes in a dose- and time-dependent manner [89]. nC_{60} (THF)-mediated hemolysis preceded the shrinkage of erythrocytes and increasing of cellular surface roughness. The hemolytic activity may be restored by use of N-acetyl-L-cysteine (NAC) that indicates the role of ROS in this process. In the meantime, C_{60} (CD-complex) and nC_{60} (so-polymer of ethylene vinylacetate - ethylene vinylversalate) were unable to perform the process.

Administration of C_{60} may alter the toxic action of other micro-impurities. Cells, co-exposed with C_{60} and As(III), had higher accumulation of As(III), thereby demonstrating the effect of "Trojan horse", because, due to fullerene, there was no elevation of cellular toxicity [90].

An opinion is often presented in the literature [91] that the observed difference of cytotoxicity of modified fullerenes is directly proportional to the ability of a fullerene derivative to induce ROS. However, the data analysis evidences that this dependence, probably, is more complex and multifunctional.

Nitrogen-containing fullerene compounds may possess greatly varied properties in dependence on the structure up to the significant toxicity. Addition of pyridines and pyrimidines to C_{60} fullerenes significantly increased their selective neurotropic activity, but increased 3-5 times the general toxicity in comparison to fullerenyls, $C_{60}F_{36}(NH_2)_{12}$. Fullerenyls were relatively low toxic ones and had no visible toxicological effects in doses not exceeding 800 mg/kg [85], and a tertiary water-soluble ammonia salt, fluor-fullerene-pyridinium fluoride $C_{60}[FNC_5H_5]+F^-$ already in a dose of 300 mg/kg demonstrated features of acute toxicity in relation to the brain and some parenchymatous organs. In the same time, the antioxidant action $C_{60}(ONO_2)_{7\pm 2}$ decreases consequences of ischemia, induced by damages of lungs [92]. Next derivatives of C_{60} such as N-methyl (2-quinolyl) fulleropyrrolidine and N - methyl (2-indolyl) fulleropyrrolidine have also demonstrated high antioxidant properties [93].

Hexa-sulfobutyl-derivatives of fullerene were used as radical sponges [94,95]. They inhibited lipid peroxidation and acted as anti-proliferative agents in atherosclerosis.

The next derivatives of fullerene such as fullerenes modified by different amino acids and also carboxy-fullerenes are under active investigation. Cystine- C_{60} derivative [96] showed a high activity of antioxidant action against superoxide and hydroxyl radicals ($IC_{50}=0.167$ and 0.008 mg/ml, correspondingly) and prevented H_2O_2 -induced apoptosis [97]. It is known [98] that 5 $\mu\text{g}/\text{ml}$ of cystine- C_{60} -derivative decreases apoptosis of PC12 cells.

Due to hydrophobic interactions, many amino acid- C_{60} derivatives are assembled independently to form spherical aggregates. Cystine- C_{60} and arginine- C_{60} (in difference from alanine- C_{60}) may form further multilayer bubbles due to hydrogen bonds. Experiments demonstrate that the bubbles morphology significantly influences cytotoxic as well as protective effect of these compounds against H_2O_2 -induced apoptosis [99].

Amphiphilic derivatives, folic acid- C_{60} (FDD, Fig.1c) [100] and oxidized glutathione- C_{60} (OGED) [101], are also able to self-assembling into spherical aggregates. They penetrate cell membrane and, due to the antioxidant action, protect cells of rat pheochrom cells (PC12), treated by H_2O_2 [100], from cytotoxicity without demonstration of visible toxicity. In that, the size of FDD-aggregates was lower than those for cystine- C_{60} and arginine- C_{60} derivatives that caused a greater amount of FDD in cells [96,99] and significantly increased the anti-apoptosis activity. Moreover, OGFED is able to overcome blood brain barrier (BBB) [101].

C3 is able to remove superoxide radical with rate constant of $2 \cdot 10^6 \text{ mole}^{-1}\text{s}^{-1}$ that is approximately 100 times lower than action of SOD. Such a value is in the range, described for manganese-containing SOD mimetic [102]. It was shown that a reaction between C3 and superoxide radical has a catalytic mechanism. Taking into account that C3

localizes in mitochondria, it is possible to consider that C3 functionally substitutes Mn-SOD, acting as biologically active SOD-mimetic.

Significant differences between lipophilic and hydrophilic particles of (C₃/D₃-C₆₀) are observed under consideration of different variants of C₆₀ modification by residues of hexacarboxylic acid (C₃ or D₃ conformations) [103]. Antioxidant action of fullerenes against lipid peroxidation and membrane disruption caused by radicals that are produced in xanthine/xanthine oxidase and Fenton reactions was used as a model system. Lipophilic derivatives of C₆₀ demonstrated protective effect that was greater than those of natural antioxidant-vitamin E. Anti-apoptotic functions of such fullerene derivatives may be also independent from their anti-ROS-acceptor role [17]. So, tris-carboxy-C₆₀ is a strong inhibitor of apoptosis in human skin epithelial cells (HEK), blocking G₀/G₁ of the cell cycle and causing the cell senescence. In that, there is decreasing of the expression level of ubiquitin ligase HERC5, participating in the innate immune response to virus and bacterial infections. In these conditions, no changes in proliferation were observed in cells, treated by hexa-carboxy-C₆₀ and γ -CD-C₆₀.

It is known that 10 μ mole/l of carboxyfullerenes decreases apoptosis of PBMC cells [5] and 1.2 mmole/l of a derivative, protocatechic acid-C₆₀ [104], decreases apoptosis of PC12 cells. It is obvious that the level of concentration depends on the chemical structure of a derivative that determines its further behavior in the field of aggregation and binding with various sites of biological components (and then toxicity) and on specificity of a cell line. Obviously, the conformational component plays an important (probably, major) role in these processes.

Comparison of antioxidant properties of PEG (polyethyleneglycol)-C₆₀, PVP (polyvinylpyrrolidone) - C₆₀, CD-C₆₀, C₆₀, containing OH-groups (Fig.1a) and C₆₀-isostearic acid [105] in relation to human skin keratinocytes (HaCaT) was conducted. All these fullerenes exhibit a powerful antiradical acceptor potential. However, the influence of side radicals (of the modifier) was expressed in necessary concentration of preparations: PEG-C₆₀ - 200-2700 μ mole/l, PVP-C₆₀ - 400-2700 μ mole/l, OH-containing fullerenes - 9 μ mole/l, C₆₀-isostearic acid - 18 μ mole/l.

PVP-C₆₀ and poly(2-alkyl-2-oxazolin)-C₆₀ (POX-C₆₀) form a homopolymer and casual co-polymer forming nanocomplexes with the same antioxidant activity [106]. However, only POX-C₆₀-complex was effectively captured by catecholaminergic neurons and attenuated elevation of intraneuron superoxide, induced by stimulation of angiotensin II.

Fullerols inspire a particular interest due to the absence of visible toxicity that are declared by many authors and due to unique physico-chemical properties. Hydroxylated, water-soluble C₆₀ due to its antioxidant properties inhibits catabolic stress-induced production of matrix metalloproteinases MMP-1, MMP-3 and MMP-13, and also apoptosis and preliminary aging in human chondrocytes. This makes it possible to use it as a protective agent against osteoarthritis [107].

Hydroxylated fullerene (C₆₀HyFn) in a dose of 4 μ g/kg decreasing oxidative stress in rats that suffer from STZ (streptozotocin)-produced diabetes, prevented complications such as testicular dysfunction and disruption of spermatogenesis [108]. Usage of C₆₀HyFn did not cause toxicity and led to elevation of some important polyunsaturated fatty acids. Hydrated fullerenes in concentration of 30 nmole/L protected nervous system tissues, preventing loss of astrocytes [93], and C₆₀(OH)₁₆₋₁₈ in concentration up to 50 mg/L did not influence on *Danio rerio* [109]. The ability of C₆₀(OH)₃₆ 8H₂O to hamper the oxidative stress in adipocytes leads to inhibition of lipogenesis-linked activation of macrophages in the adipose tissue [110]. Under certain conditions such highly hydroxylated fullerenes may sufficiently decrease the fat accumulation, ROS generation and infiltration of macrophages.

Fullerol C₆₀(OH)₂₄ demonstrated an acceptor activity in relation to superoxide radical in xanthine/xanthine oxidase system [111]. It did not cause genotoxic effects and demonstrated cytoprotective properties in a wide range of concentrations. However, other data [93] demonstrated that this fullerol of 7 nm in size and concentration of 1-100 μ mole/mL causes morphological alterations in cells of endothelium vessels, damages of cytosole (changing the level of lactate dehydrogenase) and inhibition of cell growth. This is accompanied by accumulation of polyubiquitinated proteins that activates autophagy. C₆₀(OH)₂₄ showed *in vivo* and *in vitro* a cytotoxic effect in relation to HepG2 cell line under doxorubicin (DOX)-induced hepatotoxicity. However, under even very strong oxidative stress cytotoxic effects of the fullerol were overcome by its protective role as a strong antioxidant [112].

Another fullerol C₆₀(OH)₂₂, in the range of nanomolar concentrations caused inhibition of a cell growth. The effect depended on the definite cell line, dose and time. In the same time C₆₀(OH)₂₂ as well as C₆₀(OH)₂₄ significantly suppressed the DOX-induced cytotoxicity at any concentrations independently from the time of fullerol addition. It is considered that these properties of C₆₀(OH)₂₂ are provided by its high acceptor activity in relation, namely, to OH-

radicals [113]. Fullerenes are cytotoxic for pigment epithelial cells of human retina in concentrations of more than 10 $\mu\text{mole/L}$ [114].

A gel, containing 75 $\mu\text{mole/L}$ of C_{60} , was utilized for treatment of acne as a receptor of radicals, which formation is a key moment in the disease development. As an antioxidant, fullerenes inhibited development of acne [115], seemingly, due to decreasing of neutrophilic infiltration.

Interestingly, fullerenes, dissolved with help of 1% Tween-80 (0.1 mL/10g of weight), were an effective preparation against action of polynarcotics. A lethal effect was observed under simultaneous action of methamphetamine and morphine due to increase of poly(ADP-ribose) polymerase (PARP) immunoreactivity of cells that both narcotics induce. These events are sufficiently linked with growth of hydroxyl radical number and they are sufficiently decreased by preliminary treatment by fullerene [116]. Moreover, this effect was stronger than those in the case of mepacrine (inhibitor of phospholipase A2) usage.

II. FULLERENES AND BRAIN CELLS

Fullerene and its derivatives, for example, FDD, OGD [101] and some other water-soluble fullerenes [117] are able to penetrate through the BBB and influence on the brain cells proliferation, in particular, through the oxidative stress [118].

As an example, *in vitro* modeling of interaction with BBB cells under oxidative stress, induced by H_2O_2 , was demonstrated for nanoparticles of water-soluble carboxyfullerene $\text{C}_{60}(\text{C}(\text{COOH})_2)_2$ [119]. As a model, brain cells of the endothelium microvessels (CMECs) were used because apoptosis of CMECs under action of ROS plays a key role in BBB dysfunction. It was revealed [120] that particles of $\text{C}_{60}(\text{C}(\text{COOH})_2)_2$ selectively and predominantly penetrate into oxidized (not normal) CMECs, keeping their integrity at the expense of H_2O_2 -induced F-actin depolymerization suppression.

Intraperitoneal and intravenous introduction of the suspension (0.3 mg C_{60}/kg) of C_{60} -PVP [23], less than 10 nm in size, inhibited differentiation of embryonic stem cells and brain cells that caused morphological alterations in murine embryos and correlated with increasing of cytotoxicity. In that, disruption of yolk sac function and embryonic morphogenesis were observed. Because proliferation was restored in addition of antioxidant enzymes, then the relationship between functional changes and effects of ROS is obvious.

After injection into hippocampus C_{60} -PVP [121] prevents disorders in the development of long-term memory, caused by a protein biosynthesis inhibitor- glutarimide antibiotics cycloheximide. The mechanism of inhibition is concluded in neurons apoptosis due to infringement of the protein synthesis chain that disrupts storage of information. Action of the fullerene complex is based on the antioxidant property, however, it is suggested [122] that absorption features of C_{60} may play a significant role in the prevention of neuronal apoptosis.

A derivative, malonic acid- C_{60} , promotes protection of dopamine in neurons of mesencephalon from toxin-induced degeneration [123]. Carboxyfullerenes demonstrated *in vivo* their efficiency in the prevention of neurodegeneration in the case of amyotrophic lateral sclerosis [82]. They also decrease the oxidative stress caused by iron in dopaminergic nigrostriatal way [124].

C_{60} -methionine derivative [125] (FMD) protected human neuroblastoma cells (SH-SY5Y) against neurotoxicity of a lead (in pretreatment with the 500 $\mu\text{mole/L}$ lead for 72 h.), which is realized through the oxidative stress. The protective effect was expressed in increasing of cells survival, anti-apoptotic action, elevation of a reduced glutathione (GSH) level and decreasing of a malonyl dialdehyde (MDA) level. The effect was higher for FMD in comparison with other amino acid- C_{60} derivatives, in particular, for β -alanine- C_{60} and cystine- C_{60} that, seemingly, is linked with specific properties of an amino acid. A hexacarboxylic acid- C_{60} derivative inhibits excitotoxic death of brain cortex neurons [4].

Water-soluble fullerols are considered to be a main achievement of nanotechnologies for neuromedicine [126]. They inhibit activity of glutamate receptors (GluR) from 10 to 80% under concentrations from 10 to 100 $\mu\text{mole/L}$. The most sensitive one was AMPA (2-amino-3-(3-hydroxy-5-methylisoxazol-4-yl) methylisopropionic acid) -receptor (up to 90% inhibition in concentration of fullerol of 100 $\mu\text{mole/L}$), then KA (kainate) and NMDA (N-methyl-D-aspartate)-receptors are followed, where 73% and 54% of activity, correspondingly, are inhibited [127].

Treatment of SK-N-MC cells by a single non-toxic concentration of water-soluble C_{60} -bis-adduct (Fig.3) caused modulation of adenosine receptors expression. Elevation of expression of A2A and A2B receptors mRNA and elevation of A1 and A2A protein level was observed [88], but no influence on the cells viability was revealed.

Fullerene derivatives have been studied regarding their possible usage in therapy of Alzheimer's disease (AD). To compare antioxidant activity in relation to β -amyloid ($A\beta$)-induced cytotoxicity, the effect of PEG- C_{60} nanoparticles and its pentoxifyllin carrying hybrid (PTX- C_{60}) on neuro-2A cells was studied [128]. $A\beta$ -generated free radicals may alter a structure of the lipid membrane in order to modulate neuronal systems of signal transduction. It is known that autophagy is induced by ER (endoplasmatic reticulum)-stress that is stimulated by ROS, level of cytosole Ca^{2+} , hypoxia etc. [129] and is linked with activity of AMP-activated cellular protein kinase (AMPK) [130]. All these processes are included into $A\beta$ -induced reactions in neuro-2A-cells [131]. C_{60} in a dose of 7.2 nmole inhibits process of fibril formation interfering formation of amyloid fibrils by $A\beta_{25-35}$ peptide [132]. Simultaneously, a cognitive process in rats is improved. Action of PTX in this case is included in activation of AMP-pathway that leads to decreasing of ID4 (inhibitor of differentiation 4) genes expression in cell lines, derived from astrocytes, and is expressed in decrease of cellular apoptosis. Cytoprotective effect of fullerene derivatives is linked with inhibition of autophagy [128], caused by activation of AMPK. Namely, hybrid molecule was more effective. Detailed evaluation of therapeutics possibilities of fullerene derivatives for the treatment of AD was performed in [133].

It is not possible to forget that a like neuroprotective functions, neurotoxic activities of fullerenes are possible and their manifestations have not been fully studied yet. Systematic introduction of carboxyfullerenes (10 mg/kg) did not prevent neurotoxicity, caused by known neurotoxin – 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine (MPTP) [134]. Moreover, even a dose-dependent increasing of striaric dopamine exhaustion was observed. In a dose of 30 mg/kg carboxyfullerene promoted death of MPTP-treated mice. However, during local application the same carboxyfullerene may act as neuroprotector agent [134].

Toxicity of fullerenes for brain cells is linked with oxidative stress, lipid peroxidation and glutathione depletion [53]. These processes participate in dopamine-mediated cell death in case of Parkinson disease (PD). Moreover, it is necessary to consider that transfer of β -amyloid-binding compounds through BBB is elevated due to of drugs encapsulation in polymer nanoparticles [135,136].

III. INFLUENCE OF FULLERENES ON SIGNALING PATHWAYS INVOLVED IN APOPTOSIS

Research in the field of fullerene derivatives is mostly directed to expand classes of their derivatives, while the mechanism and depth of impact on proliferation processes as well as relationship with action of other pro- and anti-apoptotic factors have not been revealed. As a result, there is no certain prognostic model to predict which derivatives of fullerene and under which conditions will exhibit anti-apoptotic properties and which derivatives may be considered as probable antitumor agents. Moreover, pro- or anti-apoptotic effects of fullerenes and their derivatives are mediated through different signaling pathways that may intersect. So, the grouping given below is conditional, but it helps to stress some relationships.

It is known [137] that dendrite single-substituted fullerenes with great number of OH-groups in the substitute and also C3-carboxyfullerenes with mirror reflection, pentaC3-exo and pentaC3-endo position are inhibitors of cell death. However, there is a question is this justified for all cell types and to what degree is it universal?

Ionic homeostasis

It was shown that water-soluble fullerenes influence ionic homeostasis [138]. Some derivatives of C_{60} demonstrated properties of blockators of ionic channels in biomembranes with exclusion of CIC3 channels [139]. Inhibition is reversible and, as suggested, interaction proceeds in extracellular domains of a channel. To assess a possibility of nanoparticles penetration through a membrane, Gibbs energy of C_{60} transfer along axis perpendicular to the membrane bilayer surface was calculated [140] that demonstrated the possibility of C_{60} penetration in principle. It is considered [141] that C_{60} possesses by affinity to transmembrane domain of K^+ -ionic channels and K^+ -ionic flow may be inhibited by addition hydrophobic medicinal preparations, causing a cell death.

Porphyrinfullerenes [142] are able to penetrate through lymphocytes cell membrane and BBB. As whole, water-soluble derivatives fullerenes readily penetrate through biological membranes of different types of cells [143].

Bcl-2 family

Expression of proteins of Bcl-2 family (Bcl-2 and Bcl-xl) was increased after pretreatment of cells by C_{60} - (glucosamine)₆. As a result, decreasing of oxidative stress and apoptosis, caused by kidney ischemia/perfusion and also decreasing of superoxide radical generation and improvement of kidney hemodynamics took place [144].

It is necessary to note that *ced-3*, *ced-4*, *ced-9* genes, responsible in *Caenorhabditis elegans* for analogs of Bcl family proteins, are subjected to modulation in consumption, as a food, of $C_{60}(OH)_{19-24}$ with average size of nanoparticles equal to 4.7 and 40.1 nm [145]. Activation of apoptosis is observed in the result that cause decreasing of animal lifetime. At high doses of nanoparticles (100 μ g/mL) a prompt reduction of viability takes place.

Mitogen-activated protein kinases (MAPK)

MAPKinases perform a control of cell division and are critical regulators of transcription [146], interacting with some signaling pathways in apoptosis [43].

There are indications in some works that the toxic action of nC₆₀ is similar to effect of single wall carbon nanotubes [147] and may be linked with activation of MAPKinases that, in turn, interrelated with activation of the transcription factor NF-κB. The process was studied on keratinocytes and it depends on a dose of a preparation [148].

Water-soluble fullerenes, on the contrary, were studied *in vitro* and *in vivo* [149] as protective agents against induction of osteoclasts genesis through RANK (receptor activator NF-κB)-signaling pathway and destruction of bone osteoclasts under arthritis. Fullerenes are able to down-regulate the RANK-induced differentiation of osteoclasts and suppress degradation of rat bone osteoclasts under intra-articular introduction of 20 μL of water-soluble C₆₀ (C=1 μmole/L).

C₆₀(C(COOH)₂)₂ nanoparticles inhibit apoptosis linked with JNK (c-Jun amino-terminal kinases)-signaling pathways [119], because they decrease JNK-phosphorylation, AP-1 (activator protein 1) and caspase-3 activation (Fig.4). In that, no influence on ERK activation was observed. There are data that the protective role of nC₆₀ may be mediated through activation of p38 [150,151], however, no contribution of nC₆₀-activation of JNK or p38 MAPK into cell necrosis was observed [3].

Anoikis

C₃-fullero-tris-methanedicarboxylic acids (C₃-tris-MDC) are able to cause resistance to such a form of apoptosis as anoikis (“homeless state”) in epithelial cells through mechanism of “trophic” effect with participation of actin of microfilaments [152].

Cytoskeleton damage

C₆₀(OH)_x was cytotoxic in millimolar concentrations in relation to renal proximal channel cells (LLC-PK1) [153], but there was no links with oxidative stress. Cells death, induced by fullerol, was caused by cytoskeleton damage and accumulation of autophagy vacuoles. Moreover, a loss of mitochondrial membrane potential and depletion of ATP was observed. Authors consider the cytoskeleton damage to be a primary cause.

Caspases and ionic composition changes

Caspase pathway of apoptosis is the most prevalent one, and modulations of multiple components of a biosystem may lead to activation of the caspase cascade [44]. However, these pathways were discussed in a very few works on fullerenes. More often separate factors able to cause activation of caspases are considered or activation of caspase-3 is established.

Effect of a substitution was considered in the case of apoptosis of human THP1 monocytes under action of free different fullerene derivatives: CD-C₆₀, hexa-adduct of carboxyfullerene C₆₀[C(CO₂H)₂]₆ (hexaC₆₀, T_h-symmetry) and tris-adduct of carboxyfullerene C₆₀[C(CO₂H)₂]₃ (trisC₆₀, C₃-symmetry) [154]. It was revealed that a substitute is especially important in characterization of compounds regarding apoptotic function. In the same time, dispersion characteristics are more important for redox-function. Preliminary treatment of cells by hexa-C₆₀ and tris-C₆₀ prevented valinomycin-induced apoptosis in difference from CD-C₆₀. In that, there was no dependence of apoptotic influence of preparations through cell ROS responses, caused by them. Nevertheless, it is necessary to consider the cell redox status, because some fullerenes are able, partly, by increasing expression of antioxidant proteins, protect cells from apoptosis, caused by oxidative stress. Thus, carboxylated fullerenes and encapsulated C₆₀ demonstrate opposite effects on pro-apoptotic caspase pathway [154].

Oxidative stress, mitochondrial depolarization and, as a result, activation of the caspase cascade were observed *in vitro* under action of nanoC₆₀, prepared by MES (THF), on the growth of B16 murine melanoma tumor cells [155].

On the contrary, nanoparticles of C₆₀(C(COOH)₂)₂ expressed an ability to inhibit cleavage of PARP and release of mitochondrial cytochrome C (Cyt C) that led to effective inhibition of the caspase-dependent apoptosis, caused by oxidative stress [119].

Transforming growth factor (TGF-β)

Derivatives of C₆₀-hexacarboxylic acid were described [156], as blocking agents of apoptosis TGF-β signaling in human hepatoma cells and there are differences in actions of regio-isomers on the same or the different cell types [156,157].

Lysosomal-mitochondrial pathway

Bis-adduct of fullerene with malonic acid, $C_{60}(C(COOH)_2)_2$, may inhibit apoptosis, initiated by TNF- α in HeLa cells by stabilization of lysosomes [158]. During internalization of $C_{60}(C(COOH)_2)_2$, expression of molecular chaperone Hsp70 is up-regulated and this promotes cells viability by inhibition of lysosomal membranes permeability (Fig.5). Acidic medium inside lysosomes is an important factor that makes significant, but temporal, influence on the aggregation state of nanoparticles. Size distribution varies, in average, from 100 to 10 nm and even to single molecules. Disaggregated particles may incorporate into lysosomal membranes, forming fullerene-containing mixed bilayer membranes. This stabilizes lysosomes and decrease permeability of lysosomal membranes, in particular, for destructing factors. Decrease of cathepsin release from lysosomes and inhibition of apoptosis, induced by TNF- α , takes place.

Mitochondrial deficiency

Cytotoxic effect of fullerols attracts a special attention. Studies [159], executed on rat hepatocytes, demonstrated that $C_{60}(OH)_{24}$ exhibits cytotoxicity, depending on drug concentration (0-0.25 mmole) and duration of exposition (0-3 hours), that is accompanied by formation of bubbles, loss of cellular ATP and GSH, elevation of MDA and oxidized glutathione (GSSG) levels. All this indicate the lipid peroxidation process. However, $C_{60}(OH)_{12}$ in concentration of 0.125 mmole demonstrated significantly lesser cytotoxicity at all parameters. It is considered that mitochondria represent the main target for fullerenes, so a mitochondrial deficiency, depolarization and inhibition of ATP synthesis are observed at an early stage. Lipid peroxidation in the result of oxidative stress is observed at more late stage. It is obvious that the toxic effect of fullerols depends on the hydroxyl group amount.

Pro-inflammatory cytokine

Derivatives of amino acid- C_{60} composition showed interesting concentration dependence in their effects on human epidermis keratinocytes. They in concentrations of 0.4 and 0.04 mg/mL caused decreasing of cells viability and initiated the pro-inflammatory reaction, but in concentrations less than 0.04 mg/mL, initiated loss of activity of cytokines and supported survival of cells [160].

Strong anti-inflammatory properties were demonstrated by fulleropyrrolidone-thalidomide- C_{60} and C_{60} -fulleropyrrolidine-xanthine dyads [161]. In the same time, fullerol $C_{60}(OH)_{24}$ influences negatively on endothelial cells, because it has pro-inflammatory and pro-apoptotic effect on HUVECs culture [162], and $C_{60}(OH)_x$ activated macrophages and promoted releasing of TNF- α [163].

A mixture of nanocrystal suspension of C_{60}/C_{70} , prepared by MES (THF), demonstrated an ability to modulate a TNF-induced oxidative stress and subsequent apoptosis [164].

Attempts to utilize fullerenes as modulator of action of known drugs are prospective. For example, a hybrid C_{60} -dexamethasone preserves an anti-inflammatory activity of dexamethasone, but has a sufficiently more lower immunosuppressive effect [165].

Dendrite cells play a key role in immune defense of the organism, in particular, in inhibition of tumor growth. $[Gd@C_{82}(OH)_{22}]_n$ may cause phenotypic maturation of dendrite cells. These nanoparticles are the powerful activator for dendrite cells as well as for Th1 immune response. The mechanism proceeds through stimulation of cytokines, including IL-12, p70; predominant increasing of IFN- γ , IL-1 β , and IL-2 production and up-regulation of co-stimulators CD80, CD83, CD86 [166].

NO-generation

Mechanochemically solubilized C_{60} is able to act as a cytoprotector against NO-induced cytotoxicity, but indirectly [167]. It seems that there is neutralization of superoxide radicals, produced by mitochondria, which interrupt a chain of peroxynitrite formation and caspases activation. However, it was shown [168] that, namely, the induction of peroxynitrite may be a critical event for genotoxicity of nano C_{60} . Immune system of mollusks in a great degree serves as a target for n C_{60} , dissolved in artificial seawater [169]. The process is mediated by NO release simultaneously with activation of p38.

A pristine fullerene, in difference from carbon nanotubes, does not stimulate release of NO [170] in murine macrophages that show very low consumption of nanoparticles.

Under action of n C_{60} on tumor cells in murine melanoma *in vivo*, the tumor growth was increased, despite accumulation of nanoparticles in cells and their cytotoxic effect [155]. The process was accompanied by elevation of production of immunoregulatory free NO radicals in splenocytes and reduction of splenocyte-mediated proliferative responses on T- and B-cell mitogens: concanavalin A and bacterial lipopolysaccharide.

To study protective activities of fullerene against NO-induced cytotoxicity (NO was released into solution by sodium nitroprusside, SNP), soluble fullerenes were used and they were modified by β -alanine, valine and folic acid, which possess a direct acceptor activity in relation to NO-radicals [171]. A contact of rat pheochromocytoma with 1 mmole/L SNP caused a notable decrease of the mitochondrial membrane potential, activity of SOD, catalase (CAT) and glutathione peroxidase (Gpx), decreased viability of cells and increased the level of intracellular accumulation of NO and MDA. Increasing of caspase-3 activity was also observed. Nevertheless, pretreatment of cells by amino acid- C_{60} -derivatives before their contact with SNP sufficiently inhibited all these NO-induced events. The degree of inhibition depended on the morphology of aggregation of fullerene nanoparticles (a visible toxicity was absent). Morphology of aggregates also influenced their protective effect against H_2O_2 -induced apoptosis [99].

Sulfonate derivatives of C_{60} increased NO content in the blood plasma after their intravenous administration [172]. In the same time, fullerols $C_{60}(OH)_{24}$ and $C_{60}(OH)_x$ demonstrated a direct NO quenching [111,163]. In the result, there was reducing of NO-induced decrease of CAT, glutathione-S-transferase (GST) and Gpx activities in the denudeated fraction of interstitial testicular cells of adult rats after SNP injection. It is suggested [160-224] that *in vivo* a competition between arginase and inducible NO-synthase (iNOS) for arginine as a substrate may take place. In the case of $C_{60}(OH)_x$ the elevation of arginase and suppression of NO was really observed.

It is interesting that a pure graphene is able to cause apoptosis through depletion of mitochondrial potential and elevation of intracellular ROS, activation of MAPKinases (JNK, ERK, p38) and TGF- β -signaling pathway, activating pro-apoptotic proteins of Bcl-2 family and also PARP and caspase-3 [173].

IV.EFFECTS OF FULLERENES ON PROTEINS AND DNA

Nanoparticles of modified fullerenes may potentially influence on proteins and biosystems, because their effects on cellular ionic homeostasis [174] and replication of viruses [138] are described. It is known a possibility of polyhydroxylated C_{60} [6,175], C_{60} -(glucosamine)₆ [176], carboxyfullerenes [4,5,152,156,177] and C_{60} -(cystine)₅ [98] derivatives to be involved in the relationships and reactions with proteins and biostructures that leads to modulation of apoptosis in different types of cells, including neurons.

Interacting with proteins, fullerene and its derivatives are able to alter their activity and properties [178,179]. An attempt was made to model interactions of C_{60} and a set of proteins that represent targets for drugs [180], in order of future developing of bioconjugate materials. Moreover, such modeling provides a possibility to obtain information about potential toxicity of conjugates as well as C_{60} -compounds. In modeling the data on antibacterial [181], neuroprotector [4,182] activity of fullerenes, their participation in DNA cleavage [183] and apoptosis [156], and also inhibition of amyloid structures [184] and ionic channels [139] were utilized.

Using docking [180], structures of C_{60} complexes with NOS, GST, and also with some proteins, in particular, β -secretase-1 and hypoxanthine phosphoribosyl transferase [185] were obtained. There are experimental data that fullerene derivatives participate in the inhibition of NOS [186], and glutathione reductase (Gsr) [187], and also of cysteine and serine proteases [67]. Anti-HIV (inhibitory) activity of fullerene derivatives [138,188,189], and also formation of fullerene-specific antibodies [190] are known.

It becomes evident on this basis that C_{60} -derivatives possess a possibility to readily interfere biochemical processes in the organism due to their binding with proteins and enzymes. From one side, it is important to learn how to use these possibilities in therapeutics purposes, predicting by docking the most favorable targets with help of fullerene derivatives. However, these processes may be dangerous for the organism, when unpredictably disrupting proliferation and interfering signaling pathways of apoptosis, which are not completely studied.

Interactions of fullerenes with BSA (bovine serum albumin) [191,192], HSA (human serum albumin, $K_{bind.}=1.2 \cdot 10^7$ mole⁻¹, Fig.6) [193] and lysozyme [194] are of particular importance. BSA and HSA execute a function of universal transporter of proteins and biologically active microelements (metals) in the organism [43].

A high affinity to HSA was demonstrated by organic phosphate-containing fullerols $C_{60}O_m(OH)_n(C(PO_3Et_2)_2)_L$, where $m \sim 8$, $n \sim 12$, $L \sim 1$ [196]. In that there was alteration of the tertiary structure of HSA in the direction of larger compactness together with elevation of content of α -helices and β -strands and increase of polarity of microenvironment of Trp residue. Hence, HSA may provide delivery of nanoparticles to organs and tissues. However, simultaneously a competition is established regarding other physiologically important components due to changes of conformational state of the molecule and a simple competition.

It is still known a little regarding real effects of fullerenes on structure and functions of proteins moreover, despite some proteins interact with fullerenes, no crystal structure of such complexes was obtained to the date. However, it

was demonstrated using docking [195] a high similarity between physico-chemical properties and geometry of the surface for sites of binding of fullerene with HIV-protease, BSA and HSA (Fig.6).

Carboxyfullerene forms a complex with β -lactoglobulin (a typical representative of a family of proteins of barrier fluid) [197], and a possibility of carboxyfullerene transfer from this complex to albumin was shown that expands our understanding of a mode of nanoparticles transport in a biological system.

During direct determination of mitochondrial Mg^{2+} -ATPase activity, it was revealed that the enzyme is inhibited by fullerol with $IC_{50} = 7,1 \pm 0,3 \mu\text{mole/L}$ [198] and this process is a concentration-dependent one.

Usage of water-soluble hexasulfonate derivatives of fullerene C_{60} decreased concentration of lactate dehydrogenase in the blood [172].

Fullerolpyrrolidines are non-competitive inhibitors in relation to acetylcholinesterase, IC_{50} varied in a range 15.6-31.4 $\mu\text{mole/L}$ [199].

Combined carboxyfullerene- C_{60} exhibited inhibitory activity against cysteine and serine proteinases. IC_{50} were equal: against calpain – 3.6 $\mu\text{mole/L}$; cathepsin – 10.5 $\mu\text{mole/L}$; papain – 43 $\mu\text{mole/L}$; trypsin – 5.6 $\mu\text{mole/L}$; plasmin – 3.2 $\mu\text{mole/L}$. Inhibition of thrombin was equal to 24% in concentration of preparation of 10 $\mu\text{mole/L}$ [67,180]. As inhibitors, such fullerenes may be used in treatment of burns and as radioprotectors, when a strong activation of proteinases takes place.

Water-soluble fullerenes that have, as a side radical, alcoholic, amine or amino acid ones act as inhibitors of Zn-containing enzyme, carboanhydrase (CAs) [200]. 13 isoforms of CAs showed different inhibition profile by these compounds. The mechanism is linked with corking of the entrance to the active center of the enzyme by the fullerene grid (the same size ~ 1 nm in both cases), while side fragments interact with amino acid residues of the active center, including His64, His94, His96, Val121 and Thr200.

C_{60} is a non-competitive inhibitor of GST in relation to etacrynic acid with $K_i = 48.8 \pm 0.25 \mu\text{mole/l}$ [201]. Under the same conditions, Gsr is effectively inhibited by only carboxyl derivatives of fullerene [187].

An uracil adduct of C_{60} was obtained [202] that is able to bind complementarily with adenine, adenosine and ATP. Also, porphyrin-fullerenes may exhibit various properties, for example, forming metallocomplexes. They are able to incorporate into proteins, for example, apo-myoglobin, with preparation of C_{60} -myoglobin [203].

DNA

One of the most known apoptosis pathways runs through DNA damages that activates p53-signaling pathway [204].

Water-soluble derivatives of C_{60} may sharply increase efficiency of polymerase reaction under relatively low DNA concentration [205] that is able to disturb a norm replication of DNA *in vivo* and initiate different signaling pathways of apoptosis.

Mono- and bis-methane phosphonates of C_{60} act in some cases as inhibitors of DNA- restriction endonucleases [15]. Inhibition of DNA-endonuclease *Exo III* by bis-methane phosphonate- C_{60} (BMPF) depends clearly on the fullerene dose and inhibition of PCR (polymerase chain reaction) with Taq DNA-polymerase ($IC_{50} = 2.7 \mu\text{mole/l}$) with help of BMPF was decreased with increasing of concentration of Taq polymerase in PCR system [206]. It is suggested that this process does not correlate with ROS and depends on the binding of the fullerene adduct with the enzyme and not with DNA.

Investigation of binding of bacterial DNA with nano C_{60} and fullerol showed [207] that at $\text{pH} < 9$ the fullerol size is suitable for binding with DNA. In this case elevation of thermal and enzymatic (in relation to DNAase I and Hind III restrictase) DNA stability in a dose-dependent manner takes place. Necessary for protection concentration of fullerol was equal to 50-500 $\text{ng}/\mu\text{L}$. In the same time, a concentration of not less than 500 $\text{ng}/\mu\text{L}$ was required to provide a protection from enzymatic cleavage during biding of DNA with nano C_{60} . Also dependence between effect of fullerene nanoparticles size on the result of impact on DNA and toxicity was observed. For example, nano C_{60} in size of 100 nm are unsuitable for binding with DNA.

DNA-fullerene binding is able to change a tertiary structure of DNA molecules and complicate their recognition by proteins because this process is based on the certain shape of the DNA molecule [208]. Seemingly, further it must affect the signaling pathway of apoptosis that involves p53.

Certain changes were demonstrated in [209] on the complex $C_{60}(OH)_{24}$ -DNA, where fullerol bound strongly to phosphates outside the native DNA chain and to nitrogen bases pairs in the main channel of DNA double helix.

It is known that $C_{60}(OH)_{24}$ only in concentration of $\sim 0.71 \mu\text{mole/L}$ begins to demonstrate an ability to inhibit DNA synthesis, while under lesser concentrations it inhibits proliferation. Thus, its effect is more cytostatic than cytotoxic. To understand various effects of fullerols and physical and chemical mechanism of their action, different components of fullerol were separated by isoelectric focusing [210]. Four fractions were identified with a general formulae $C_{60}(OH)_m(O)_n$, where m and n values may vary, according to their isoelectric point. Experiments, performed *in vitro*, showed that fullerols possess low toxicity. However, a fraction with the greatest density of the negative charge [$(-1.913 \pm 0.008) 10^{-4} \text{ C}$] and smaller size ($3.92 \pm 0.71 \text{ nm}$) causes DNA damage. In that, a free negative surface charge is determined by a degree of C@O functionalization. Hence, to obtain good biocompatibility, it is necessary to perform separation and purification of the raw fullerol.

Antivirus activity of fullerenes

Fullerene derivatives are able to be accommodated inside hydrophobic cavity of HIV-protease, thereby preventing access of substrates into catalytic center of the enzyme. Relatively long ago it was found [138] that a derivative fenetylaminosuccinate- C_{60} possess anti-HIV activity with $K_{\text{inh}}=5.3 \mu\text{mole/L}$. Dendrite derivatives of fullerene [83] showed EC_{50} equal to $0.22 \mu\text{mole/L}$ in lymphocytes of a patient with HIV-1 acute phase. To degrade selectively the HIV-1 protease, C_{60} -carbohydrate hybrid was used [211]. A docking was performed to analyze interactions of mono- and bis-adducts of fullerene C_{60} in order to investigate the binding of fullerene based inhibitors to the HIV-1 protease [212].

A number of works has shown [213,214] that prerequisites for antiviral activity in relation to HIV-1 and HIV-2 strains are, possibly, specific relative position of two substitutes and positive charges in the neighborhood of the carbon grid. This enhances attention of researchers to studying structure/activity relationships in C_{60} derivatives to reveal anti-HIV activity. It was found that only trans-2-isomers are prospective ones and presence of quaternary pyrrolidone nitrogen affect solubility [214]. Positive charges near C_{60} -framework may increase antivirus activity, while extension of solubilizing chains induces cytotoxicity. Trans-isomer of bis-fulleropyrrolidine- C_{60} showed interesting antivirus properties in relation to HIV, and no regiostereomers of this compound exhibited inhibitory activity in relation to DNA or RNA of viruses different from HIV.

A mechanism of HIV reversed transcriptase inhibition was registered for some fullerene derivatives [215]. However, at the moment it is hardly to talk about specificity of anti-HIV mechanism.

Recently due to development of conjugates synthesis, C_{60} -Curdlan (linear 1,3- β -glucan) sulfate [216] was obtained in order combine anti-HIV activity of both components [217].

Adduct of fullerene with polyvinylpyrrolidone possesses antivirus effect in relation to influenza A and B virus [218]. Under effective dose of $C_{60} \sim 5 \mu\text{g/mL}$, its effect is stable during all cycle of the virus amplification.

V. FULLERENES AND GENOTOXICITY

Oligonucleotide microchips were used identifying and profiling of gene expression in rat lungs after inhalation of C_{60} [219] and up-regulation of genes linked with inflammatory response, oxidative stress, apoptosis and metalloendonuclease activity was observed. Moreover, some genes, linked with immune system are subjected to up-regulation. In general, no severe pulmonary toxicity was observed. Fullerenes in lungs [220,221] are captured by alveolar macrophages, which, in such a manner, execute their protective role.

C_{60} in doses from 0 to $200 \mu\text{g/mL}$ and long exposition did not cause apoptosis of epithelial cells of murine lungs and mutations in *CII* gene, however, proliferation was slowed down (in G1 phase), DNA breaks are occurred, but amount of mutations was insufficient [222].

Comparative gene microchip analysis firstly showed on zebrafish that hydroxylated fullerenes causes de-regulation of circadian rhythm gene and also alter the regulation of genes, participating in activity of kinases, vesicular transport and immune response [223].

Emergence of genotoxic effect of fullerenes was registered at concentration of $2.24 \mu\text{g/mL}$ of medium in *Drosophila melanogaster* larvae [93] and human lymphocytes culture [74], where the strict concentration dependence was observed. It is suggested that the genotoxicity is a result of the lipid peroxidation induction.

Changes in the profile of expression of genes after intratracheal instillation with C₆₀ in various dosages were characterized. In particles dose of 1 mg the next genes: *Cxcl2*, *Cxcl6*, *Orm1* and *Spp1* participate in the «inflammatory reaction» and *Mmp7* and *Mmp12* genes were highly expressed for more than 6 months in matrix metalloprotease activity [220]. Expression of some genes (predominantly, «inflammatory») correlated with the dose of introduced drug that makes it possible to use them as biomarkers.

Mechanism of immunotoxicity was studied [224] on murine BAL cells. In instillation of n-C₆₀ elevation of pro-inflammatory cytokines such as IL-1, TNF- α and IL-6, and also Th1-cytokines (IL-12 and IFN- γ) in BAL-culture took place. Moreover, expression of MHC (major histocompatibility complex) of class II (H2-Eb1) molecules was stronger than those for MHC of class I (H2-T23). Increase of T-cell dissemination, cellular infiltration and damage of a tissue, linked with genes expression in the pulmonary tissue were also observed in the experimental period that indicates inflammatory response to appearance of nanoparticles. However, fine dispersed fullerene in size of 33 nm and in amount of 0.1-0.2 mg during intratracheal instillation did not cause increase of inflammatory neutrophils level that was observed in a small amount after introduction of 1 mg of nanoC₆₀ [87]. Also in these cases there was no significant elevation of expression of CINC-1, -2 $\alpha\beta$ and -3 (cytokine-induced neutrophil chemoattractant) in lungs.

According to [225] fullerene C₆₀ (water-dispersive) adversely affects on cell activity of nonspecific immunity link by inhibiting the enzymatic activity of myeloperoxidase and the expression of molecules CD54. Hence this is the impact of C₆₀ on the various phases and mechanisms of phagocytosis.

Modulating effect of fullerene in the combined molecules was demonstrated under usage of fullerene C₆₀ conjugates with immunomodulating peptide Thr-Lys-Pro-Arg (TA) in the form of NH₂-TA-C₆₀ (161 nm) and C₆₀-TA-COOH (132 nm) in concentration of 2-20 μ mole/L at murine model *in vitro*. In comparison with the peptide TA enhancement of phagocytosis, chemotaxis activity and expression of MHC of class II. Conjugates were resistant in relation to degradation by leucine aminopeptidase, did not exhibit toxicity in relation to macrophages and on all parameters are the candidates for vaccine adjuvant [226]. Immunoactivation was observed in the case of nanocomposites of fullerene C₆₀-PVP and C₆₀-Tween-80 (0.01-0.1% solutions) [227] that showed a brightly expressed anti-inflammatory effect.

Gene modulating effect of fullerols C₆₀(OH)_x (20 ppm) in comparison with other (non-carbon) nanoparticles was studied on human primary skin fibroblasts (HDF) [228]. The greatest changes of gene expression (in times in relation to control) were observed for *TLR3* (2.59), *TLR6* (1.84), *IL1RAP* (1.66), *IKBKB* (1.2), *NFKB2* (1.29), *CCL2* (0.62), *NLRC4* (0.52), however, smaller changes were visible regarding expression of other genes. Thus, fullerols greatly influence on cytokines balance and, seemingly, increasing immunity in these conditions. In the same time, there were phosphorylation of p38 and JNK (MAPK) that leads to NF- κ B activation and inflammatory response. This cascade is a result of ROS action and IL-6 up-regulation. It is obvious that the balance of cytokines may shift to any direction under changes of nanoparticles parameters or concentration.

Stable C₆₀ nanoparticles, prepared in the form of suspensions with help of 0.1% sodium carboxymethylcellulose or 0,1% water solution of Tween-80, did not demonstrate genotoxic ability in bacterial reversible mutation analysis, *in vitro* analysis of chromosomal aberrations and *in vivo* micro-nuclear analysis [78]. However, usage of fullerol C₆₀(OH)₂₄ indirectly in some cases caused decreasing of chromosomal aberrations frequency [229]. In the meantime, recent investigations showed that fullerols are perspective for inhibition of enzymes involved in processes of DNA replication, transcription etc., and leading to dysfunction of DNA with subsequent increasing of the mutation rate [207]. Authors suggest that the direct damage of DNA by fullerene derivatives is not great, but emerging ROS induce inflammatory processes and genetic damages [230]. In that, namely, peroxyinitrite represented a signal event for genotoxicity and emergence of mutations (deletions) in *gpt* delta transgenic mouse primary embryo fibroblasts [168].

It is necessary to remember that all these processes are dose- and size (morphology)-dependent ones.

VI. ANTITUMOR EFFICIENCY OF FULLERENE DERIVATIVES

Due to the ability to cause under definite conditions cell death, fullerenes are potential antitumor agents. However, predicting of pro- and antitumor properties of various derivatives of fullerenes, moreover, for different types of cells, is still a hard task. In this connection, computer approaches and DFT method for calculation of physico-chemical properties of fullerene derivatives are actively utilized. Calculations by method of DFT of stabilization of ortho-, meta- and para- substitutes in relation to aldehyde group in a series of 1-(4,5, and 6-selenyl-derivatives of 3-formyl-phenyl) pyrrolidine fullerene molecules C₆₀-C₂H₄N-[3-(CHO)C₆H₃SeX], where X= CN, CH₂CH(NH₂)CO₂H and CH₂CH₂CH(NH₂)CO₂H, in order to determine effect of such structural changes on electronic structure of fullerene

derivatives [231] determining their oxidative properties, may serve as an example. In case of selen-cyanide systems, general tendency that ortho-structure is more stable was kept. Similar picture is observed in the case of selen-methionine group. However, in the case of selen-cysteine the para-structure was the most low energy one, probably due to close position of a proximal NH₂-group to a point of interaction Se-O. Such fullerene systems may be effective molecular acceptors of free electrons in biological systems.

Some forms of fullerols demonstrated antitumor activity *in vivo* in relation of hepatocarcinoma H22 [163] and *in vitro* cytotoxicity against human liver carcinoma cells (HepG2) [91,154]. The effect depended on the dose and size of particles. It was shown through two lysosomal enzymatic markers (arginase and acid phosphatase of peritoneal macrophages, which were obtained from animals, treated *in vivo*) [163] that usage of fullerol C₆₀(OH)_x stimulates significant activation of macrophages and inhibits the tumor growth. Production of TNF-α by fullerol-stimulated macrophages is sufficiently increased under action of C₆₀(OH)_x in the dose range of 0.2-2 mg/kg. In that, no acute toxicity was observed at concentrations up to 5 mg/kg.

Daily doses of 0.08 and 0.4 mg/mL were used during investigation of fullerol C₆₀(OH)₂₀ for antitumor and antimetastatic activity [232] on EMT-6, metastatic model of breast cancer the average size of nanoparticles at pH 7 varied from 223 to 364 nm, correspondingly. In addition to expected modulation of oxidative stress, decreasing of expression of a few angiogenic factors in tumor tissues was observed. In particular, expression of CD-31 (PECAM-1 – molecule of adhesion of endothelial cells thrombocytes) and vascular density was decreased in tumors treated by fullerol. By this mode C₆₀(OH)₂₀ inhibits a tumor growth and suppresses metastasizing through signaling pathway linked with tyrosine kinase.

Bis-(N,N-dimethyl-pyrrolidin iodide)-C₆₀ [233] demonstrated cytotoxicity in relation to pro-myeloleukemic cells (HL-C60).

Endofullerenes are of special interest regarding cancer therapeutics. At present, antitumor effect of hydroxylated metallofullerene, containing gadolinium [Gd@C₈₂(OH)_x], which action is linked with accumulation of immune cells, was studied in detail [14]. However, presence of this compound in the tumor *in vitro* was not practically detected that suggests interrelationship of tumor growth inhibition and antitumor immunity activation [234]. It is known that up-regulated immune response *in vivo* is a favorable strategy for clinical treatment of cancer.

Gd@C₈₂(OH)₂₂ [235,236] of 2 nm in size and with mean size of its aggregates of 100 nm [14] possesses immunomodulating activity *in vivo* and *in vitro*. It may stimulate macrophages and T-cells to release of a few cytokines IL-2, IL-4, IL-5, TNF-α and IFN-γ that promote inhibition of tumor growth *in vivo*. Specific immunomodulating effects on T-cells and macrophages include polarization of cytokine balance to Th1 (T-helper cellular type 1) cytokines by decrease of Th2 cytokines production (IL-4, IL-5 and IL-6). Amount of Th1 cytokines (IL-2, IFN-γ and TNF-α) in serum samples are increased. Modulation depends on a drug dose. Under low concentrations nanoparticles of Gd@C₈₂(OH)₂₂ slightly influence activity of immune cells *in vitro*. In high concentrations they notably increase immune responses and stimulate release of cytokines that facilitates removal of abnormal cells. In particular, increasing of TNF-α level that plays key role in cellular immune processes is observed. Gd @ C₈₂(OH)₂₂-nanoparticles are more effective than some clinical antitumor preparations in inhibition of tumor growth in mice.

Many tumor cells (in particular, murine hepatoma H22 cells) have pro-coagulant activity that promotes local activation of the coagulation system. So, changes in blood coagulation were assessed after introduction of [Gd@C₈₂(OH)₂₂] to mice. Elevation of APTT (activated partial thromboplastin time) and decrease of PT (protrombin time) in comparison with control group was revealed. Moreover, an elevated concentration of fibrinogen was observed in the blood plasma after introduction of [Gd@C₈₂(OH)₂₂]-nanoparticles [234]. Thus, this drug decreased *in vivo* blood supply of the tumor [166] and simultaneously down-regulated more than 10 angiogenic factors at mRNA level that was further confirmed at the protein level [237]. In that, all indicators on activity of enzymes linked with oxidative stress: Gpx, SOD, GST; CAT – demonstrated tendency of approaching normal level. In the same time, reduction of MDA level took place. This indicates high antioxidant properties of Gd-containing fullerene [238]. Moreover, the preparation promotes activation of dendrite cells enhancing the immune response [166]. Nanoparticles of [Gd@C₈₂(OH)₂₂]_n are the first fullerene derivative promoting maturation of dendrite cells as well as antigen-specific immune response.

Experiments on transplantation of malignant tumors to group of mice demonstrated efficiency of C₆₀ usage to inhibit growth of such tumors [239]. This is linked not only with antioxidant activity of C₆₀, but also with blockage of specific cell receptors, in particular, endothelia growth factor.

A review [240] was devoted to possible use of C₆₀ and C₇₀-fullero-pyrrolidines that include antitumor as well as antioxidant drugs [27].

Water-soluble co-polymer fullerene-acrylamide demonstrated *in vitro* activity against bone marrow tumor cells [241].

As concerned biodistribution of metal-containing fullerols, then, after intravenous injection to mice, Gd@C₈₂(OH)₄₀ was accumulated, predominantly, in kidney, liver, lungs and spleen [242]. [Gd@C₈₂(OH)₂₂]-nanoparticles were accumulated in bone tissue, pancreas, kidney, spleen and liver and only very small amount was directed to lungs [234]. Fullerol, containing holmium, ¹⁶⁶Ho@C₈₂(OH)_y was distributed among different organs and tissues with exclusion of brain and adipocytes after 1 hour from its administration into the organism [117].

Using ^{99m}Tc, biodistribution of C₆₀(OH)_x(O)_y was studied [243]. The compound was fastly absorbed by tissues, especially by coronal bone, breastbone, spine, liver and spleen. Its clearance was slow in all tissues with exception of the brain.

In the last time, conjugates of fullerenes with known drugs for enhancement and/or deliverance of drugs became a widespread event. For example, the fullerene-doxorubicin was used [244]. However, uptake of conjugates, containing fullerenes, and their biodistribution often substantially differs from those for free molecules of the drugs that require a detailed investigation. For example, DOX is localized in a cell nucleus, but its conjugate with fullerene is in a cytoplasm.

Comparative study of nC₆₀ and nC₆₀(Nd) as agents that modify proliferation in HeLa cancer cells was performed [245]. Nanoparticles of nC₆₀(Nd) elevated chemotherapeutics susceptibility of cancer cells, and the mechanism of these processes included not only common for nC₆₀ ROS-attacks, but also enhancing of autophagy. Elevation of chemotherapeutic activity proceeded already at low dose of the preparation without significant cytotoxicity. It is suggested that, while entering cells, the nanoparticles are captured by autophagosomes, however, their inability to be degraded leads to block of the way after autophagosome/lysosome fusion due to autolysosomes accumulation. However, this needs an experimental confirmation.

VII. PORPHYRIN-FULLERENES AND TARGETED DELIVERY OF DRUGS

Over the last years a great attention is paid to using different fullerenes for drug transport. It is still relatively novel, although extensively developing field in the fullerene chemistry and their usage that needs new ideas and thorough investigations. Some contemporary examples of the drug delivery [246] and also a general strategy of nanoparticles utilization as transporters [34] have been described.

Concerning fullerenes, the most attractive conception is contained in the idea that there is a multiunit with properties of transporter and a few attached medicinal parts (for example, photosensibilizator) and also there is a lead unit possessing selectivity to target cells [247]. In another variant, an additional sensibilizing compound may be used as a transporter. In that, nanoparticles must be hydro- and lipophilic ones and be free of acute toxicity.

One of the prospective approaches in this field is the use of porphyrinfullerenes. Fullerenes are known as excellent electron acceptors due to their low energy of reorganization. A direct interaction of porphyrins and fullerene [248,249] and a strong interaction of bis-porphyrins with fullerenes in the solution are known. There are two methodologies for the system of electron transfer of photoexcited state of porphyrin and fullerene: covalent [250] and coordination [251] ones. Porphyrins, in turn, possess properties of receptors and are able to form complexes with metals cations yielding metalloporphyrins. A class of dendrite porphyrins, containing a central atom of zinc inside and 32 ammonia groups, was a strongest singlet oxygen inducing cytotoxic agent [252]. It was possible to form various supramolecular complexes of fullerenes and metalloporphyrins such as, for example, were shown on Fig.7a [37]. A constant of association of relative flexible bis-porphyrins with C₆₀ approximately is equal to 5 10³ mole⁻¹ (in toluene) [253] and is varied in dependence on the central metal ion. For example, in the case of Rh(III) the association constant is equal to 2.5 10⁷ mole⁻¹. In general, this value is greater for C₇₀ than for C₆₀ due to alteration of fullerene "ball" conformation [37] and associated differences in symmetry and a number of non-equivalent bonds [19,254]. Such supramolecules, in addition to their own modulating effects on cells, may serve as transporters for targeting delivery of fullerene conjugates (and fullerenes themselves) to biological targets in the organism. To perform a targeting transport of different drugs [85] a high electro negativity of fullerene molecules (C₆₀ and C₇₀) are exploited.

Moreover, porphyrinfullerenes are exclusively attractive carriers for targeting delivery of drugs to organs and tissues, because they under a certain design possess an elevated bioavailability and non-toxic degradation products.

Porphyrinfullerene derivative PMC16 (Fig.7b) [255] was used for delivery of magnetic nuclei of ^{25}Mg that may increase more than two times the ATP yield [256] (that is very important in conditions of hypoxia-induced acidosis). PMC16 decreased the acidosis consequences, arisen during chemotherapy, and did not exhibit toxicity in a wide range of concentrations. After introduction into the organism, PMC16 was preferably accumulated in the myocardium and in a less degree in lymphocytes and brain. For comparison: accumulation of nC_{60} is observed, mainly, in the liver and spleen and sufficiently lower in other organs [257]. Being a cation exchanger, PMC16 prolonged release of metal ions in acidosis conditions only. PMC16 was also used for delivery of zinc isotopes to lymphocytes of healthy donors and patients with ALL-T, ALL-B and AML [142] that led in case of ALL-B and AML to increasing of apoptosis of cancer cells in specific conditions.

Construction of supramolecules, where fullerene derivatives or their complexes are used as guest molecules, is a relatively new scientific field in the drug delivery. *In vitro* supramolecules demonstrated themselves as good thermo-reacting co-polymer of poly-(N – isopropylacrylamide)–CD as a host molecule and adamantyl- C_{60} as a guest molecule [259]. Due to fullerene, these molecules possess acceptor ability in relation to OH-radicals and effectively inhibited oxidative damages of mitochondria.

Cationic micelles of block-copolymer poly (N-vinylcaprolactam)- C_{60} delivered fullerene to cells for further utilization in PDT [260].

Amphiphilic derivatives of C_{70} , in rational way packed into lipid bilayers with loading up to 65%, represent a new system of drug delivery [261]. In particular, lyposomal malonyl-fullerenes- C_{70} (ALM) were used. ALM have been effective quenching agents for superoxide and hydroxyl radicals and inhibited lipid peroxidation, while maintaining an integrity of lipid bilayer structure.

Use of fullerols as drug transporters may bring positive results in cancer therapy. So, fullerol ($\text{C}_{60}(\text{OH})_8$)-DOX conjugates inhibited proliferation of cancer cell lines *in vitro* through blocking of G2/M stage of cell cycle [262]. Fullerols may serve as transporters of radioactive nuclides *in vivo*. $\text{C}_{60}(\text{OH})_{20}$, labeled by $^{99\text{m}}\text{Tc}(\text{CO})_3$ [263] as a complex was delivered quickly to all tissues, excepting the brain, and was kept their without loss of activity during 3 h. Rate of radioactivity clearance took place during 24 h. It is suggested to use in future a fullerol platform for cisplatin delivery.

To transport through ophthalmologic barrier, fullerols $\text{C}_{60}(\text{OH})_{22-26}$ are used, however, it is necessary to remember that water-soluble fullerenes demonstrated cytotoxicity in relation to epithelial cells of human retina (HLE-B-3) [264,265]. However, $\text{C}_{60}(\text{OH})_{22-24}$ -nanoparticles are considered as transporters of photosensibilizers to eye (and skin) cells in PDT of tumors [265-267].

VIII. PHOTOTOXICITY OF FULLERENES. PHOTODYNAMIC THERAPY

During light absorption fullerenes may actively generate ROS that make them possible agent in PDT. Depending on functional groups introduced into the molecule, fullerenes may effectively perform photoinactivation of pathogenic microbial cells and malignant cancer cells. However, under certain conditions many fullerenes are able to protect cells from UV and other irradiation. To determine a role of fullerenes in PDT, physical and chemical properties of a certain derivative as well as experimental conditions: duration and intensity of irradiation, concentration of a preparation, size of nanoparticles, aggregation are of great meaning.

Large amount of model investigations of fullerene phototoxicity are executed in frames of ecological investigations on some fish and invertebrates species in model water systems at simultaneous environmental level of UV. It is necessary to take into account the presence of organic substances, in particular, polycyclic aromatic hydrocarbons (PAHs), which bioavailability may altered by fullerenes [268]. For example, under short-term expositions C_{60} protects cell components (mitochondria, microvilli, basal inclusions) from UV+fluoranthene (model PAHs) phototoxicity. However, long-term effects (21 days) even at low concentrations of C_{60} lead to cells damage in gastrointestinal tract of *Daphnia Magna*. This demonstrates an exclusive importance of interactions between stress agents and water nanoparticles in water medium. Such synergism must be accounted in medical usage of fullerenes.

The method of PDT is widely used in cancer treatment (for example, zinc-containing photosensibilizers are well-known [43]). Two types of effects are distinguished in PDT [269], where acting agents are preferably superoxide radicals or singlet oxygen (Fig.2). For fullerenes the method is based on the selective accumulation of photosensibilizer in tumor cells and its ability to generate both agents (singlet oxygen as well as superoxide radical) under irradiation with light beam of definite wavelength. The ability of fullerene to generate in these conditions ROS with rate of $10 \text{ nmole mL}^{-1} \text{ min}^{-1}$ was confirmed by EPR method [270].

In the present time, photosensibilizers on the basis of carbon nanostructures did not pass clinical trials and are not allowed for their practical use [271]. Nevertheless, investigations are widely carried out with hope for their further usage.

Incubation of tumor cells with fullerenes with subsequent irradiation by light causes apoptosis in different types of cancer through 4-6 hours after irradiation [272]. Mono-adducts of C₆₀ (in comparison with bis- or tris-adducts) demonstrated the greatest activity in relation to cancer cells due to their high cell uptake and enhanced localization in mitochondria.

C₆₀ with a few polyester chains are also prefer to be accumulated in cancer cells and under irradiation at $\lambda=400-505$ nm cause a tumor necrosis [273].

The next derivatives phenylalanine-C₆₀, folic acid-C₆₀, L-arginine-C₆₀ in concentration of 5 $\mu\text{g/mL}$ did not cause cytotoxicity in the dark during long time and possessed selectivity regarding tumor cells (HeLa) [274]. Decrease of mitochondrial membrane potential, cells viability and activity of SOD, CAT, Gpx and also increase of MDA were observed after irradiation by visible light. Finally, the apoptosis proceeded through enhancement of caspase-3 activity.

C₆₀-pyroferoforbids with different amount of substitutes showed different phototoxicity in relation to leukemic human T-lymphocytes (Jurkat cells) [247]. Effects were determined by difference in their physical and chemical properties and, as consequence, by different intracellular uptakes.

Photoinduced cleavage of super-helix DNA pBR322 predominantly at guanine bases [269] was observed after incubation with carboxyfullerenes-C₆₀ and irradiation by visible light (but no in the darkness). Triads tetraphenylporphyrin (TPP)-PVP-C₆₀ [275] demonstrated good results in PDT. No dark cytotoxicity in relation of human lymphoblast cell line (K562, chronic leukemia) was observed under use of 1 $\mu\text{mole/L}$ of sensibilizer. However, apoptosis of cancer cells without participation of caspase-3 took place under triad concentrations of 0.05-5 $\mu\text{mole/L}$ with irradiation by light at $\lambda=436$ nm (1000 mJ/sm^2 , 20-200 mW/sm^2

PEG-fullerene-C₆₀ demonstrated phototoxicity ($\lambda=400-600$ nm, 140 J/cm^2) in relation to human fibrosarcoma cells HT1080, significantly decreasing their viability (from 50 to 30% in dependence on the conditions). Normal fibroblasts in the same conditions kept viability of 85-93%. It is considered that PEG-fullerenes C₆₀ possess good potential to be used in PDT with very small side effects for normal cells [276]. PEG-conjugated fullerenes, containing Gd³⁺ ion, were used for photodynamic therapy in combination with magnetic resonance tomography [277].

Co-polymers of C₆₀-N-vinylpyrrolidone demonstrated the best phototoxicity in relation to HeLa and murine osteogenic sarcoma cells under irradiation by light in comparison with other complexes of fullerene [278]. Cell apoptosis is already observed under low concentrations (5 $\mu\text{g/mL}$) of such a copolymer and more than 30% of cells perished at concentration of 100 $\mu\text{g/mL}$. A cell membrane was a target for damages. It is considered that calcium as a secondary messenger participates in this photo-induced process.

Fullerol C₆₀(OH)₂₄ is a powerful photosensibilizer [279] at the expense of superoxide radicals and singlet oxygen generation through effective resonance transfer of energy [280]. C₆₀(OH)₂₄ is used in PDT in ophthalmology, because it is able to overcome eye barriers and some correlation between its intracellular distribution and progression in damage of human lens and retina *in vivo* is observed [281].

Fullerol (C₆₀(OH)₁₉(ONA)₁₇18H₂O) also exhibited in water solutions a high phototoxicity in relation to cells and cellular compounds [282,283] that was explained by authors [279] as a result of possible accumulation of ROS-products.

Photosensibilizing and photoprotector role of fullerenes are intersected. Some works are devoted to joint role of fullerene as pro-oxidant as well as antioxidant in UVB-induced damages of different tissues, i.e. synergic or cumulative effects in PDT [13]. This problem is very important in production of cosmetic anti-sunburn means and also at some skin diseases. It is known that under UV-irradiation a set of events, leading to pro-apoptotic alterations (generation of ROS, cell rounding, bubbling of its surface etc.) and anoikis, is observed at the cellular level [284]. In mice sebaceous glands fulfill a role of a main site for ROS generation in UV-irradiation [285]. Use of C₆₀ did not lead to emergence of toxicity, but index of ROS and index of apoptosis were decreased. More significant decrease was detected at simultaneous use of ascorbic acid (AA) and fullerene that, possibly, is provided by binding of fullerene with AA and decreasing of the Fenton reaction yield due to intercalation of AA to the heme pocket. Thus,

use of a sum of fullerene+AA in combination with UV-irradiation is an effective remedy against oxidative damage of a skin.

C₆₀, incorporated into phospholipid membrane (74.5 nm, C=150 ppm), in the case of its introduction before or after UVA irradiation (10 J/cm²), restores viability of cells, decreasing at 30% the level of ROS [286]. Quick excretion of the drug is its shortage. Liposome-fullerene (250 ppm, with C= 0.75 ppm C₆₀) under full absence of own toxicity significantly inhibited damage caused by chronic UVA irradiation of skin (by 4 κJ/cm², in total 76 J/cm²) [287]. Carboxyfullerenes sufficiently decreased human keratinocytes proliferation blocking induced by UVB-irradiation [288], simultaneously decreasing amount of cells with depolarized mitochondria. The mechanism of action included interference of the preparation to the process of ROS generation by depolarized mitochondria, however, without participation of Bcl-2. In the same time, a cationic adduct of C₆₀ (Fig.7c,d) incorporated into liposome demonstrated high efficiency of HeLa cells damage in PDT [257].

Water-soluble nanoparticles containing fullerene-C₆₀ incorporated into liposomes (LPF, 75.6 nm, up to 0.3% C₆₀ by weight) showed dose-dependent protective effect HaCaT cells against OH-radicals, emerging under UVA- (12 J/cm²) and UVB- (500mJ/cm²) irradiation. Prevention of cell morphology degeneration was observed and any protective effect took place in the presence of C₆₀ only [289].

C3 is able to protect selectively cells against intracellular and/or membrane changes in UV-irradiation [152]. It suggested that protection of epithelial cells A431 is linked with ability of, namely, this compound to maintain a network of cytoskeleton components and the integrity of coordination linkage, but only in the case, when fullerene derivative presents during irradiation. It is not excluded that the ability of a preparation to capture the superoxide radical before its conversion into OH- radical is a substantial component of the antioxidant action. However, C3 have been localized in the cell membrane.

Skin keratinocytes (HaCaT) were used for probing of different fullerols: C₆₀(OH)₄₄ 8H₂O (SHH-F), C₆₀(OH)₆₋₁₂ (LH-F) and C₆₀(OH)₃₂₋₃₄ 7H₂O (HH-F) [290] in conditions of UVA and UVB-irradiation. Acceptor activity in relation to ROS was higher for HH-F and SHH-F, than for LH-F that determined a degree of their cytoprotector effect, which was greater in the case of SHH-F. In that, protective effect of SHH-F in relation to UVB- induced damages was higher than those in relation to UVA. Thus, SHH-F is a highly effective cytoprotector under UV-irradiation.

Water-soluble derivative (γ-CD)₂/C₆₀ is used in two ways: as drug transporter through eye barriers and photosensibilizer in PDT for tumors treatment [291]. 2 μmole/L of (γ-CD)₂/C₆₀ have already been toxic for HLE B-3 cells in UVA irradiation. However, the effect was not observed under irradiation by visible light and in the darkness. In the meantime, aggregated nanoparticles did not provide such an effect even at 30 μmole/L. Namely singlet oxygen is an important intermediate product of phototoxicity of monomer (γ-CD)₂/C₆₀ and its production decreases with particles aggregation degree growth. Photodynamic activity of C₇₀-γ-cyclodextrine complex was sufficiently greater than those for C₆₀ [15]. Similar picture was observed in comparison of photodynamic activity (λ>400 nm) of C₆₀ and C₇₀ incorporated into lipid membrane in relation to HeLa cells [292]. Authors consider that this is linked with a simple difference in their ability to generate singlet oxygen, due to asymmetry of C₇₀.

Investigation of [293] phototoxicity of three fullerene hexa-cis-adducts in combination with different amount of introduced photosensibilizers: bis (3¹,3²-didehydrophytychlorine) fullerene [5:1]-hexa-adduct (FHP1), fullerene [5:1]-hexa-adduct with six 3¹,3²- didehydrophytychlorine groups (FHP6) and fullerene[6:0]-hexa-adduct with 12- 3¹,3²- didehydrophytychlorine units (FHP12) was performed and FHP6 was the most prospective one. The degree of intracellular uptake, which was depended on the size and asymmetry of the fullerene complex, that affected the singlet oxygen quantum yield, had the greatest meaning.

Some fullerols cause phototoxicity of human retina pigment epithelial cells [114]. Early apoptosis proceeds already at concentrations >5 μmole/L and a dose of 8.5 J/cm² of visible light (quantum yield of singlet oxygen is 0.05).

High phototoxicity was observed under action of 1 μmole/L of diad porphyrin-C₆₀ (P-C₆₀), incorporated into liposomes, on the cell line of human larynx carcinoma Hep-2. Death of 80% of cells was observed under 54 J/cm². [294]. There is a first case, when a high ability of such compounds to form a photo-induced state with isolated (separated) charge was stressed. The cells viability depended on the light level. High cytotoxic effect was kept even in the argon atmosphere. Damages were caused through the mechanism of ¹O₂ – mediated photoreaction process, or through ROS attack under low oxygen concentration in dependence on sensibilizer localization site micro-environment. Caspase-3 pathway-dependent apoptosis (58% of apoptotic cells) was substituted by predominance of necrotic events under anaerobic conditions.

A fullerene-carbohydrate hybrid that did not require any additions was used for selective degradation of a target protein, HIV-1 protease, under irradiation by UV or visible light [211]. Carbohydrate-C₆₀ acts through generation of singlet oxygen during laser irradiation with $\lambda=355$ nm [295]. PDT did not reveal cytotoxicity in usage of fullerene glyco-conjugates against normal fibroblasts targeting, namely, cancer cells [296]. In that, carbohydrate-substituted C₆₀ inhibited lipid peroxidation in the blood plasma [297].

Cationic micelles of block-copolymers of poly(N-vinylcaprolactam)-C₆₀ demonstrated themselves as a potential antiviral drug [298] and, simultaneously, strong photosensibilizer under UV-irradiation [260].

Using BMPF (4 μ mole/L, irradiation by green light) and HeLa cells, it was shown [299] that ROS are involved in the process of sensitization of apoptosis and necrosis together with calcium ions. Simultaneously, BMPF enhances lipid peroxidation increasing the MDA level in dependence on dose and duration of irradiation [269]. Presence of extracellular calcium promotes activity of fullerene derivatives, but its removal does not interrupt their membrane damaging activity that indicates for existence of both calcium-dependent and calcium-independent ways in the process.

Simultaneous action of C₆₀ (12-50 nm, 50 μ mole/L) colloid solution and irradiation in UV- and visible light range caused cytotoxic effects on leukemic T-lymphocytes and Jurkat cell line (but not thymocytes) [300]. The process was accompanied by elevated activation of caspase-3 and depended on C₆₀ concentration.

A hypothesis regarding usage of fullerenes to mediate PDT of intraperitoneal carcinomatosis was checked [301], applying N-methyl-pyrrolidine-C₆₀, placed into Cremophore-EL-micelles, and illumination by white light through the abdominal wall. Domination of necrotic events over apoptosis was observed.

Dendrite C₆₀ [11] inhibited growth of cells in the darkness and was slightly phototoxic under UV-irradiation due to independent formation of aggregates. A method of preparation of fullerene nanoparticles played a great role in its phototoxic activity [260].

PVP-C₆₀ [302] repressed the changes, caused by UVA HaCaT cells in the form of translocations of the transcription factor NF- κ B into cytoplasm to the nucleus of keratinocytes. Protective effect and apoptosis abnormal signaling pathways blocking were observed.

Thus, photosensibilizing action of fullerene derivatives may be performed at the expense of ¹O₂ (C₆₀) and ROS enhancing (C₆₀(OH)₁₈). In that it is necessary to take into account the result of differential interaction with various structures, for example, binding to molecules of the lipid membrane, increasing of OH-generation etc. The effect may be additionally enhanced by using of light gathering antenna molecule [273]. The greatest phototoxicity is characteristic for mono-substituted C₆₀ adducts that have no an ability to self-aggregation, especially in the presence of calcium ions. The process is a dose- and concentration-dependent and the method of preparation of fullerene nanoparticles is of a great importance.

It is necessary to account that fullerenes may be more effective in PDT of hypoxia tumors (where the oxygen level is low), because they are able to change a mechanism of cell damage.

In the recent works [303] an idea of fullerene association with porphyrin structures in order to develop new photosensibilizers with increased generation of singlet oxygen and improved penetration into tumors is discussed. Some modern aspects of fullerene-mediated PDT, in particular, therapeutic perspectives of their use, are considered in the review [304]. In particular, using of this method for rescuing of mice, whose wounds were infected by gram-negative bacteria, was demonstrated.

IX. IONIZING RADIATION AND FULLERENES

It is known that impact of radiation in great degree is determined by generation of larger amounts of ROS. Cytotoxic or, on the contrary, antioxidant effects of fullerenes may play a substantial role in these conditions, increasing or compensating impact of ionizing radiation on the organism. Moreover, properties of the fullerenes may also be changed under action of ionizing radiation.

NanoC₆₀ in certain conditions inhibit tumor cells growth, expressing properties of a sensibilizer under radiotherapy, enhancing apoptosis [305].

Radioprotective (anti-radical) activity of hydrated fullerene C₆₀HyFn and its labile nanosize clusters at concentrations of 10⁻¹¹-10⁻⁶ mole/L was developed under X-ray irradiation of DNA (1-7 Gy *in vitro*) [306]. Using 8-

oxoguanine as a marker of DNA oxidative damage, it was shown that C₆₀HyFn at concentrations of 10⁻⁷-10⁻⁶ mole/L protects nucleic acids against radical-induced damage. An optimal radioprotective concentration of C₆₀HyFn was equal to 1 mg/kg *in vivo* (mice, intraperitoneally, 1 hour before or 15 min. after irradiation) at the lethal dose of 7 Gy. Lysine-C₆₀ did not express visible toxicity in relation to human lymphoblastic cells AHH-1 and was used for preliminary treatment of cells before γ -irradiation (>400 mg/L). This sufficiently increased the cell survival after irradiation, decreasing the apoptosis level in a dose-dependent manner [307].

Dendro-C₆₀-fullerene – the fullerene derivative, containing 18 carboxyl groups (C₆₀DF) [308], is able to protect human lymphocytes and intestine cells from consequences of impact of high doses of γ -irradiation. The process includes decrease of ROS level, inhibition of radiation-induced apoptosis and cells necrosis, DNA damages, oxidative stress. However, the process is not selective, because no difference between modest protective effect of the dendro-fullerene on normal fibroblasts and tumor cells was observed *in vivo*. Dose modifying factor (DMF) was equal of 1.1 [309]. LD_{50/30} for mice, received 300 mg/kg of the dendro-fullerene before irradiation, was equal to 10.09 Gy in comparison with 8.29 Gy for control group. No protective effect was described for a dose of the drug of 200 mg/kg. Protective effect, linked with antioxidant effects of C₆₀DF in more wide range of usage, was observed under irradiation of *Danio rerio* embryos in a dose of 20 Gy and 40 Gy [310].

γ -Irradiation is able to influence on cytotoxicity of n-C₆₀ (THF) [7]. Irradiated fullerene not only did not cause oxidative stress and induce ERK-dependent death of different mammalian cells, but, on the contrary, protected cells against oxidative stress, induced by starting THF-nC₆₀ or hydrogen peroxide. Thus, γ -irradiation is able to alter physical and chemical properties of n-C₆₀, leading to full loss of its cytotoxicity and transforming it into a cytoprotective agent. In this case, it is not possible to exclude a possibility of functionalization of fullerene surface by products of water radiolysis that, in addition to a possible role of the residual THF in cytotoxicity, is able to cause the loss of cytotoxicity and acquisition of new properties by the fullerene.

Observed protective effect of polyhydroxylated derivatives of fullerene attracts great attention, however, the mechanism of action is not clarified to the moment, excluding its antioxidant action. It is known [311,312] that fullerene derivatives, predominantly, linked with mitochondria. It is also possible that nanosize particles may be captured by reticule-endothelial cells [313]. Namely, localization of one or another fullerene derivatives, may serve as decisive factor of their effector activity. In the next work [314] C₆₀(OH)₂₄ was utilized in a dose of 40 mg/kg for 2 weeks before irradiation of the whole body of mice with a lethal dose of γ -irradiation (⁶⁰Co). In comparison with control group animals receiving the drug demonstrated decrease of mortality due to elevation of immunity, decrease of oxidative damages and improvement of mitochondria functioning (restoration of mitochondrial membrane potential).

Fullerols, exhibiting protective effect in relation to heart and liver against chronic toxicity induced by doxorubicin [315], possessed also radioprotective effect during X-ray irradiation of animals (8 Gy) [316].

C₆₀-PVP and γ -CD-C₆₀, acting as antioxidants, are able to be stabilizers of radioprotective properties of β -carotene, preventing its oxidation [317] that increases the interest to them as compensating drugs during cancer treatment using radiotherapy.

There is information [318] about the possibility of combined action sulfa-containing drugs as radioprotectors and derivatives of C₆₀, which in some cases results to reduce the side effects of the original drug (amifostine).





Fig.1. Hydroxylated fullerene (a), spheric cluster (C₆₀@{H₂O}_n)_m (b), C₆₀-folic acid (c).

Type 1: O₂^{-•}



Type 2: ¹O₂^{*}




Fig.2. Mechanisms of singlet oxygen and superoxide radical generation by fullerenes.




Fig.3. C₆₀-derivative and adenosine receptors in SK-N-MC cell line of human neuroepithelioma. Copy from [88].




Fig.4. A scheme of events in CMECs, according to [119], induced by hydrogen peroxide and treatment by C₆₀(C(COOH)₂)₂ nanoparticles. Changes are mediated by decreasing of JNK phosphorylation and inhibition of ROS production. There is down-regulation of the signaling pathway with decreasing of c-Jun and caspase-3 activation and inhibition of PARP cleavage and release of a mitochondrial cytochrome C.




Fig.5. The effect of $C_{60}(C(COOH)_2)_2$ on lysosomal and mitochondrial apoptotic pathway, according [158].




Fig.6. A site of carboxyfullerene binding with HSA, predicted by docking. Copy from [195].




Fig.7. (a)- Structure of supramolecular aggregate, octaamidoporphyrin with fullerene C₆₀. (b)-bakminsterfullerene (C₆₀) – 2 – (butadiene – 1- yl) tetra (o – γ – amino butyryl–o-phthalyl) ferroporphyrin (PMC16), containing 4 ions of magnetic isotope of ²⁵Mg, (c)- Complex of C₆₀ with a liposome, possessing, due to compounds 1-4 (d), different surface charge, according to [258].

In the same time, it is necessary to note that fullerenes as drugs transporters, in some cases, are much suitable than carbon nanotubes, possessing different (depending on shape and size), but significant and sustainable toxicity, in particular, to lungs [147,320].

It is very important that a system may be developed with fullerenes participation, when a single molecule combines acting component (for example, fullerene as generator of singlet oxygen), and a system of its targeting delivery to the target molecule (for example, cationic groups of side radical bind to anionic groups in DNA). So, it is absolutely necessary to develop new approaches to design preparations as well as experiments on nanomaterials assessment on animals.

Use of fullerenes in photo- and radio-therapy of cancer seems to be very perspective. In addition, searching of anti-tumor drugs among fullerenes is successfully continued.

A large amount of fullerene derivatives applications in medicine was described in [321]. At present, there are enough patents on concrete use of fullerene derivatives, for example, for treatment of granule cells and basophiles – mediated disease [322].

As a whole, active application of any nanoparticles in medicine must be under strict control till all their consequences (including remote ones) for the organism will be revealed in detail. One of the main research fields is revealing of preparation effects on apoptosis signaling pathways. At present, there are data regarding undoubtedly positive possibilities of fullerene nanoparticles as well as their potentially dangerous properties [323]. For example, neurons stopped to react on chemical signals and enter latent state in the presence of magnetic nanoparticles of less than 10 nm in size. Fullerenes in some conditions *in vitro* are able to kill liver, skin and brain cells [324,325]. Accumulation of nanoparticles in a cell medium under biodegradation is dangerous that may disrupt organelles integrity and even cause genetic mutations [323]. In the meantime, use of fullerene nanoparticles may bring a serious success today in therapy of such complicated diseases as cancer, HIV, AD or PD. Seemingly the greatest progress in near future will be seen in the field of construction of supramolecular nanoplatfroms with filling by differently acting drugs and drug delivery with developing of targeted therapy and personalized medicine.

Acknowledgment

This work was supported by RFBR grant № 10-08-00680. We wish to thank Professor Kuznetsov D.A. for valuable advice and Arkhangelsky S.E. – for the technical assistance.

REFERENCES

- [1] Duncan, R.; Gaspar, R. Nanomedicine(s) under the microscope. *Mol. Pharmaceutics*, **2011**, *8*, 2101–2141.
- [2] Kateb, B.; Chiu, K.; Black, K.L.; Yamamoto, V.; Khalsa, B.; Ljubimova, J.Y.; Ding, H.; Patil, R.; Portilla-Arias, J.A.; Modo, M.; Moore, D.F.; Farahani, K.; Okun, M.S.; Prakash, N.; Neman, J.; Ahdoot, D.; Grundfest, W.; Nikzad, S.; Heiss, J.H. Nanoplatfroms for constructing new approaches to cancer treatment, imaging, and drug delivery: What should be the policy? *NeuroImage*, **2011**, *54*, S106-S124.
- [3] Harhaji, L.; Isakovic, A.; Raicevic, N.; Markovic, Z.; Todorovic-Markovic, B.; Nikolic, N.; Vranjes-Djuric, S.; Markovic, I.; Trajkovic, V. Multiple mechanisms underlying the anticancer action of nanocrystalline fullerene. *Eur. J. Pharmacol.*, **2007**, *568*, 89-98.
- [4] Dugan, L.L.; Turetsky, D.M.; Du, C.; Lobner, D.; Wheeler, M.; Almlı, R.; Shen, C.K.F.; Luh, T.Y.; Choi, D.W.; Lin, T.S. Carboxyfullerenes as neuroprotective agents. *Proc. Natl. Acad. Sci. U.S.A.*, **1997**, *94*, 9434-9439.
- [5] Monti, D.; Moretti, L.; Salvioli, S.; Strafolacince, E.; Malorni, W.; Pellicciari, R.; Schettini, G.; Bisaglia, M.; Pincelli, C.; Fumelli, C.; Bonafe, M.; Franceschi, C. C₆₀ carboxyfullerene exerts a protective activity against oxidative stress-induced apoptosis in human peripheral blood mononuclear cells. *Biochem. Biophys. Res. Commun.*, **2000**, *277*, 711-717.
- [6] Dugan, L.L.; Gabrielsen, J.K.; Yu, S.P.; Lin, T.S.; Choi, D.W. Buckminsterfullerenol free radical scavengers reduce excitotoxic and apoptotic death of cultured cortical neurons. *Neurobiol. Dis.*, **1996**, *3*, 129-135.
- [7] Isakovic, A.; Markovic, Z.; Nikolic, N.; Todorovic-Markovic, B.; Vranjes-Djuric, S.; Harhaji, L.; Raicevic, N.; Romcevic, N.; Vasiljevic-Radovic, D.; Dramicanin, M.; Trajkovic, V. Inactivation of nanocrystalline C₆₀ cytotoxicity by γ -irradiation. *Biomaterials*, **2006**, *27*, 5049-5058.
- [8] Isakovic, A.; Markovic, Z.; Todorovic-Markovic, B.; Nikolic, N.; Vranjes-Djuric, S.; Mirkovic, M.; Dramicanin, M.; Harhaji, L.; Raicevic, N.; Nikolic, Z.; Trajkovic, V. Distinct cytotoxic mechanisms of pristine versus hydroxylated fullerene. *Toxicol. Sci.*, **2006**, *91*, 173–183.
- [9] Yamawaki, H.; Iwai, N. Cytotoxicity of water-soluble fullerene in vascular endothelial cells. *Am. J. Physiol. Cell Physiol.*, **2006**, *290*, C1495–C1502.

- [10] Yang, X.L.; Fan, C.H.; Zhu, H.S. Photo-induced cytotoxicity of malonic acid [C₆₀]fullerene derivatives and its mechanism. *Toxicol. InVitro*, **2002**, *16*, 41–46.
- [11] Rancan, F.; Rosan, S.; Boehm, F.; Cantrell, A.; Brellreich, M.; Schoenberger, H.; Hirsch, A.; Moussa, F. Cytotoxicity and photocytotoxicity of a dendritic C₆₀ mono-adduct and a malonic acid C₆₀ tris-adduct on Jurkat cells. *J. Photochem. Photobiol., B.*, **2002**, *67*, 157–162.
- [12] Bosi, S.; L.Feruglio, L.; T.Da Ros, T.; G.Spalluto, G.; B.Gregoretto, B.; M.Terdoslavich, M. Hemolytic effects of water-soluble fullerene derivatives. *J. Med. Chem.*, **2004**, *47*, 6711–6715.
- [13] Tabata, Y.; Ishii, T.; Aoyama, T.; Oki, R.; Hirano, Y.; Ogawa, O.; Ikada, Y. In: *Perspectives of Fullerene Nanotechnology*. Ed. E.Osawa, Kluwer Academic Publ., Dordrecht; Boston; London, **2001**.
- [14] Chen, C.; Xing, G.; Wang, J.; Zhao, Y.; Li, B.; Tang, J.; Jia, G.; Wang, T.; Sun, J.; Xing, L.; Yuan, H.; Gao, Y.; Meng, H.; Chen, Z.; Zhao, F.; Chai, Z.; Fang, X. Multihydroxylated [Gd@C₈₂(OH)₂₂]_n nanoparticles: antineoplastic activity of high efficiency and low toxicity. *Nano Lett.*, **2005**, *5*, 2050–2057.
- [15] Darwish, A.D. Fullerenes. *Annu. Rep. Prog. Chem., A*, **2010**, *106*, 356–375.
- [16] Shirinkin, S.V.; Volkova, T.O.; Nemova, N.N. In: “*Medicinal nanotechnology. The using of fullerenes in a therapy of respiratory diseases*”, Ed. Karelsky Sci. Centre RAS, Petrazavodsk. **2009**.
- [17] Gao, J.; Wang, H.L.; Shreve, A.; Iyer, R. Fullerene derivatives induce premature senescence: A new toxicity paradigm or novel biomedical applications. *Toxicol. Appl. Pharmacol.*, **2010**, *244*, 130–143.
- [18] Jung, H.; Wang, C.; Jang, W. Nano-C₆₀ and hydroxylated C₆₀: Their impacts on the environment, *Toxicol. Environ. Health Sci.*, **2009**, *1*, 132-139.
- [19] Sidorov, L.N.; Yurovskaya, M.A.; Borschevsky, A.Ya.; Trushkov, I.V.; Ioffe, I.N. *Fullerenes*, Ed. Examination, Moscow. **2005**.
- [20] Hirsch, A.; Brettreich, M. *Fullerenes: chemistry and reactions*. Wiley-VCH, Weinheim. **2005**.
- [21] Piotrovsky, L.B. *Fullerenes in biology*. Ed. Rostock. St. Petersburg. **2006**.
- [22] Troshin, P.A.; Lyubovskaya, Z.N. Organic chemistry of fullerenes: basic reaction, types of fullerene compounds and prospects for their practical application. *Adv. Chem. (Russ)*, **2008**, *77*, 324-369.
- [23] Ema, M.; Kobayashi, N.; Naya, M.; Hanai, S.; Nakanishi, Reproductive and developmental toxicity studies of manufactured nanomaterials. *J. Reprod. Toxicol.*, **2010**, *30*, 343-352..
- [24] Markovic, Z.; Trajkovic, V. Biomedical potential of the reactive oxygen species generation and quenching by fullerenes (C₆₀). *Biomaterials*, **2008**, *29*, 3561-3573.
- [25] Grausova, L.; Vacik, J.; Vorliceck, V.; Svorcik, V.; Slepicka, P.; Bilkova, P.; Vandrovцова, M.; Lisa, V.; Bacakova, L. Fullerene C₆₀ films of continuous and micropatterned morphology as substrates for adhesion and growth of bone cells. *Diamond Relat. Mater.*, **2009**, *18*, 578-586.
- [26] Aschberger, K.; Johnston, H.J.; Stone, V.; Aitken, R.J.; Tran, C.L.; Hankin, S.M.; Peters, S.A.; Christensen, F.M. Review of fullerene toxicity and exposure – Appraisal of a human health risk assessment, based on open literature. *Regul. Toxicol. Pharmacol.*, **2010**, *58*, 455-473.
- [27] Da Ros, T.; Spalluto, G.; Prato, M. Biological applications of fullerene derivatives: a brief overview. *Croat. Chem. Acta*, **2001**, *74*, 743-755.
- [28] Da Ros, T. Twenty Years of Promises: Fullerene in Medicinal Chemistry. *Carbon Mater.: Chem. Phys.*, **2008**, *1*, 1-21.
- [29] Bakry, R.; Vallant, R.M.; Najam-ul-Haq, M.; Rainer, M.; Szabo, Z.; Huck, C.; Bonn, G.K. Medicinal applications of fullerenes. *Int. J. Nanomed.*, **2007**, *4*, 639–649.
- [30] Satoh, M.; Takayanagi, I. Pharmacological Studies on Fullerene (C₆₀), a Novel Carbon Allotrope, and Its Derivatives. *J. Pharmacol. Sci.*, **2006**, *100*, 513-518.
- [31] Chawla, P.; Chawla, V.; Maheshwari, R.; Saraf, A.; Saraf, K. Fullerenes: from carbon to nanomedicine. *Mini Reviews in Med. Chem.*, **2010**, *10*, 662-677.
- [32] Partha, R.; Conyers, J.L. Biomedical applications of functionalized fullerene-based nanomaterials. *Int. J. Nanomed.*, **2009**, *4*, 261–275.
- [33] Yan, L.; Zhao, F.; Li, S.; Hu, Z.; Zhao, Y. Low-toxic and safe nanomaterials by surface-chemical design, carbon nanotubes, fullerenes, metallofullerenes, and graphenes. *Nanoscale*, **2011**, *3*, 362–382.
- [34] Parveen, S.; Misra, R.; Sahoo, S.K. Nanoparticles: a boon to drug delivery, therapeutics, diagnostics and imaging. *Nanomedicine: Nanotechnol. Biol. Med.*, **2012**, *8*, 147-166.
- [35] Huh, A.J.; Kwon, Y.J. “Nanoantibiotics”: A new paradigm for treating infectious diseases using nanomaterials in the antibiotics resistant era. *J. Control. Release*, **2011**, *7*, 128-145.
- [36] Koifman, O.I.; Mamardashvili, I.Z.; Antipin, I.S. *Synthetic receptors based on porphyrins and their conjugates with klicks [4] arenes*, Ed. Nauka, Moscow. **2006**.
- [37] Satake, A.; Kobuke, Y. Dynamic supramolecular porphyrin system. *Tetrahedron*, **2005**, *61*, 13-41.
- [38] Tykhomyrov, A.A.; Nedzvetsky, V.S.; Klochkov, V.K.; Andrievsky, G.V. Nanostructures of hydrated C₆₀ fullerene (C₆₀HyFn) protect rat brain against alcohol impact and attenuate behavioral impairments of alcoholized animals. *Toxicology*, **2008**, *246*, 158-165.

- [39] Spohn, P.; Hirsch, C.; Hasler, F.; Bruinink, A.; Krug, H.F.; Wick, P. C₆₀ fullerene: a powerful antioxidant or a damaging agent? The importance of an in-depth material characterization prior to toxicity assays. *Environ. Pollut.*, **2009**, *157*, 1134–1139.
- [40] Yokel, R.A.; MacPhail, R.C. Engineered nanomaterials: exposures, hazards, and risk prevention. *J. Occup. Med. Toxicol.*, **2011**, *6*, 7-34.
- [41] Buseck, P.R. Geological fullerenes: review and analysis. *Earth Planet. Sci. Lett.*, **2002**, *203*, 781-792.
- [42] Shinohara, N.; Gamo, M.; Nakanishi, J. Fullerene C₆₀: inhalation hazard assessment and derivation of a period-limited acceptable exposure level. *Toxicol. Sci.*, **2011**, *123*, 576-589.
- [43] Orlova, M.A.; Orlov, A.P. Role of zinc in an organism and its influence on processes leading to apoptosis. *Br. J. Med. Med. Res.*, **2011**, *1*, 239-305.
- [44] Portt, L.; Norman, G.; Clapp, C.; Greenwood, M. Anti-apoptosis and cell survival. *Biochim. Biophys. Acta*, **2011**, *1813*, 238-259.
- [45] Shvedova, A.A.; Kagan, V.E.; Fadeel, B. Close encounters of the small kind: adverse effects of man-made materials interfacing with the nano-cosmos of biological systems. *Annu. Rev. Pharmacol. Toxicol.*, **2010**, *50*, 63-88.
- [46] Usenko, C.Y.; Harper, S.L.; Tanguay, R.L. Fullerene C₆₀ exposure elicits an oxidative stress response in embryonic zebrafish. *Toxicol. Appl. Pharmacol.*, **2008**, *229*, 44-55.
- [47] Yamakoshi, Y.; Umezawa, N.; Ryu, A.; Arakane, K.; Miyata, N.; Goda, Y. Active oxygen species generated from photoexcited fullerene (C₆₀) as potential medicines: O₂^{•-} versus ¹O₂. *J. Am. Chem. Soc.*, **2003**, *125*, 12803-12809.
- [48] Scrivens, W.A.; Tour, J.M.; Creek, K.E.; Pirisi, L. Synthesis of ¹⁴C-labeled C₆₀, its suspension in water, and its uptake by human keratinocytes. *J. Am. Chem. Soc.*, **1994**, *116*, 4517-4518.
- [49] Maeda, R.; Noiri, E.; Isobe, H.; Homma, T.; Tanaka, T.; Negishi, K.; Doi, K.; Fujita, T.; Nakamura, E. A water-soluble fullerene vesicle alleviates angiotensin II-induced oxidative stress in human umbilical venous endothelial cells. *Hypertension Res.*, **2008**, *31*, 141-151.
- [50] Fortner, J.D.; Lyon, D.Y.; Sayes, C.M.; Boyd, A.M.; J.C.Falkner, J.C.; E.M.Hotze, E.M.; L.B.Aleman, L.B.; Y.J.Tao, Y.J.; Guo, W.; Ausman, K.D.; Colvin, V.L.; Hughes, J.B. C-60 in water: nanocrystal formation and microbial response. *Environ. Sci. Technol.*, **2005**, *39*, 4307–4316.
- [51] Henry, T.B.; Menn, F.M.; Fleming, J.T.; Wilgus, J.; Compton, R.N.; Saylor, G.S. Attributing effects of aqueous C-60 nano-aggregates to tetrahydrofuran decomposition products in larval zebrafish by assessment of gene expression. *Environ. Health Perspect.*, **2007**, *115*, 1059–1065.
- [52] Colvin, V.L. The potential environmental impact of engineered nanomaterials. *Nat. Biotechnol.*, **2003**, *21*, 1166-1170.
- [53] Sayes, C.M.; Gobin, A.M.; Ausman, K.D.; Mendez, J.; West, J.; Colvin, V.L. Nano-C₆₀ cytotoxicity is due to lipid peroxidation. *Biomaterials*, **2005**, *26*, 7587-7595.
- [54] Oberdorster, E.; Zhu, S.; Blickley, T.; McClellan-Green, P.; Haasch, M. Ecotoxicology of carbon-based engineered nanoparticles: effects of fullerene (C₆₀) on aquatic organisms. *Carbon*, **2006**, *44*, 1112-1120.
- [55] Andrievsky, G.; Klochkov, V.; Derevyanchenko, L. Is the C₆₀ fullerene molecule toxic?! *Fullerenes Nanotubes Carbon Nanostructures*, **2005**, *13*, 363-376.
- [56] Andrievsky, G.; Klochkov, V.; Bordyuh, A.; Dovbeshko, G. Comparative analysis of two aqueous–colloidal solutions of C₆₀ fullerene with help of FTIR reflectance and UV–vis spectroscopy. *Chem. Phys. Lett.*, **2002**, *364*, 8-17.
- [57] Gharbi, N.; Pressac, M.; Hadchouel, M.; Szwarc, H.; Wilson, S.R.; Moussa, F. [60]fullerene is a powerful antioxidant *in vivo* with no acute or subacute toxicity. *Nano Lett.*, **2005**, *5*, 2578-2585.
- [58] Kolosnjaj, J.; Szwarc, H.; Moussa, F. Toxicity studies of fullerenes and derivatives. *Adv. Exp. Med. Biol.*, **2007**, *620*, 168-180.
- [59] Jia, G.; Wang, H.; Yan, L.; Wang, X.; Pei, R.; Yan, T.; Zhao, Y.; Guo, X. Cytotoxicity of carbon nanomaterials: single-wall nanotube, multi-wall nanotube, and fullerene. *Environ. Sci. Technol.*, **2005**, *39*, 1378–1383.
- [60] Sayes, C.M.; Marchione, A.A.; Reed, K.L.; Warheit, D.B. Comparative pulmonary toxicity assessments of C₆₀ water suspensions in rats: few differences in fullerene toxicity *in vivo* in contrast to *in vitro* profiles. *Nano Lett.*, **2007**, *7*, 2399-2406.
- [61] Baker, G.L.; Gupta, A.; Clark, M.L.; Valenzuela, B.R.; Staska, L.M. Inhalation toxicity and lung toxicokinetics of C60 fullerene nanoparticles and microparticles. *Toxicol. Sci.*, **2008**, *101*, 122-131.
- [62] Horie, M.; Nishio, K.; Kato, H.; Shinohara, N.; Nakamura, A.; Fujita, K.; Kinugasa, S.; Endoh, S.; Yamamoto, K.; Yamamoto, O.; Niki, E.; Yoshida, Y.; Iwahashi, H. *In vitro* evaluation of cellular responses induced by stable fullerene C₆₀ medium dispersion. *J. Biochem.*, **2010**, *148*, 289-298.
- [63] Osuna, S.; Swart, M.; Solà, M. On the Mechanism of action of fullerene derivatives in superoxide dismutation. *Chemistry – Eur. J.*, **2010**, *16*, 3207-3214.
- [64] Zhou, S.; Burger, C.; Chu, B.; Sawamura, M.; Nagahama, N.; Toganoh, M. Spherical bilayer vesicles of fullerene-based surfactants in water: a laser light scattering study. *Science*, **2001**, *291*, 1944-1947.

- [65] Sawamura, M.; Kawai, K.; Matsuo, Y.; Kanie, K.; Kato, T.; Nakamura, E. Stacking of conical mesogens with a fullerene apex into polar columns in crystals and liquid crystals. *Nature*, **2002**, *419*, 702-705.
- [66] Yin, J.J.; Lao, F.; Fu, P.P.; Wamer, W.G.; Zhao, Y.; Wang, P.C. The scavenging of reactive oxygen species and the potential for cell protection by functionalized fullerene materials. *Biomaterials*, **2009**, *30*, 611-621.
- [67] Tokuyama, H.; Yamago, S.; Nakamura, E.; Shiraki, T.; Sugiura, Y. Photoinduced biochemical-activity of fullerene carboxylic-acid. *J. Am. Chem. Soc.*, **1993**, *115*, 7918-7919.
- [68] Husebo, L.O.; Sitharaman, B.; Furukawa, K.; Kato, T.; Wilson, L.J. Fullerenols revisited as stable radical anions. *J. Am. Chem. Soc.*, **2004**, *126*, 12055-12064.
- [69] Yamakoshi, Y.N.; Yagami, T.; Sueyoshi, S.; Miyata, N. Acridine adduct of [60]fullerene with enhanced DNA-cleaving activity. *J. Org. Chem.*, **1996**, *61*, 7236-7237.
- [70] Andersson, T.; Nilsson, K.; Sundahl, M.; Westman, G.; Wennerstrom, O. C₆₀ embedded in gamma-cyclodextrin—a water-soluble fullerene. *J. Chem. Soc. Chem. Commun.* **1992**, *8*, 604-606.
- [71] Makha, M.; Purich, A.; Raston, C.L.; Sobolev, A.N. Structural diversity of host-guest and intercalation complexes of fullerene C₆₀. *Eur. J. Inorg. Chem.*, **2006**, 507-517.
- [72] Deguchi, S.; Mukai, S.A.; Tsudome, M.; Horikoshi, K. Facile generation of fullerene nanoparticles by hand-grinding. *Adv. Mater.*, **2006**, *18*, 729-732..
- [73] Quaranta, A.; Zhang, Y.; Filippone, S.; Yang, J.; Sinay, P.; Rassat, A.; Edge, R.; Navaratnam, S.; McGarvey, D.; Land, E.J.; Brettreich, M.; Hirsch, A.; Bensasson, R.V. Photophysical studies of six amphiphilic 2:1 cyclodextrin:[60]fullerene derivatives. *Chem. Phys.*, **2006**, *325*, 397-403.
- [74] Dhawan, A.; Taurozzi, J.S.; Pandey, A.K.; Shan, W.Q.; Miller, S.M.; Hashsham, S.A.; Tarabara, V.V. Stable colloidal dispersions of C₆₀ fullerenes in water: evidence for genotoxicity. *Environ. Sci. Technol.* **2006**, *40*, 7394-7401.
- [75] Deguchi, S.; Alargova, R.G.; Tsujii, K. Stable dispersions of fullerenes, C₆₀ and C₇₀, in water: preparation and characterization. *Langmuir*, **2001**, *17*, 6013-6017.
- [76] Lyon, D.Y.; Adams, L.K.; Falkner, J.C.; Alvarez, P.J. Antibacterial activity of fullerene water suspensions: effects of preparation method and particle size. *Environ. Sci. Technol.*, **2006**, *40*, 4360-4366.
- [77] Brant, J.A.; Labille, J.; Bottero, J.Y.; Wiesner, M.R. Characterizing the impact of preparation method on fullerene cluster structure and chemistry. *Langmuir*, **2006**, *22*, 3878-3885.
- [78] Shinohara, N.; Matsumoto, K.; Endoh, S.; Maru, J.; Nakanishi, J. *In vitro* and *in vivo* genotoxicity tests on fullerene C₆₀ nanoparticles. *Toxicol. Lett.*, **2009**, *191*, 289-296.
- [79] Seki, M.; Fujishima, S.; Gondo, Y.; Inoue, Y.; Nozaka, T.; Suemura, K.; Takatsuki, M. Acute toxicity of fullerene C₆₀ in aquatic organism. *Environ. Sci.*, **2008**, *21*, 53-62.
- [80] Cook, S.M.; Aker, W.G.; Rasulev, B.F.; Hwang, H.M.; Leszczynski, J.; Jenkins, J.; Shockley, V. Choosing safe dispersing media for C₆₀ fullerenes by using cytotoxicity tests on the bacterium *Escherichia coli*. *J. Hazard Mater.*, **2010**, *176*, 367-373.
- [81] Cho, M.; Fortner, J.D.; Hughes, J.B.; Kim, J.H. *Escherichia coli* inactivation by water-soluble, ozonated C₆₀ derivative: kinetics and mechanisms. *Environ. Sci. Technol.*, **2009**, *43*, 7410-7415.
- [82] Bosi, S.; Da Ros, T.; Spalluto, G.; Prato, M. Fullerene derivatives: an attractive tool for biological applications. *Eur. J. Med. Chem.*, **2003**, *38*, 913-923.
- [83] Brettreich, M.; Hirsch, A. A highly water-soluble dendro[60]fullerene. *Tetrahedron Lett.* **1998**, *39*, 2731- 2734.
- [84] Markovic, Z.; Todorovic-Markovic, B.; Kleut, D.; Nikolic, N.; Vranjes-Djuric, S.; Misirkic, M.; Vucicevic, L.; Janjetovic, K.; Isakovic, A.; Harhaji, L.; Babic-Stojic, B.; Dramicanin, M.; Trajkovic, V. The mechanism of cell-damaging reactive oxygen generation by colloidal fullerenes. *Biomaterials*, **2007**, *28*, 5437-5448.
- [85] Karkischenko, N.N. Nanosafly: new approaches to risk assessment and toxicity of nanomaterials. *Biomedicine (Russ)*, **2009**, #1, 5-27.
- [86] Nishimura, T.; Kubota, R.; Tahara, M.; Nagaoka-Hamano, M.; Shimizu, K.; Hirose, A.; Tokunaga, H. Biological effects of fullerene C₆₀ in mouse embryonic stem cells. *Toxicol. Lett.*, **2006**, *164S*, S214.
- [87] Morimoto, Y.; Hirohashi, M.; Ogami, A.; Oyabu, T.; Myojo, T.; Nishi, K.; Kadoya, C.; Todoroki, M.; Yamamoto, M.; Murakami, M.; Shimada, M.; Wang, W.; Yamamoto, K.; Fujita, K.; Endoh, S.; Uchida, K.; Shinohara, N.; Nakanishi, J.; Tanaka, I. Inflammogenic effect of well-characterized fullerenes in inhalation and intratracheal instillation studies. *Part. Fibre Toxicol.*, **2010**, *7*, 4-22.
- [88] Giust, D.; Leon, D.; Ballesteros-Yanez, I.; Da Ros, T.; Albasanz, J.L.; Martín, M. Modulation of adenosine receptors by [60]fullerene hydrosoluble derivative in SK-N-MC cells. *ACS Chem. Neurosci.*, **2011**, *2*, 363-369.
- [89] Trpkovic, A.; Todorovic-Markovic, B.; Kleut, D.; Misirkic, M.; Janjetovic, K.; Vucicevic, L.; Pantovic, A.; Jovanovic, S.; Dramicanin, M.; Marcovic, Z.; Trajkovic, V. Oxidative stress-mediated hemolytic activity of solvent exchange-prepared fullerene (C₆₀) nanoparticles. *Nanotechnol.*, **2010**, *21*, 375102.
- [90] Costa, C.L.A.; Chaves, I.S.; Ventura-Lima, J.; Ferreira, J.L.R.; Ferraz, L.; de Carvalho, L.; Monserrat, J.M. *In vitro* evaluation of co-exposure of arsenium and an organic nanomaterial (fullerene, C₆₀) in zebrafish hepatocytes. *Comp. Biochem. Physiol. C*, **2012**, *155*, 206-212.

- [91] Sayes, C.; Fortner, J.; Guo, W.; Lyon, D.; Boyd, A.; Ausman, K.; Tao, Y.; Sitharaman, B.; Wilson, L.; Hughes, J.; West, J.; Colvin, V. The differential cytotoxicity of water soluble fullerenes. *Nano Lett.*, **2004**, *4*, 1881-1887
- [92] Lai, Y.L.; Murugan, P.; Hwang, K.C. Fullerene derivative attenuates ischemia-reperfusion-induced lung injury. *Life Sci.*, **2003**, *72*, 1271-1278.
- [93] Kolesnichenko, A.V.; M.A. Timofeyev, M.A.; M.V. Protopopova, M.V. Toxicity of nanomaterials – 15 years of research. *Russ. Nanotechnol. (Russ)*, **2008**, *3*, 54-61.
- [94] Hsu, H.C.; Chiang, Y.Y.; Chen, W.J.; Lee, Y.T. Water-soluble hexasulfobutyl-[60]-fullerene inhibits plasma lipid peroxidation by direct association with lipoproteins. *J. Cardiovasc. Pharmacol.*, **2000**, *36*, 423-427.
- [95] Lee, Y.T.; Chiang, L.Y.; Chen, W.J.; Hsu, H.C. Water-soluble hexasulfobutyl-[60]-fullerene inhibits low-density lipoprotein oxidation in aqueous and lipophilic phases. *Proc. Soc. Exp. Biol. Med.*, **2000**, *224*, 69-75.
- [96] Hu, Z.; Guan, W.C.; Tang, X.Y.; Huang, L.Z.; Xu, H. Synthesis of water-soluble cystine C₆₀ derivative with catalyst and its active oxygen radical scavenging ability. *Chinese Chem. Lett.*, **2007**, *18*, 51-54.
- [97] Hu, Z.; Xing, H.P.; Zhu, Z.; Wang, W.; Guan, W.C. Synthesis of cystine C₆₀ derivative and its protective effects on hydrogen peroxide-induced apoptosis in PC12 cells. *Chinese Chem. Lett.*, **2007**, *18*, 145-148.
- [98] Hu, Z.; Guan, W.; Wang, W.; Huang, L.; Xing, H.; Zhu, Z. Protective effects of a novel cystine C₆₀ derivative on hydrogen peroxide-induced apoptosis in rat pheochromocytoma PC12 cells. *Chem. Biol. Interact.*, **2007**, *167*, 135-144.
- [99] Hu, Z.; Guan, W.; Wang, W.; Huang, L.; Tang, X.; Xu, H.; Zhu, Z.; Xie, X.; Xing, H. Synthesis of amphiphilic amino acid C₆₀ derivatives and their protective effect on hydrogen peroxide-induced apoptosis in rat pheochromocytoma cells. *Carbon*, **2008**, *46*, 99-109.
- [100] Hu, Z.; Guan, W.; Wang, W.; Zhu, Z.; Wang, Y. Folacin C₆₀ derivative exerts a protective activity against oxidative stress-induced apoptosis in rat pheochromocytoma cells. *Bioorg. Med. Chem. Lett.*, **2010**, *20*, 4159-4162.
- [101] Hu, Z.; Liu, S.; Wei, Y.; Tong, E.; Cao, F.; Guan, W. Synthesis of glutathione C₆₀ derivative and its protective effect on hydrogen peroxide-induced apoptosis in rat pheochromocytoma cells. *Neurosci. Lett.*, **2007**, *429*, 81-86.
- [102] Ali, S.S.; Hardt, J.I.; Quick, K.L.; Kim-Han, J.S.; Erlanger, B.F.; Huang, T.T.; Epstein, C.J.; Dugan, L.L. A biologically effective fullerene (C₆₀) derivative with superoxide dismutase mimetic properties. *Free Radic. Biol. Med.*, **2004**, *37*, 1191-1202.
- [103] Wang, I.C.; Tai, L.A.; Lee, D.D.; Kanakamm, P.P.; Shen, C.K.; Luh, T.Y.; Cheng, C.H.; Hwang, K.C. C₆₀ and water-soluble fullerene derivatives as antioxidants against radical-initiated lipid peroxidation. *J. Med. Chem.*, **1999**, *42*, 4614-4620.
- [104] Guan, S.; Bao, Y.; Jiang, B.; An, L. Protective effect of protocatechuic acid from *Alpinia oxyphylla* on hydrogen peroxide-induced oxidative PC12 cell death. *Eur. J. Pharmacol.*, **2006**, *538*, 73-79.
- [105] Xiao, L.; H. Takada, H.; K. Maeda, K.; M. Haramoto, M.; N. Miwa, N. Antioxidant effects of water-soluble fullerene derivatives against ultraviolet ray or peroxy lipid through their action of scavenging the reactive oxygen species in human skin keratinocytes. *Biomed. Pharmacotherapy*, **2005**, *59*, 351-358.
- [106] Tong, J.; Zimmerman, M.C.; Li, S.; Yi, X.; Luxenhofer, R.; Jordan, R.; Kabanov, A.V. Neuronal uptake and intracellular superoxide scavenging of a fullerene (C₆₀)-poly(2-oxazoline)s nanoformulation. *Biomaterials*, **2011**, *32*, 3654-3665.
- [107] Alcaraz, M.J.; Megirias, J.; Garcia-Arnandis, I.; Clerigues, V.; Guillern, M.I. New molecular targets for the treatment of osteoarthritis. *Biochem. Pharmacol.*, **2010**, *80*, 13-21.
- [108] Bal, R.; Turk, G.; Tuzcu, M.; Yilmaz, O.; Ozercan, I.; Kuloglu, T.; Gur, S.; Nedzvetsky, V.S.; Tykomyrov, A.A.; Andrievsky, G.V.; Baydas, G.; Naziroglu, M. Protective effects of nanostructures of hydrated C₆₀ fullerene on reproductive function in streptozotocin-diabetic male rats. *Toxicology*, **2011**, *282*, 69-81.
- [109] Zhu, L.; Chang, D.W.; Dai, L.; Hong, Y. DNA damage induced by multiwalled carbon nanotubes in mouse embryonic stem cells. *Nano. Lett.*, **2007**, 3592-3597.
- [110] Xiao, L.; Aoshima, H.; Saitoh, Y.; Miwa, N. Highly hydroxylated fullerene localizes at the cytoskeleton and inhibits oxidative stress in adipocytes and a subcutaneous adipose-tissue equivalent. *Free Radic. Biol. Med.*, **2011**, *51*, 1376-1389.
- [111] Mirkov, S.M.; Djordjevic, A.N.; Andric, N.L.; Andric, S.A.; Kostic, T.S.; Bogdanovic, G.M.; Vojinovic-Miloradov, M.B.; Kovacevic, R.Z. Nitric oxide-scavenging activity of polyhydroxylated fullereneol, C₆₀(OH)₂₄. *Nitric Oxide*, **2004**, *11*, 201-207.
- [112] Injac, R.; Perse, M.; Obermajer, N.; Djordjevic-Milic, V.; Prijatelj, M.; Djordjevic, A.; Cerar, A.; Strukelj, B. Potential hepatoprotective effects of fullereneol C₆₀(OH)₂₄ in doxorubicin-induced hepatotoxicity in rats with mammary carcinomas. *Biomaterials*, **2008**, *29*, 3451-3460.
- [113] Bogdanovic, G.; Koji, V.; Dordevic, A.; Canadanovic-Brunet, J.; Vojinovic-Miloradov, M.; Balti, V.V. *Toxicol. In Vitro*, **2004**, *18*, 629-637. Modulating activity of fullerol C₆₀(OH)₂₂ on doxorubicin-induced cytotoxicity.
- [114] Wielgus, A.R.; Zhao, B.; Chignell, C.F.; Hu, D.N.; Roberts, J.E. Phototoxicity and cytotoxicity of fullerol in human retinal pigment epithelial cells. *Toxicol. Appl. Pharmacol.*, **2010**, *242*, 79-90.
- [115] Inui, S.; Aoshima, H.; Nishiyama, A.; Itami, S. Improvement of *acne vulgaris* by topical fullerene application: unique impact on skin care. *Nanomed. Nanotechnol. Biol. Med.*, **2011**, *7*, 238-241.

- [116]Mori, T.; Ito, S.; Namiki, M.; Suzuki, T.; Kobayashi, S.; Matsubayashi, K.; Sawaguchi, T. Involvement of free radicals followed by the activation of phospholipase A2 in the mechanism that underlies the combined effects of methamphetamine and morphine on subacute toxicity or lethality in mice: Comparison of the therapeutic potential of fullerene, mepacrine, and cooling. *Toxicology*, **2007**, *236*, 149-157.
- [117]Cagle, D.W.; Kennel, S.J.; Mirzadeh, S.; Alford, J.M.; Wilson, L.J. *In vivo* studies of fullerene-based materials using endohedral metallofullerene radiotracers. *Proc. Natl. Acad. Sci. USA*, **1999**, *96*, 5182-5187.
- [118]Oberdorster, E. Manufactured nanomaterials (fullerenes, C₆₀) induce oxidative stress in the brain of juvenile largemouth bass. *Environ. Health Perspect.*, **2004**, *112*, 1058-1062.
- [119]Lao, F.; Chen, L.; Li, W.; Ge, C.; Qu, Y.; Sun, Q.; Zhao, Y.; Han, D.; Chen, C. Fullerene nanoparticles selectively enter oxidation-damaged cerebral microvessel endothelial cells and inhibit JNK related apoptosis. *AcsNano*, **2009**, *3*, 3358-3368.
- [120]Rouzer, C.A. Nanoparticle tumorigenicity. *Chem. Res. Toxicol.*, **2010**, *23*, 4-5.
- [121]Podolsky, I.Ya.; Kondratieva, E.V.; Scheglov, I.V.; Dumpis, M.A.; Piotrovsky, L.B. C₆₀ with polyvinylpyrrolidone to prevent violation of the formation of long-term memory. *Solid State Physics (Russ)*, **2002**, *44*, 552-559.
- [122]Zaporotskova, I.V.; Chernozatonskiy, L.A. The mechanism the positive effect of fullerene on the recovery of spatial memory. *Bull. New Med. Technol.(Russ.)*, **2005**, *12*, 117-119.
- [123]Dugan, L.L.; Lovett, E.G.; Quick, K.L.; Lotharius, J.; Lin, T.T.; O'Malley, K.L. Fullerene-based antioxidants and neurodegenerative disorders. *Parkinson. Relat. Disord.*, **2001**, *7*, 243-246.
- [124]Lin, J.; Wu, C. Surface characterization and platelet adhesion studies on polyurethane surface immobilized with C₆₀. *Biomaterials*, **1999**, *20*, 1613- 1620.
- [125]Chen, T.; Li, Y.; Zhang, J.; Xu, B.; Lin, Y.; Wang, C.; Guan, W.; Wang, Y.; Xu, S. Protective effect of C₆₀-methionine derivate on lead-exposed human SH-SY5Y neuroblastoma cells. *J. Appl. Toxicol.*, **2011**, *31*, 255-261.
- [126]Linazasoro, G. Potential applications of nanotechnologies to Parkinson's disease therapy. *Parkinson. Relat. Disor.*, **2008**, *14*, 383-392.
- [127]Schloss, J.V.; Wu, J.Y. Polyhydroxylated C₆₀, fullerenols, as glutamate receptor antagonists and neuroprotective agents. *J. Neurosci. Res.*, **2000**, *62*, 600-607.
- [128]Lee, C.M.; Huang, S.T.; Huang, S.H.; Lin, H.W.; Tsai, H.P.; Wu, J.Y.; Lin, C.M.; Chen, C.T. C₆₀ fullerene-pentoxifylline dyad nanoparticles enhance autophagy to avoid cytotoxic effects caused by the β -amyloid peptide. *Nanomed. Nanotechnol. Biol. Med.*, **2011**, *7*, 107-114.
- [129]Szaraz, P.; Banhegyi, G.; Benedetti, A. Altered redox state of luminal pyridine nucleotides facilitates the sensitivity towards oxidative injury and leads to endoplasmic reticulum stress dependent autophagy in HepG2 cells. *Int. J. Biochem. Cell Biol.*, **2010**, *42*, 157-166.
- [130]Hoyer-Hansen, M.; Bastholm, L.; Szyniarowski, P.; Campanella, M.; Szabadkai, G.; Farkas, T. Control of macroautophagy by calcium, calmodulin-dependent kinase kinase- β , and Bcl-2. *Mol. Cell*, **2007**, *25*, 193-205.
- [131]Lu, T.; Kao, P.; Lee, C.; Huang, S.; Lin, C. C₆₀ Fullerene nanoparticle prevents β -amyloid peptide induced cytotoxicity in neuro 2A cells. *J. Food Drug Anal.*, **2011**, *19*, 151-158.
- [132]Podolski, I.Y.; Podlubnaya, Z.A.; Kosenko, E.A.; Mugantseva, E.A.; Makarova, E.G.; Marsagishvili, L.G. Effects of hydrated forms of C60 fullerene on amyloid 1-peptide fibrillization in vitro and performance of the cognitive task. *J. Nanosci. Nanotechnol.*, **2007**, *7*, 1479-1485.
- [133]Brambilla, D.; Droumaguet, B.L.; Nicolas, J.; Hashemi, S.H.; Wu, L.P.; Moghimi, S.M.; Couvreur, P.; Andrieux, K. Nanotechnologies for Alzheimer's disease: diagnosis, therapy, and safety issues. *Nanomed.: Nanotechnol. Biol. Med.*, **2011**, *7*, 521-540.
- [134]Lin, A.M.; Yang, C.H.; Ueng, Y.; Luh, T.Y.; Liu, T.Y.; Lay, Y.P.; Ho, L.T. Differential effects of carboxyfullerene on MPP+/MPTP-induced neurotoxicity. *Neurochem. Int.*, **2004**, *44*, 99-105.
- [135]Hartig, W.; Paulke, B.R.; Varga, C.; Seeger, J.; Harkany, T.; Kacza, J. Electron microscopic analysis of nanoparticles delivering thioflavin-T after intrahippocampal injection in mouse: implications for targeting betaamyloid in Alzheimer's disease. *Neurosci. Lett.*, **2003**, *338*, 174-176.
- [136]Gross, C.G. Neurogenesis in the adult brain: death of a dogma. *Nature Rev. Neurosci.*, **2000**, *1*, 67-72.
- [137]Lebovits, R.; Rosenblum, M. Substituted fullerenes and their use as inhibitors of cell death. *US Patent*, #0197950 A1, **2009**.
- [138]Friedman, S.H.; De Camp, D.L.; Sijbesma, R.P.; Srdanov, G.; Wudl, F.; Kenyon, G.L. Inhibition of the HIV-1 protease by fullerene derivatives: model building studies and experimental verification. *J. Am. Chem. Soc.*, **1993**, *115*, 6506-6509.
- [139]Park, K.H.; Chhowalla, M.; Iqbal, Z.; Sesti, F. Single-walled carbon nanotubes are a new class of ion channel blockers. *J. Biol. Chem.*, **2003**, *278*, 50212-50216.
- [140]Redmill, P.S.; McCabe, C. Molecular dynamics study of the behavior of selected nanoscale building blocks in a gel-phase lipid bilayer. *J. Phys. Chem. B*, **2010**, *114*, 9165-9172.
- [141]Kraszewski, S.; Tarek, M.; Treptow, W.; Ramseyer, C. Affinity of C60 neat fullerenes with membrane proteins: a computational study on potassium channels. *ACS Nano*, **2010**, *4*, 4158-4164.

- [142]Orlova, M.A.; Osipova, E.Y.; Roumiantsev, S.A. Effect of ⁶⁷Zn-nanoparticles on leukemic cells and normal lymphocytes. *Br. J. Med. Med. Res.*, **2012**, *2*, 21-30.
- [143]Andreev, I.; Petrukhina, A.; Garmanova, A.; Babakhin, A.; Andreev, S.; Romanova, V.; Troshin, P.; Troshina, O.; DuBuske, L. Penetration of fullerene C₆₀ derivatives through biological membranes. *Fullerenes Nanotubes & Carbon Nanostructures*, **2008**, *16*, 89-102.
- [144]Hsu, S.; Chien, C. Forced expression of Bcl-2 and Bcl-X(L) by novel water-soluble fullerene, C-60(glucosamine)(6), reduces renal ischemia/reperfusion-induced oxidative stress. *Fullerene Sci. Technol.*, **2001**, *9*, 77-88.
- [145]Cha, Y.J.; Lee, J.; Choi, S.S. Apoptosis-mediated *in vivo* toxicity of hydroxylated fullerene nanoparticles in soil nematode *Caenorhabditis elegans*. *Chemosphere*, **2012**, *87*, 49-54.
- [146]Johnson, G.L.; Lapadat, R.L. Mitogen-activated protein kinase pathways mediated by ERK, JNK and p38 protein kinases. *Science*, **2002**, *298*, 1911-1912.
- [147]Fatkhutdinova, L.M.; Khaliullin, T.O.; Zalyalov, R.R. Toxicity of engineered nanoparticles. *Kazanskii Med. J. (Russ.)*, **2009**, *90*, 578-584.
- [148]Manna, S.K.; Sarkar, S.; Barr, J.; Wise, K.; Barrera, E.V.; Jejelowo, O.; Rice-Ficht, A.C.; Ramesh, G.T. Single-walled carbon nanotube induces oxidative stress and activates nuclear transcription factor-κB in human keratinocytes. *Nano Lett.*, **2005**, *5*, 1676-1684.
- [149]Yudoh, K.; Karasawa, R.; Masuko, K.; Kato, T. Water-soluble fullerene (C₆₀) inhibits the osteoclast differentiation and bone destruction in arthritis. *Int. J. Nanomed.*, **2009**, *4*, 233-239.
- [150]Okada, T.; Otani, H.; Wu, Y.; Kyo, S.; Enoki, C.; Fujiwara, H.; Sumida, T.; Hattori, R.; Imamura, H. Role of F-actin organization in p38 MAPkinase-mediated apoptosis and necrosis in neonatal rat cardiomyocytes subjected to simulated ischemia and reoxygenation. *Am. J. Physiol. Heart Circ. Physiol.*, **2005**, *289*, H2310-H2318.
- [151]Luschen, S.; Scherer, G.; Ussat, S.; Ungefroren, H.; Adam-Klages, S. Inhibition of p38 mitogen-activated protein kinase reduces TNF-induced activation of NF-κB, elicits caspase activity, and enhances cytotoxicity. *Exp. Cell Res.*, **2004**, *293*, 196-206
- [152]Straface, E.; Natalini, B.; Monti, D.; Franceschi, C.; Schettini, G.; Bisaglia, M.; Fumelli, C.; Pincelli, C.; Pellicciari, R.; Malorni, W. C₃-Fullerene-tris-methanodicarboxylic acid protects epithelial cells from radiation-induced anoikia by influencing cell adhesion ability. *FEBS Lett.*, **1999**, *454*, 335-340.
- [153]Johnson-Lyles, D.N.; Peifley, K.; Lockett, S.; Neun, B.W.; Hansen, M.; Clogston, J.; Stern, S.T.; McNeil, S.M. Fullerenol cytotoxicity in kidney cells is associated with cytoskeleton disruption, autophagic vacuole accumulation, and mitochondrial dysfunction. *Toxicol. Appl. Pharmacol.*, **2010**, *248*, 249-258.
- [154]Rebecca, M.; Hsing-Lin, W.; Jun, G.; Srinivas, I.; Gabriel, M.A.; Jennifer, M.; Andrew, S.P.; Yuping, B.; Chun-Chih, W.; Zhong, C.; Yuan, G.; Rashi, I. Impact of physicochemical properties of engineered fullerenes on key biological responses. *Toxicol. Appl. Pharmacol.*, **2009**, *234*, 58-67.
- [155]Zogovic, N.S.; Nikolic, N.S.; Vranjes-Djuric, S.D.; Harhaji, L.M.; Vucicevic, L.M.; Janjetovic, K.D.; Misirkic, M.S.; Todorovic-Markovic, B.M.; Markovic, Z.M.; Milonjic, S.K.; Trajkovic, V.S. Opposite effects of nanocrystalline fullerene (C₆₀) on tumour cell growth *in vitro* and *in vivo* and a possible role of immunosuppression in the cancer-promoting activity of C₆₀. *Biomaterials*, **2009**, *30*, 6940-6946.
- [156]Huang, Y.L.; Shen, C.K.F.; Luh, T.Y.; Yang, H.C.; Hwang, K.C.; Chou, C.K. Blockage of apoptotic signaling of transforming growth factor-beta in human hepatoma cells by carboxy-fullerene. *Eur. J. Biochem.*, **1998**, *254*, 38-43.
- [157]Hsu, S.C.; Wu, C.C.; Luh, T.Y.; Chou, C.K.; Han, S.H.; Lai, M.Z. Apoptotic signal of Fas is not mediated by ceramide. *Blood*, **1998**, *91*, 2658-2663.
- [158]Li, W.; Zhao, L.; Wei, T.; Zhao, Y.; Chen, C. The inhibition of death receptor mediated apoptosis through lysosome stabilization following internalization of carboxyfullerene nanoparticles. *Biomaterials*, **2011**, *32*, 4030-4041.
- [159]Nakagawa, Y.; Suzuki, T.; Ishii, H.; Nakae, D.; Ogata, A. Cytotoxic effects of hydroxylated fullerenes on isolated rat hepatocytes via mitochondrial dysfunction. *Arch. Toxicol.*, **2011**, *85*, 1429-1440.
- [160]Rouse, J.G.; Yang, J.Z.; Barron, A.R.; Monteiro-Riviere, N.A. Fullerene-based amino acid nanoparticle interactions with human epidermal keratinocytes. *Toxicol. in Vitro*, **2006**, *20*, 1313-1320.
- [161]Huang, S.T.; Liao, J.S.; Fang, H.W.; Lin, C.M. Synthesis and anti-inflammation evaluation of new C₆₀fulleropyrrolidines bearing biologically active xanthine. *Bioorg. Med. Chem. Lett.*, **2008**, *18*, 99-103.
- [162]Gelderman, M.P.; Simakova, O.; Clogston, J.D.; Patri, A.K.; Siddiqui, S.F.; Vostal, A.C.; Simak, J. Adverse effects of fullerenes on endothelial cells: Fullerenol C₆₀(OH)₂₄ induced tissue factor and ICAM-1 membrane expression and apoptosis *in vitro*. *Int. J. Nanomed.*, **2008**, *3*, 59-68.
- [163]Zhu, J.; Ji, Z.; Wang, J.; Sun, R.; Zhang, X.; Gao, Y.; Sun, H.; Liu, Y.; Wang, Z.; Li, A.; Ma, J.; Wang, T.; Jia, G.; Gu, Y. Tumor-inhibitory effect and immunomodulatory activity of fullerol C₆₀(OH)_x. *Small*, **2008**, *4*, 1168-1175.

- [164] Harhaji, L.; Isakovic, A.; Vucicevic, L.; Janjetovic, K.; Misirkic, M.; Markovic, Z.; Todorovic-Markovic, B.; Nikolic, N.; Vranjes-Djuric, S.; Nikolic, Z.; Trajkovic, V. Modulation of tumor necrosis factor-mediated cell death by fullerenes. *Pharm. Res.*, **2008**, *25*, 1365-1376.
- [165] Liu, R.L.; Cai, X.Q.; Wang, J.D.; Li, J.G.; Huang, Q.; Li, W.X. Research on the bioactivities of C₆₀-dexamethasone. *J. Nanosci. Nanotechnol.*, **2009**, *9*, 3171-3176.
- [166] Yang, D.; Zhao, Y.; Guo, H.; Li, Y.; Tewary, P.; Xing, G.; Hou, W.; Oppenheim, J.J.; Zhang, N. [Gd@C82(OH)22]*n* Nanoparticles induce dendritic cell maturation and activate Th1 immune responses. *ACS Nano*, **2010**, *4*, 1178-1186.
- [167] Misirkic, M.S.; Todorovic-Markovic, B.M.; Vucicevic, L.M.; Janjetovic, K.D.; Jokanovic, V.R.; Dramicanin, M.D.; Markovic, Z.M.; Trajkovic, V.S. The protection of cells from nitric oxide-mediated apoptotic death by mechanochemically synthesized fullerene (C₆₀) nanoparticles. *Biomaterials*, **2009**, *30*, 2319-2328.
- [168] Xu, A.; Chai, Y.; Nohmi, T.; Hei, T.K. Genotoxic responses to titanium dioxide nanoparticles and fullerene in *gpt* delta transgenic MEF cells. *Part. Fibre Toxicol.*, **2009**, *6*, 3-16.
- [169] Ciacci, L.C.C.; Vallottoc, D.; Galloa, G.; Marcominic, A.; Pojanac, G. *In vitro* effects of suspensions of selected nanoparticles (C₆₀ fullerene, TiO₂, SiO₂) on *Mytilus hemocytes*. *Aquatic Toxicol.*, **2010**, *96*, 151-158.
- [170] Fiorito, S.; Serafino, A.; Andreola, F.; Bernier, P. Effects of fullerenes and single-wall carbon nanotubes on murine and human macrophages. *Carbon*, **2006**, *44*, 1100-1105.
- [171] Hu, Z.; Huang, Y.; Guan, W.; Zhang, J.; Wang, F.; Zhao, L. The protective activities of water-soluble C₆₀ derivatives against nitric oxide-induced cytotoxicity in rat pheochromocytoma cells. *Biomaterials*, **2010**, *31*, 8872-8881.
- [172] Huang, S.S.; Tsai, S.K.; Chin, C.L.; Chiang, L.Y.; Hsieh, H.M.; Teng, C.M.; Tsai, M.C. Neuroprotective effect of hexasulfobutylated C₆₀ on rats subjected to focal cerebral ischemia. *Free Rad. Biol. Med.*, **2001**, *30*, 643-649.
- [173] Li, Y.; Liu, Y.; Fu, Y.; Wei, T.; Guyader, L.L.; Gao, G.; Liu, R.S.; Chang, Y.Z.; Chen, C. The triggering of apoptosis in macrophages by pristine graphene through the MAPK and TGF-beta signaling pathways. *Biomaterials*, **2012**, *33*, 402-411.
- [174] Kotelnikova, R.A.; Kotelnikov, A.I.; Bogdanov, G.N.; Romanova, V.S.; Kuleshova, E.F.; Parnes, Z.N. Membranotropic properties of the water soluble amino acid and peptide derivatives of fullerene C₆₀. *FEBS Lett.*, **1996**, *389*, 111-114.
- [175] Chen, Y.W.; Hwang, K.C.H.; Yen, C.C.; Lai, Y.L. Fullerene derivatives protect against oxidative stress in RAW 264.7 cells and ischemia-reperfused lungs. *Am. J. Physiol.*, **2004**, *287*, 21-26.
- [176] Chien, C.T.; Chen, C.F.; Hsu, S.M.; Chiang, L.Y.; Lai, M.K. Forced expression of bcl-2 and bcl-xL by novel water-soluble fullerene, C₆₀(glucosamine)₆, reduces renal ischemia/reperfusion-induced oxidative stress. *Fuller Nanotub Car.*, **2001**, *9*, 77-88.
- [177] Bisaglia, M.; Natalini, B.L.; Pellicciari, R.; Straface, E.; Malorni, W.; Monti, D. Carboxyfullerenes as neuroprotective agents C₃-fullerotrimer-methanodicarboxylic acid protects cerebellar granule cells from apoptosis. *J. Neurochem.*, **2000**, *74*, 1197-1204.
- [178] Nel, A.; Xia, T.; Madler, L.; Li, N. Toxic potential of materials at the nanolevel. *Science*, **2006**, *311*, 622-627.
- [179] Maynard, A.D.; Aitken, R.J.; Butz, T.; Colvin, V.; Donaldson, K.; Oberdorster, G.; Philbert, M.A.; Ryan, J.; Seaton, A.; Stone, V. Safe Handling of Nanotechnology. *Nature*, **2006**, *444*, 267-269.
- [180] Calvaresi, M.; Zerbetto, F. Baiting Proteins with C₆₀. *ACS Nano*. **2010**, *4*, 2283-2299.
- [181] Da Ros, T.; M.Prato, M.; F.Novello, F.; M.Maggini, M.; E.Banfi, E. Easy access to water-soluble fullerene derivatives via 1,3-dipolar cycloadditions of azomethine ylides to C₆₀. *J. Org. Chem.*, **1996**, *61*, 9070-9072.
- [182] Jin, H.; Chen, W.Q.; Tang, X.W.; Chiang, L.Y.; Yang, C.Y.; Schloss, J.V.; Wu, J.Y. Polyhydroxylated C₆₀, Fullerenols, as glutamate receptor antagonists and neuroprotective agents. *J. Neurosci. Res.*, **2000**, *62*, 600-607.
- [183] Boutorine, A.S.; Tokuyama, H.; Takasugi, M.; Isobe, H.; Nakamura, E.; Helene, C. Fullerene-oligonucleotide conjugates: photoinduced sequence-specific DNA cleavage. *Angew. Chem. Int. Ed.*, **1994**, *33*, 2426-2465.
- [184] Kim, J.E.; Lee, M. Fullerene inhibits β -amyloid peptide aggregation. *Biochem. Biophys. Res. Commun.*, **2003**, *303*, 576-579.
- [185] Gupta, S.; Dhawan, A.; Shanker, R. *In silico* approaches: prediction of biological targets for fullerene derivatives. *J. Biomed. Nanotechnol.*, **2011**, *7*, 91-92.
- [186] Wolff, D.J.; Barbieri, C.M.; Richardson, C.F.; Schuster, D.I.; Wilson, S.R. Trisamine C₆₀-fullerene adducts inhibit neuronal nitric oxide synthase by acting as highly potent calmodulin antagonists. *Arch. Biochem. Biophys.*, **2002**, *399*, 130-141
- [187] Mashino, T.; Okuda, K.; Hirota, T.; Hirobe, M.; Nagano, T.; Mochizuki, M. Inhibitory effect of fullerene derivatives on glutathione reductase. *Fullerene Sci. Technol.*, **2001**, *9*, 191-196.
- [188] Marcorin, G.L.; Da Ros, T.; Castellano, S.; Stefancich, G.; Bonin, I.; Miertus, S.; Prato, M. Design and synthesis of novel [60]fullerene derivatives as potential HIV aspartic protease inhibitors. *Org. Lett.*, **2000**, *2*, 3955-3958.
- [189] Schuster, D.I.; Wilson, S.R.; Schinazi, R.F. Anti-human immunodeficiency virus activity and cytotoxicity of derivatized Buckminsterfullerenes. *Bioorg. Med. Chem. Lett.*, **1996**, *6*, 1253-1256..

- [190] Braden, B.C.; Goldbaum, F.A.; Chen, B.X.; Kirschner, A.N.; Wilson, S.R.; Erlanger, B.F. X-Ray crystal structure of an anti-Buckminsterfullerene antibody fab fragment: biomolecular recognition of C₆₀. *Proc. Natl. Acad. Sci. U.S.A.*, **2000**, *97*, 12193-12197.
- [191] Rozhkov, S.P.; Goryunov, A.S.; Sukhanova, G.A.; Borisova, A.G.; Rozhkova, N.N.; Andrievsky, G.V. Protein interaction with hydrated C₆₀ fullerene in aqueous solutions. *Biochem. Biophys. Res. Commun.*, **2003**, *303*, 562-566.
- [192] Belgorodsky, B.; Fadeev, L.; Kolsenik, J.; Gozin, M. Formation of a soluble stable complex between pristine C₆₀-fullerene and a native blood protein. *ChemBioChem.*, **2006**, *7*, 1783-1789.
- [193] Belgorodsky, B.; Fadeev, L.; Ittah, V.; Benyamini, H.; Zelnor, S.; Huppert, D.; Kotlyar, A.B.; Gozin, M. Formation and characterization of stable human serum albumin-tris-malonic acid [C₆₀]fullerene complex. *Bioconjugate Chem.*, **2005**, *16*, 1058-1062.
- [194] Yang, S.T.; Wang, H.; Guo, L.; Gao, Y.; Liu, Y.; Cao, A. Interaction of fullereneol with lysozyme investigated by experimental and computational approaches. *Nanotechnology*, **2008**, *19*, 395-401.
- [195] Benyamini, H.; Shulman-Peleg, A.; Wolfson, H.J.; Belgorodsky, B.; Fadeev, L.; Gozin, M. Interaction of C₆₀-fullerene and carboxyfullerene with proteins: docking and binding site alignment. *Bioconjugate Chem.*, **2006**, *17*, 378-386.
- [196] Zhang, X.F.; Shu, C.Y.; Xie, L.; Wang, C.; Zhang, Y.Z.; Xiang, J.F.; Li, L.; Tang, Y.L. Protein conformation changes induced by a novel organophosphate-containing water-soluble derivative of a C₆₀ fullerene nanoparticle. *J. Phys. Chem. C*, **2007**, *111*, 14327-14333.
- [197] Belgorodsky, B.; Fadeeva, L.; Kolsenik, J.; Gozin, M. Biodelivery of a fullerene derivative. *Bioconjugate Chem.*, **2007**, *18*, 1095-1100.
- [198] Ueng, T.H.; Kang, J.J.; Wang, H.W.; Cheng, Y.W.; Chiang, L.Y. Suppression of microsomal cytochrome P450-Dependent monooxygenases and mitochondrial oxidative phosphorylation by fullereneol, a polyhydroxylated fullerene C₆₀. *Toxicol. Lett.*, **1997**, *93*, 29-37.
- [199] Pastorin, G.; Marchesan, S.; Hoebeke, J.; Da Ros, T.; Ehret-Sabatier, L.; Briand, J.P.; Prato, M.; Bianco, A. Design and activity of cationic fullerene derivatives as inhibitors of acetylcholinesterase. *Org. Biomol. Chem.*, **2006**, *4*, 2556-2562.
- [200] Innocenti, A.; Durdagi, S.; Doostdar, N.; Strom, T.A.; Barron, A.R.; Supuran, C.T. Nanoscale enzyme inhibitors: fullerenes inhibit carbonic anhydrase by occluding the active site entrance. *Bioorg. Med. Chem.*, **2010**, *18*, 2822-2828.
- [201] Iwata, N.; T. Mukai, T.; Y.N. Yamakoshi, Y.N.; S.Hara, S.; T. Yanase, T.; M. Shoji, M.; T. Endo, T.; N. Miyata, N. Effect of C₆₀, a fullerene, on the activities of glutathione S-transferase and glutathione-related enzymes. *Fullerenes, Nanotubes & Carbon Nanostructures*, **1998**, *6*, 213-226.
- [202] Marczak, R.; Hoang, V.T.; Noworyta, K.; Zandler, M.E.; Kutner, W.; D'Souza, F. Molecular recognition of adenine, adenosine and ATP at the air-water interface by a uracil appended fullerene. *J. Mater. Chem.*, **2002**, *12*, 2123-2129.
- [203] Ito, M.; Nakashima, N. Design, synthesis and photophysical properties of C₆₀-modified proteins. *J. Mater. Chem.*, **2002**, *12*, 2026-2033.
- [204] Szeltukhin, A.O.; Chumakov, P.M. Casual and induced functions of p53 gene. *Success of Biol. Chem. (Russ.)*, **2010**, *50*, 447-516.
- [205] Liang, Y.; Luo, F.; Lin, Y.; Zhou, Q.F.; Jiang, G.B. C₆₀ affects DNA replication *in vitro* by decreasing the melting temperature of DNA templates. *Carbon*, **2009**, *47*, 1457-1465.
- [206] Kang, F.; Song, G.G. Inhibition of Taq DNA polymerase and DNA exonuclease ExoIII by an aqueous nanoparticle suspension of a bis-methanophosphonate fullerene. *Mater. Sci. Forum*, **2011**, *685*, 345-351.
- [207] An, H.; Jin, B. DNA exposure to buckminsterfullerene (C₆₀): toward DNA stability, reactivity, and replication. *Environ. Sci. Technol.*, **2011**, *45*, 6608-6616.
- [208] Rohs, R.; West, S.M.; Sosinsky, A.; Liu, P.; Mann, R.S.; Honig, B. The role of DNA shape in protein-DNA recognition. *Nature*, **2009**, *461*, 1248-1253.
- [209] Pinteala, M.; Dascalu, A.; Ungurenasu, C. Binding fullereneol C₆₀(OH)₂₄ to dsDNA. *Int. J. Nanomed.*, **2009**, *4*, 193-199.
- [210] Li, J.; Zhang, M.; Sun, B.; Xing, G.; Song, Y.; Guo, H.; Chang, Y.; Ge, Y.; Zhao, Y. Separation and purification of fullereneols for improved biocompatibility. *Carbon*, **2012**, *50*, 460-469.
- [211] Tanimoto, S.; Sakai, S.; Matsumura, S.; Takahashi, D.; Toshima, K. Target-selective photo-degradation of HIV-1 protease by a fullerene-sugar hybrid. *Chem. Commun.*, **2008**, 5767-5769.
- [212] Durdagi, S.; Mavromoustakos, T.; Chronakis, N.; Papadopoulos, M.G. Computational design of novel fullerene analogues as potential HIV-1 PR inhibitors: Analysis of the binding interactions between fullerene inhibitors and HIV-1 PR residues using 3D QSAR, molecular docking and molecular dynamics simulations. *Bioorg. Med. Chem.*, **2008**, *16*, 9957-9974.
- [213] Bosi, S.; Da Ros, T.; Spalluto, G.; Balzarini, J.; Prato, M. Synthesis and anti-HIV properties of new water-soluble bis-functionalized [60]fullerene derivatives. *Bioorg. Med. Chem. Lett.*, **2003**, *13*, 4437-4440.

- [214] Marchesan, S.; Da Ros, T.; Spalluto, G.; Balzarini, J.; Prato, M. Anti-HIV properties of cationic fullerene derivatives. *Bioorg. Med. Chem. Lett.*, **2005**, *15*, 3615–3618.
- [215] Mashino, T.; Shimotohno, K.; Ikegami, N.; Nishikawa, D.; Okuda, R.; Takahashi, R.; Nakamura, S.; Mochizuki, M. Human immunodeficiency virus-reverse transcriptase inhibition and hepatitis C virus RNA-dependent RNA polymerase inhibition activities of fullerene derivatives. *Bioorg. Med. Chem. Lett.*, **2005**, *15*, 1107–1109.
- [216] Ungurenasu, C.; Pinteala, M. Syntheses and characterization of water-soluble C₆₀-Curdlan sulfates for biological applications. *J. Polym. Sci. Polym. Chem.*, **2007**, *45*, 3124–3128.
- [217] Lehtovaara, B.C.; Gu, F.X. Pharmacological, structural, and drug delivery properties and applications of 1,3-β-glucans. *J. Agric. Food Chem.*, **2011**, *59*, 6813–6828.
- [218] Shimanovsky, N.S. Nanotechnologies in contemporary pharmacology. *Int. Med. J. (Russ.)*, **2009**, №1, 131–135.
- [219] Fujita, K.; Morimoto, Y.; Ogami, A.; Myojyo, T.; Tanaka, I.; Shimada, M.; Wang, W.; Endoh, S.; Uchida, K.; Nakazato, T.; Yamamoto, K.; Fukui, H.; Horie, M.; Yoshida, Y.; Iwahashi, H.; Nakanishi, J. Gene expression profiles in rat lung after inhalation exposure to C₆₀ fullerene particles. *Toxicology*, **2009**, *258*, 47–55.
- [220] Fujita, K.; Morimoto, Y.; Endoh, S.; Uchida, K.; Fukui, H.; Ogami, A.; Tanaka, I.; Horie, M.; Yoshida, Y.; Iwahashi, H.; Nakanishi, J. Identification of potential biomarkers from gene expression profiles in rat lungs intratracheally instilled with C₆₀ fullerenes. *Toxicology*, **2010**, *274*, 34–41.
- [221] Xia, T.; Kovochich, M.; Brant, J.; Hotze, M.; Sempf, J.; Oberley, T.; Sioutas, C.; Yeh, J.I.; Wiesner, M.R.; Nel, A.N. Comparison of the abilities of ambient and manufactured nanoparticles to induce cellular toxicity according to an oxidative stress paradigm. *Nano Lett.*, **2006**, *6*, 1794–1807.
- [222] Jacobsen, N.R.; Pojana, G.; White, P.; Møller, P.; Cohn, C.A.; Korsholm, K.S.; Vogel, U.; Marcomini, A.; Loft, S.; Wallin, H. Genotoxicity, cytotoxicity, and reactive oxygen species induced by single-walled carbon nanotubes and C₆₀ fullerenes in the FE1-Muta™ Mouse lung epithelial cells. *Environ. Mol. Mutagen.*, **2008**, *49*: 476–487.
- [223] Jovanovic, B.; Ji, T.; Palic, D. Gene expression of zebrafish embryos exposed to titanium dioxide nanoparticles and hydroxylated fullerenes. *Ecotoxicol. Environ. Safety*, **2011**, *74*, 1518–1525.
- [224] Park, E.J.; Kim, H.; Kim, Y.; Yi, J.; Choi, K.; Park, K. Carbon fullerenes (C₆₀s) can induce inflammatory responses in the lung of mice. *Toxicol. Appl. Pharmacol.*, **2010**, *244*, 226–233.
- [225] Kaidashev, I.P. Influence of C₆₀ fullerene on activity of phagocytic cells. *Exp. Clinic. Pharmacol. (Russ.)*, **2011**, #6, 26–29
- [226] Xu, Y.; Zhu, J.; Xiang, K.; Li, Y.; Sun, R.; Ma, J.; Sun, H.; Liu, Y. Synthesis and immunomodulatory activity of [60]fullerene-tuftsin conjugates. *Biomaterials*, **2011**, *32*(36), 9940–9949.
- [227] Vengerovich, N.G.; Tunin, M.A.; Antonenkova, E.V.; Konshakov, Yu.A.; Bolekhan, A.V.; Zaitseva, O.B.; Stukov, A.N.; Boyarkin, M.N.; Popov, V.A.. Biological activity of nanobiocomposites fullerene C₆₀. *Immunology (Russ)*, **2011**, *12*, 161–177.
- [228] Romoser, A.A.; Figueroa, D.E.; Soresh, A.; Scribner, K.; Chen, P.L.; Porter, W.; Criscitiello, M.F.; Sayes, C.M. Distinct immunomodulatory effects of a panel of nanomaterials in human dermal fibroblasts. *Toxicol. Lett.*, **2011**, *6*, 7–34.
- [229] Mrdanovic, J.; Solajic, S.; Bogdanovic, V.; Stankov, K.; Bogdanovic, G.; Djordjevic, A. Effects of fullereneol C₆₀(OH)₂₄ on the frequency of micronuclei and chromosome aberrations in CHO-K1 cells. *Mutation Res.*, **2009**, *680*, 25–30.
- [230] Nielsen, G.D.; Roursgaard, M.; Jensen, K.A.; Poulsen, S.S.; Larsen, S.T. Biology and toxicology of fullerenes and their derivatives. *Basic Clin. Pharmacol. Toxicol.*, **2008**, *103*, 197–208.
- [231] Jalbout, A.F.; Hameed, A.J.; Trzaskowski, B. Study of the structural and electronic properties of 1-(4, 5 and 6-selenenyl derivatives-3-formyl-phenyl) pyrrolidinofullerenes, *J. Organometal. Chem.*, **2007**, *692*, 1039–1047.
- [232] Jiao, F.; Liu, Y.; Qu, Y.; Li, W.; Zhou, G.; Ge, C.; Li, Y.; Sun, B.; Chen, C. Studies on anti-tumor and antimetastatic activities of fullereneol in a mouse breast cancer model, *Carbon*, **2010**, *48*, 2231–2243.
- [233] Nishizawa, C.; Hashimoto, N.; Yokoo, S.; Funakoshi-Tago, M.; Kasahara, T.; Takahashi, K.; Nakamura, S.; Mashino, T. Pyrrolidinium-type fullerene derivative-induced apoptosis by the generation of reactive oxygen species in HL-60 cells, *Free Radic. Res.*, **2009**, *43*, 1240–1247.
- [234] Wang, J.; C.Chen, C.; B.Li, B.; H.Yu, H.; Y.Zhao, Y.; J.Sun, J.; Y.Li, Y.; G.Xing, G.; H.Yuan, H.; J.Tang, J.; Z.Chen, Z.; H.Meng, H.; Y.Gao, Y.; C.Ye, C.; Z.Chai, Z.; C.Zhu, C.; B.Ma, B.; X.Fang, X.; L.Wanc, L. Antioxidative function and biodistribution of [Gd@C₈₂(OH)₂₂]_n nanoparticles in tumor-bearing mice, *Biochem. Pharmacol.*, **2006**, *71*, 872–881.
- [235] Liu, Y.; Jiao, F.; Qiu, Y.; Li, W.; Qu, Y.; Li, Y. Immunostimulatory properties and enhanced TNF-α mediated cellular immunity for tumor therapy by C₆₀(OH)₂₀ nanoparticles, *Nanotechnology*, **2009**, *20*, 415102–415111.
- [236] Liu, Y.; Jiao, F.; Qiu, Y.; Li, W.; Lao, F.; Zhou, G.Q. The effect of Gd@C₈₂(OH)₂₂ nanoparticles on the release of Th1/Th2 cytokines and induction of TNF-α mediated cellular immunity, *Biomaterials*, **2009**, *30*, 3934–3945.

- [237] Meng, H.; Xing, G.; Sun, B.; Zhao, F.; Lei, H.; Li, W.; Song, Y.; Chen, Z.; Yuan, H.; Wang, X.; Long, J.; Chen, C.; Liang, X.; Zhang, N.; Chai, Z.; Zhao, Y. Potent angiogenesis inhibition by the particulate form of fullerene derivatives. *ACS Nano*, **2010**, *4*, 2773-2783.
- [238] Darwish, A.D. Fullerenes, *Annu. Rep. Prog. Chem., A*, **2009**, *105*, 363-381.
- [239] Prylutska, S.; Burlaka, A.P.; Klymenko, P.P.; Grynyuk, I.I.; Prylutsky, Yu.I.; Schütze, C.; Ritter, U. Using water-soluble C₆₀ fullerenes in anticancer therapy, *Cancer Nanotechnol.*, **2011**, *2*, 105-110.
- [240] Kharisov, B.I.; Kharissova, O.V.; Gomez, M.J.; Mendez, U.O. Recent advances in the synthesis, characterization, and applications of fulleropyrrolidines, *Ind. Eng. Chem. Res.*, **2009**, *48*, 545-571.
- [241] Darwish, A.D. Fullerenes, *Annu. Rep. Prog. Chem., A*, **2007**, *103*, 370-391.
- [242] Mikawa, M.; Kato, H.; Okumura, M.; Narazaki, M.; Kanazawa, Y.; Miwa, N. Paramagnetic water-soluble metallofullerenes having the highest relaxivity for MRI contrast agents. *Bioconjug. Chem.*, **2001**, *12*, 510-514.
- [243] Li, Q.; Xiu, Y.; Zhang, X.; Liu, R.; Du, Q.; Sun, X.; Chen, S.; Li, W. Biodistribution of fullerene derivative C₆₀(OH)_x(O)_y, *Chinese Sci. Bull.*, **2001**, *46*, 1615-1617.
- [244] Liu, J.H.; Cao, L.; Luo, P.G.; Yang, S.T.; Lu, F.; Wang, H.; Meziani, M.J.; Haque, S.A.; Liu, Y.; Lacher, S.; Sun, Y.P. Fullerene-conjugated doxorubicin in cells, *ACS Appl. Mater. Interf.*, **2010**, *2*, 1384-1389.
- [245] Wei, P.; Zhang, L.; Lu, Y.; Man, N.; Wen, L. C₆₀(Nd) nanoparticles enhance chemotherapeutic susceptibility of cancer cells by modulation of autophagy, *Nanotechnol.*, **2010**, *21*, 495101.
- [246] Montellano, A.; Da Ros, T.; Bianco, A.; Prato, M. Fullerene C₆₀ as a multifunctional system for drug and gene delivery, *Nanoscale*, **2011**, *3*, 4035-4044.
- [247] Rancan, F.; Helmreich, M.; Molich, A.; Jux, N.; Hirsch, A.; Roder, B.; Witt, C.; Bohm, F. Fullerene-porphyrin complexes as sensitizer for photodynamic therapy: uptake and photo-induced cytotoxicity on Jurkat cells, *J. Photochem. Photobiol. B*, **2005**, *80*, 1-7.
- [248] Boyd, P.D.; Hodgson, M.C.; Rickard, C.E.F.; Oliver, A.G.; Chaker, L.; Brothers, P.J.; Bolskar, R.D.; Tham, F.S.; Reed, C.A. Selective supramolecular porphyrin/fullerene interactions, *J. Am. Chem. Soc.* **1999**, *121*, 10487-10495.
- [249] Sun, Y.; Drovetskaya, T.; Bolskar, R.D.; Bau, R.; Boyd, P.D.; Reed, C.A. Fullerides of pyrrolidine-functionalized C₆₀, *J. Org. Chem.*, **1997**, *62*, 3642-3649.
- [250] Imahori, H.; Mori, Y.; Matano, Y. Nanostructured artificial photosynthesis, *J. Photochem. Photobiol., C*, **2003**, *4*, 51-83.
- [251] Da Ros, T.; Prato, M.; Guldi, D.M.; Ruzzi, M.; Pasimeni, L. Efficient charge separation in porphyrin-fullerene-ligand complexes, *Chemistry*, **2001**, *7*, 816-827.
- [252] Nishiyama, N.; Stapert, H.R.; Zhang, G.D.; Takasu, D.; Jiang, D.L.; Nagano, T.; Aida, T.; Kataoka, K. Light-harvesting ionic dendrimer porphyrins as new photosensitizers for photodynamic therapy, *Bioconjugate Chem.*, **2003**, *14*, 58-66.
- [253] Dudic, M.; Lhotak, P.; Stibor, I.; Petrickova, H.; Lang, K. (Thia)calyx[4]arene-porphyrin conjugates: novel receptors for fullerene complexation with C₇₀ over C₆₀ selectivity, *New J. Chem.*, **2004**, *28*, 85-90.
- [254] Hirsch, A. *The Chemistry of Fullerenes*. Thieme-Verlag, Stuttgart; New York, **1994**.
- [255] Rezayat, S.M.; Boushehri, S.V.S.; Salmanian, B.; Omidvari, A.H.; Tarighat, S.; Esmaeili, S.; Sarkar, S.; Amirshahi, N.; Alyautdin, R.N.; Orlova, M.A.; Trushkov, I.V.; Buchachenko, A.L.; Liu, K.C.; Kuznetsov, D.A. The porphyrin-fullerene nanoparticles to promote the ATP overproduction in myocardium: ²⁵Mg²⁺-magnetic isotope effect, *Eur. J. Med. Chem.*, **2009**, *44*, 1554-1569.
- [256] Buchachenko, A.L.; Kuznetsov, D.A.; Breslavskaya, N.N.; Orlova, M.A. Magnesium Isotope Effect in Enzymatic Phosphorylation, *J. Phys. Chem. B*, **2008**, *112*, 2548-2556.
- [257] Nikolic, N.; Vranjes-Ethuric, S.; Jankovic, D.; Ethokic, D.; Mirkovic, M.; Bibic, N.; Trajkovic, V. Preparation and biodistribution of radiolabeled fullerene C₆₀ nanocrystals, *Nanotechnol.*, **2009**, *20*, 385102.
- [258] Ikeda, A.; Doi, Y.; Nishiguchi, K.; Kitamura, K.; Hashizume, M.; Kikuchi, J.; Yogo, K.; Ogawa, T.; Takey, T. Induction of cell death by photodynamic therapy with water-soluble lipid-membrane-incorporated [60]fullerene, *Org. Biomol. Chem.*, **2007**, *5*, 1158-1160.
- [259] Wang, H.; Wang, L.; Wang, X.; Xu, J.; Luo, Q.; Liu, J. Self-assembled nanostructures from C₆₀-containing supramolecular complex: its stimuli-responsive reversible transition and biological antioxidative capacity, *New J. Chem.*, **2011**, *35*, 2632-2638.
- [260] Akiyama, M.; Ikeda, A.; Shintani, T.; Doi, Y.; Kikuchi, J.; Ogawa, T.; Yogo, K.; Takeya, T.; Yamamoto, N. Solubilisation of [60]fullerenes using block copolymers and evaluation of their photodynamic activities, *Org. Biomol. Chem.*, **2008**, *6*, 1015-1019.
- [261] Zhou, Z.; Lenk, R.P.; Dellinger, A.; Wilson, S.R.; Sadler, R.; Kepley, C.L. Liposomal formulation of amphiphilic fullerene antioxidants. *Bioconj. Chem.* **2010**, *21*, 1656-1661.
- [262] Chaudhuri, P.; Paraskar, A.; Soni, S.; Mashelkar, R.A.; Sengupta, S. Fullerenol-cytotoxic conjugates for cancer chemotherapy, *ACS Nano*, **2009**, *3*, 2505-2514.
- [263] Song, H.; Luo, S.; Wei, H.; Song, H.; Yang, Y.; Zhao, W. *In vivo* biological behavior of ^{99m}Tc(CO)₃ labelled fullerol, *J. Radioanal. Nucl. Chem.*, **2010**, *285*, 635-639.

- [264]Roberts, J.E.; Wielgus, A.R.; Boyes, W.K.; Andley, U.; Chignell, C.F. Phototoxicity and cytotoxicity of fullerol in human lens epithelial cells, *Toxicol. Appl. Pharmacol.*, **2008**, *228*, 49-58.
- [265]Zhao, B.; Yin, J.J.; Bilski, P.; Chignell, C.F.; Roberts, J.E.; He, Y.Y. Enhanced photodynamic efficacy towards melanoma cells by encapsulation of Pc4 in silica nanoparticles, *Toxicol. Appl. Pharmacol.*, **2009**, *241*, 163-172.
- [266]Prow, T.W.; Bhutto, I.; Kim, S.Y.; Grebe, R.; Merges, C.; McLeod, D.S.; Uno, K.; Mennon, M.; Rodriguez, L.; Leong, K.; Luty, G.A. Ocular nanoparticle toxicity and transfection of the retina and retinal pigment epithelium, *Nanomedicine*, **2008**, *4*, 340-349.
- [267]Bejjani, R.A.; BenEzra, D.; Cohen, H.; Rieger, J.; Andrieu, C.; Jeanny, J.C.; Gollomb, G.; Behar-Cohen, F.F. Nanoparticles for gene delivery to retinal pigment epithelial cells, *Mol. Vis.*, **2005**, *17*, 124-132.
- [268]Yang, X.Y.; Edelman, R.E.; Oris, J.T. Suspended C₆₀ nanoparticles protect against short-term UV and fluoranthene photo-induced toxicity, but cause long-term cellular damage in *Daphnia magna*, *Aquatic Toxicol.*, **2010**, *100*, 202-210.
- [269]Mroz, P.; Tegos, G.P.; Gali, H.; Wharton, T.; Sarna, T.; Hamblin, M.R. Photodynamic therapy with fullerenes, *Photochem. Photobiol. Sci.*, **2007**, *6*, 1139-1149.
- [270]Burlaka, A.P.; Sidorik, U.P.; Prylutska, S.V. Catalytic system of the reactive oxygen species on the C₆₀ fullerene basis, *Exp. Oncol.*, **2004**, *26*, 326-327.
- [271]Meshalkin, Y.P.; Bgatova, N.P. Prospects and problems of the use of inorganics nanoparticles in oncology, *J. Siber. Fed. Univ. B. (Russ.)* **2008**, *3*, 248-268.
- [272]Mroz, P.; Pawlak, A.; Satti, M. Functionalized fullerenes mediate photodynamic killing of cancer cells: Type I versus Type II photochemical mechanism, *Free Radical Biol. Med.*, **2007**, *43*, 711-719.
- [273]Yano, S.; Hirohara, S.; Obata, M.; Hagiya, Y.; Ogura, S.; Ikeda, A.; Kataoka, H.; Tanaka, M.; Joh, T. . Current states and future views in photodynamic therapy, *J. Photochem. Photobiol. C*, **2011**, *12*, 46- 67.
- [274]Hu, Z.; Zhang, C.; Huang, Y.; Sun, S.; Guan, W.; Yao Y. Photodynamic anticancer activities of water-soluble C₆₀ derivatives and their biological consequences in a HeLa cell line, *Chem.-Biol. Inter.*, **2012**, *195*, 86-94.
- [275]Ion, R.M.; Fierascu, R.C.; Neagu, M.; Constantin, C.; Stavaru, C. Porphyrin (TPP)-polyvinylpyrrolidone (PVP)-fullerene (C₆₀) triad as novel sensitizer in photodynamic therapy, *Sci. Adv. Mater.*, **2010**, *2*, 223-229.
- [276]Liao, F.; Saitoh, Y.; Miwa, N. Anticancer effects of fullerene [C₆₀] included in polyethylene glycol combined with visible light irradiation through ROS generation and DNA fragmentation on fibrosarcoma cells with scarce cytotoxicity to normal fibroblasts, *Oncol. Res. Feat. Preclin. Clin. Cancer Therap.*, **2011**, *19*, 203-216.
- [277]Liu, J.; Ohta, S.; Sonoda, A. Preparation of PEG-conjugated fullerene containing Gd³⁺ ions for photodynamic therapy, *J. Controlled Release*, **2007**, *117*, 104-110.
- [278]Jiang, G.; Li, G. Preparation, characterization, and properties of fullerene-vinylpyrrolidone copolymers, *Biotechnol. Prog.*, **2012**, *28*, 215-222.
- [279]Vileno, B.; Jeney, S.; Sienkiewicz, A.; Marcoux, P.R.; Miller, L.M.; Forró, L. Evidence of lipid peroxidation and protein phosphorylation in cells upon oxidative stress photo-generated by fullerols, *Biophys. Chem.*, **2010**, *152*, 164-169.
- [280]Badireddy, A.R.; Hotze, E.M.; Chellam, S.; Alvarez, P.; Wiesner, M.R. Inactivation of bacteriophages via photosensitization of fullerol nanoparticles, *Environ. Sci. Technol.*, **2007**, *41*, 6627-6632.
- [281]Taroni, P.; D'Andrea, C.; Valentini, G.; Cubeddu, R.; Hu, D.N.; Roberts, J.E. Fullerol in human lens and retinal pigment epithelial cells: time domain fluorescence spectroscopy and imaging, *Photochem. Photobiol. Sci.*, **2011**, *10*, 904-910.
- [282]Vileno, B.; Marcoux, P.R.; Lekka, M.; Sienkiewicz, A.; Feher, T.; Forró, L. Spectroscopic and photophysical properties of a highly derivatized C₆₀ fullerol, *Adv. Funct. Mater.*, **2006**, *16*, 120-128.
- [283]Sienkiewicz, A.; Vileno, B.; Pierzchała, K.; Czuba, M.; Marcoux, P.R.; Graczyk, A.; Fajer, P.G.; Forró, L. . Oxidative stress-mediated protein conformation changes: ESR study of spin-labelled staphylococcal nuclease, *J. Phys. Condens. Matter*, **2007**, *19*, 285201
- [284]Straface, E.; Santini, M.T.; Donelli, G.; Giacomoni, P.U.; Malorni, W. Vitamin E prevents UVB-induced cell blebbing and cell death in A431 epidermoid cells, *Int. J. Rad. Biol.*, **1995**, *68*, 579-587.
- [285]Ito, S.; Itoga, K.; Yamato, M.; Akamatsu, H.; Okano, T. The co-application effects of fullerene and ascorbic acid on UV-B irradiated mouse skin, *Toxicology*, **2010**, *267*, 27-38.
- [286]Kato, S.; Kikuchi, R.; Aoshima, H.; Saitoh, Y.; Miwa, N. Defensive effects of fullerene-C₆₀/liposome complex against UVA-induced intracellular reactive oxygen species generation and cell death in human skin keratinocytes HaCaT, associated with intracellular uptake and extracellular excretion of fullerene-C₆₀, *J. Photochem. Photobiol. B.*, **2010**, *98*, 144-151.
- [287]Kato, S.; Aoshima, H.; Saitoh, Y.; Miwa, N. Fullerene-C₆₀/liposome complex: Defensive effects against UVA-induced damages in skin structure, nucleus and collagen type I/IV fibrils, and the permeability into human skin tissue, *J. Photochem. Photobiol. B*, **2010**, *98*, 99-105.
- [288]Fumelli, C.; Marconi, A.; Salvioli, S.; Straface, E.; Malorni, W.; Offidani, A.M.; Pellicciari, R.; Schettini, G.; Giannetti, A.; Monti, D.; Franceschi, C.; Pincelli, C. Carboxyfullerenes protect human keratinocytes from ultraviolet-B-induced apoptosis, *J. Invest. Dermatol.*, **2000**, *115*, 835-841.

- [289]Kato, S.; Ajshima, H.; Saitoh, Y.; Miwa, N. Fullerene-C₆₀ Incorporated in Liposome Exerts Persistent Hydroxyl Radical-Scavenging Activity and Cytoprotection in UVA/B-Irradiated Keratinocytes, *J. Nanosci. Nanotechnol.*, **2011**, *11*, 3814-3823.
- [290]Saitoh, Y.; Miyanishi, A.; Mizuno, H.; Kato, S.; Aoshima, H.; Kokubo, K.; Miwa, N. Super-highly hydroxylated fullerene derivative protects human keratinocytes from UV-induced cell injuries together with the decreases in intracellular ROS generation and DNA damages (2011), *J. Photochem. Photobiol. B*, **2011**, *102*, 69-76.
- [291]Zhao, B.; He, Y.Y.; Chignell, C.F.; Yin, J.J.; Andley, U.; Roberts, J.E. Difference in phototoxicity of cyclodextrin complexed fullerene [(γ -CyD)₂/C₆₀] and its aggregated derivatives toward human lens epithelial cells, *Chem. Res. Toxicol.*, **2009**, *22*, 660-667.
- [292]Doi, Y.; Ikeda, A.; Akiyama, M.; Nagano, M.; Shigematsu, T.; Ogawa, T.; Takeya, T.; Nagasaki, T. Intracellular uptake and photodynamic activity of water-soluble [60]- and [70]fullerenes incorporated in liposomes, *Chemistry: Eur. J.*, **2008**, *14*, 8892-8897.
- [293]Rancan, F.; Helmreich, M.; Mölich, A.; Jux, N.; Hirsch, A.; Röder, B.; Böhm, F. Intracellular Uptake and Phototoxicity of 3¹,3²-Didehydrophytychlorin-fullerene Hexaadducts, *Photochem. Photobiology*, **2007**, *83*, 1330-1338.
- [294]Alvarez, M.; Prucca, C.; Milanesio, M.E.; Durantini, E.N.; Rivarola, V. Photodynamic activity of a new sensitizer derived from porphyrin-C₆₀ dyad and its biological consequences in a human carcinoma cell line, *Int. J. Biochem. Cell Biol.*, **2006**, *38*, 2092-2101.
- [295]Mikata, Y.; Takagi, S.; Tanahashi, M.; Ishii, S.; Obata, M.; Miyamoto, Y.; Wakita, K.; Nishisaka, T.; Hirano, T.; Ito, T.; Hoshino, M.; Ohtsuki, C.; Tanihara, M.; Yano, S. Detection of 1270 nm emission from singlet oxygen and photocytotoxic property of sugar-pendant [60] fullerenes, *Bioorg. Med. Chem. Lett.*, **2003**, *13*, 3289-3292.
- [296]Otake, E.; Sakuma, S.; Torii, K.; Maeda, A.; Ohi, H.; Yano, S.; Morita, A. Effect and mechanism of a new photodynamic therapy with glycoconjugated fullerene, *Photochem. Photobiol.*, **2010**, *86*, 1356-1363.
- [297]Horie, M.; Fukuhara, A.; Saito, Y.; Yoshida, Y.; Sato, H.; Ohi, H.; Obata, M.; Mikata, Y.; Yano, S.; Niki, E. Antioxidant action of sugar-pendant C₆₀ fullerenes, *Bioorg. Med. Chem. Lett.*, **2009**, *19*, 5902-5904.
- [298]Lebedev, V.T.; Torok, G.; Melenevskaya, E.Y.; Vinogradova, L.V.; Ivanova, I.N. Poly(N-vinylcaprolactam)-C₆₀ complexes in aqueous solution, *Fullerenes Nanotubes & Carbon Nanostructures*, **2008**, *16*, 603-609.
- [299]Qiao, X.; Huang, C.; Ying, Y.; Yang, X.; Liu, Y.; Tian, Q. Involvement of reactive oxygen species and calcium in photo-induced membrane damage in HeLa cells by a bis-methanophosphonate fullerene, *J. Photochem. Photobiol. B*, **2010**, *98*, 193-198.
- [300]Palyvoda, K.O.; Grynyuk, I.I.; Prylutsk, S.V.; Samoylenko, A.A.; Drobo, L.B.; Matyshevsk, O.P. Apoptosis photoinduction by C₆₀ fullerene in human leukemic T cells, *Ukr. Biochem. J.*, **2010**, *82*, 121-127..
- [301]Mroz, P.; Xia, Y.; Asanuma, D.; Konopko, A.; Zhiyentayev, T.; Huang, Y.Y.; Sharma, S.K.; Dai, T.; Khan, U.J.; Wharton, T.; Hamblin M.R. Intraperitoneal photodynamic therapy mediated by a fullerene in a mouse model of abdominal dissemination of colon adenocarcinoma, *Nanomed.: Nanomed. Biol. Med.*, **2011**, *7(6)*, 965-974.
- [302]Xiao, L.; Aoshima, H.; Saitoh, Y.; Miwa, N. Fullerene-polyvinylpyrrolidone clathrate localizes in the cytoplasm to prevent ultraviolet-A ray-induced DNA-fragmentation and activation of the transcriptional factor NF-kappaB, *J. Cell. Biochem.*, **2010**, *111*, 955-966.
- [303]Constantin, C.; Neagu, M.; Ion, R.; Gherghiceanu, M.; Stavaru, C. Fullerene-porphyrin nanostructures in photodynamic therapy, *Nanomedicine*, **2010**, *5*, 307-317.
- [304]Sharma, S.K.; Chiang, L.Y.; Hamblin, M.R. Photodenamic therapy with fullerenes *in vivo*: reality or a dream?, *Nanomedicine*, **2011**, *6*, 1813-1825.
- [305]Ni, J.; Q.Y.Wu, Y.G.Li, Z.X.Guo, G.S.Tang, D.Sun, F.Gao, J.M.Cai, Cytotoxic and radiosensitizing effects of nano-C₆₀ on tumor cells *in vitro*, *J. Nanoparticle Res.*, **2008**, *10*, 643-651.
- [306]Andrievsky, G.; Bruskov, V.I.; Tykhomyrov, A.A.; Gudkov, S.V. Peculiarities of the antioxidant and radioprotective effects of hydrated C₆₀ fullerene nanostructures *in vitro* and *in vivo*, *Free Radic. Biol. Med.*, **2009**, *47*, 786-793.
- [307]Huang, S.Q.; Gao, Y.; Li, F.; Cui, B.; Zhao, J.; Dong, F.; Cai, J. Synthesis of fullerene derivative C(60)-Lys and its radio-protection effects in AHH-1 cell, *J. Rad. Res. Rad. Proces.*, **2010**, *#1*, 37-41.
- [308]Theriot, A.C.; Casey, R.C.; Moore, V.C.; Mitchell, L.; Reynolds, J.O.; Burgoyne, M.; Partha, R.; Huff, J.L.; Conyers, J.L.; Jeevarajan, A. Dendro[C₆₀]fullerene DF-1 provides radioprotection to radiosensitive mammalian cells, *Rad. Envir. Biophys.*, **2010**, *49*, 437-445.
- [309]Brown, A.P.; Chung, E.J.; Urick, M.E.; Shield, W.P.; Sowers, A.L.; Thetford, A.; Shankavaram, U.T.; Mitchell, J.V.; Citrin, D.E. Evaluation of the fullerene compound DF-1 as a radiation protector, *Radiat. Oncol.*, **2010**, *5*, 34-43.
- [310]Daroczi, B.; Kari, G.; McAleer, M.F.; Wolf, J.C.; Rodeck, U.; Dicker, A.P. *In vivo* radioprotection by the fullerene nanoparticle DF-1as assessed in a Zebrafish model, *Clin. Cancer Res.*, **2006**, *12*, 7086-7091.
- [311]Foley, S.; Crowley, C.; Smahhi, M.; Bonfils, C.; Erlanger, B.F.; Seta, P.; Larroque, C. Cellular localisation of a water-soluble fullerene derivative, *Biochem. Biophys. Res. Commun.*, **2002**, *294*, 116-119.

- [312]Chirico, F.; Fumelli, C.; Marconi, A.; Tinari, A.; Straface, E.; Malorni, W.; Pellicciari, R.; Pincelli, C. Carboxyfullerenes localize within mitochondria and prevent the UVB-induced intrinsic apoptotic pathway, *Exp. Dermatol.*, **2007**, *16*, 429–436.
- [313]Dobrovolskaia, M.A.; McNeil, S.E. Immunological properties of engineered nanomaterials, *Nat. Nanotechnol.*, **2007**, *2*, 469–478.
- [314]Cai, X.; Hao, J.; Zhang, X.; Yu, B.; Ren, J.; Luo, C.; Li, Q.; Huang, Q.; Shi, X.; Li, W.; Liu, J. The polyhydroxylated fullerene derivative C₆₀(OH)₂₄ protects mice from ionizing-radiation-induced immune and mitochondrial dysfunction, *Toxicol. Appl. Pharmacol.*, **2010**, *243*, 27-34.
- [315]Injac, R.; Perse, M.; Cerne, M.; Potocnik, N.; Radic, N.; Govedarica, B.; Djordjevic, A.; Cerar, A.; Strukelj, B. Protective effects of fullereneol C₆₀(OH)₂₄ against doxorubicin-induced cardiotoxicity and hepatotoxicity in rats with colorectal cancer, *Biomaterials*, **2009**, *30*, 1184–1196.
- [316]Dordević, A.; Bogdanović, G., Fullereneol—a new nanopharmaceutic? *Arch. Oncol.*, **2008**, *16*, 42–45.
- [317]Takada, H.; Kokubo, K.; Matsubayashi, K.; Oshima, T. Antioxidant activity of supramolecular water-soluble fullerenes evaluated by β-carotene bleaching assay, *Biosci. Biotechnol. Biochem.*, **2006**, *70*, 3088-3093.
- [318]Vávrová, J.; Řezáčová, M.; Pejchal, J. Fullerene nanoparticles and their anti-oxidative effects: a comparison to other radioprotective agents, *J. Appl. Biomed.*, **2012**, *10*, 1-8
- [319]Fourches, D.; Pu, D.; Tassa, C.; Weissleder, R.; Shaw, S.Y.; Mumper, R.J.; Tropsha, A. Quantitative nanostructure-activity relationship modeling, *AcsNano*, **2010**, *4*, 5703-5712.
- [320]Kayat, J.; Gajbhiye, V.; Tekade, R.K.; Jain, N.K. Pulmonary toxicity of carbon nanotubes: a systematic report, *Nanomed. Nanotechnol. Biol. Med.*, **2011**, *7*, 40-49.
- [321]Miller, J.; Lam, M.; Lebovitz, R. *Derivatized Fullerenes: A New Class of Therapeutics and Imaging Agents*. Nanotech. L. & Bus. <http://heinonline.org/>, 423, **2007**.
- [322]Kepley, C. Use of fullerenes for the treatment of mast cell and basophil-mediated disease, US Patent 7947262, **2006**.
- [323]Bystrzejewska-Piotrowska, G.; Golimowski, J.; Urban, P.L. Nanoparticles: Their potential toxicity, waste and environmental management, *Waste Management*, **2009**, *29*, 2587-2595.
- [324]Mraz, S.J. Nanowaste: the next big threat? *Machine Design*, **2005**, *77*, 46–53.
- [325]Lewinski, N.; Colvin, V.; Drezek, R. Cytotoxicity of nanoparticles, *Small*, **2008**, *4*, 26–49.

ABBREVIATIONS

AA – ascorbic acid
 AD – Alzheimer's disease
 ALM – liposomal malonyl fullerene-C₇₀
 AMPA – 2-amino-3(3-hydroxy-5-methylisoxazol-4-yl)methyl isopropionic acid
 AMPK – AMP-activated protein kinases
 AP1 – activator protein 1 – transcription factor (leucine zipper)
 APTT – activated partial thromboplastin time
 BBB – blood brain barrier
 BMPF – bis-methanophosphonate-C₆₀
 BSA – bovine serum albumin
 CAs – carbonic anhydrases
 CAT – catalase
 C3 – tris-malonyl-C₆₀
 CD – cyclodextrin
 C₆₀-DF – C₆₀-dendrofullerene with 18 carboxyl groups
 CINC – cytokine-induced neutrophil chemoattractant
 CMECs- brain microvessel endothelial cells
 Cyt C – cytochrome C
 Curdlan – linear 1,3β-glucan
 DFT – density functional theory
 DMF – dose modifying factor
 DOX – doxorubicin
 ERK – signal regulated kinases
 ER – endoplasmatic reticulum
 FMD – C₆₀-methionine
 GSH/GSSG – reduced/oxidized glutathione
 GluR – glutamate receptors
 Gpx – glutathione peroxidase
 GST – glutathione-S-transferase
 Gsr – glutathione reductase
 HAS – human serum albumin
 HH-F – C₆₀(OH)₃₂₋₃₄ 7H₂O
 HIF – hypoxia-inducible factor responsible for the reaction to the lack of oxygen
 Hsp70 – molecular shaperon
 ID4 – inhibitor of differentiation 4
 IL12; p70; IFN-γ; IL1β; IL2; IL6 – cytokins
 JNK – c-Jun amino terminal kinases
 KA – kainate

LH-F – C₆₀(OH)₆₋₁₂
MES – solvent exchange method
MMP-1; MMP-3; MMP-13 – matrix metalloproteinases
MDA – malonic dialdehyde
MPTP – 1-methyl-4-phenyl-1,2,3,6-tetrahydropyridine
MAPK – mitogen activated protein kinases
MHC – major histocompatibility complex
NADH/NAD⁺ – reduced/oxidized nicotinamide adenine
NAC – N-acetyl-L-cysteine
NMDA – N-methyl-D-aspartate
NF-κB – nuclear factor kappa light-chain-enhancer of activated B cells (transcription factor)
NOS – NO-synthases
PAHs – polycyclic aromatic hydrocarbons
PARP – poly(ADP-ribose)polymerase
PCR – polymerase chain reaction
PD – Parkinson's disease
PDT – photodynamic therapy
PEG – polyethylene glycol
PEGAM 1 (CD31) – platelet/endothelial cell adhesion molecule 1
PMC16 – Buckminster fullerene(C₆₀)-2-(butadiene-1-yl) tetra-(*o*-γ-amino-butyryl-*o*-phtalyl) ferroporphyrin
POX – poly(2-alkyl-2-oxazoline)
PTX – pentoxifylline
PT – protrombin time
PVP – polyvinylpyrrolidone
RANK – receptor activator NF-κB
ROS – reactive oxidative species
SNP – sodium nitroferricyanide
SHH-F – C₆₀(OH)₄₄ 8H₂O
SOD – superoxide dismutase
STZ – streptozotocin
TA – peptide Thr-Lys-Pro-Arg
TGF-β – transforming growth factor β
TNF-α – tumor necrosis factor α