

Initial Symptoms of Morbus Behcet

James Patrick*

Department of Obstetrics and Gynecology, Harvard University, Cambridge, MA

*Corresponding Author: James Patrick, Department of Obstetrics and Gynecology, Harvard University, Cambridge, MA, E-mail: jpatrick@harvard.edu

CLINICAL IMAGE

In the year 2013, one 23 years old female presented with severe spontaneous pain in vagina and lesions on vulva. Macroscopic and under colposcopy I found white, solid, hard, round, painful and easy bleeding lesions on board with healthy tissue. Biopsy result was Mucosis manifestation of antibody reaction full of leukocyte, Papanikolau test then was III.

For 3-4 days pain and others symptoms disappeared without any treatment. I thought that it could be Linear IgA disease. After few months I took again Papanikolau test and was II.

3 years after this lesions in genital region, start to show up new lesions in mouth and esophagus with oral disorders. Lesions looks like aphta. They are often, painful, rounds. After exams, rheumatologist gave her another drug "Morbus Behcet". Her gynecological status now is in order Figure 1,2,3,4,5.



Figure 1 Macroscopic and under colposcopy discovered white, strong, hard, round, difficult and simple draining injuries ready for solid tissue.



Figure 2 Biopsy result was Mucosis appearance of counter acting agent response brimming with leukocyte.



Figure 3 After Papanikolaou test.



Figure 4 Before infusing dg "Morbus Behcet".



Figure 5 Status was Normal After infusing dg "Morbus Behcet".