



Scholars Research Library
Journal of Computational Methods in Molecular Design, 2021, 11 (6): 1-2
(<http://scholarsresearchlibrary.com>)



Modification of nutritional properties of microalgae for Artemia breeding

Zeynep E Cakmak*

Karamanoglu Mehmetbey Üniversitesi, Department of Bioengineering Bioengineering

*Corresponding Author: Numan Eczacioglu, Karamanoglu Mehmetbey Üniversitesi, Department of Bioengineering Bioengineering; Email: numaneczacioglu@gmail.com

ABSTRACT

Artemia (brine shrimp) is used as a live-feed stuff for seed fish in fish hatcheries and aquarium fisheries. Nutritional properties of *Artemia* are in close relationship with the nutritional facts of the microalgae it is fed by. In this study, 20 different microalgae and cyanobacteria (indigenous strains from *Dunaliella*, *Isochrysis*, *Phaeodactylum*, *Tetraselmis*, *Nannochloropsis*, *Spirulina*, *Synechocystis*, *Synechococcus*, *Chlamydomonas*, *Chlorella*, and *Scenedesmus* genus) were supplied to *A. franciscana* as food source and growth characteristics of *A. franciscana* and were followed during 10 days of growth. Seven microalgae strains were selected for *Artemia* breeding and dry weight, total protein, starch and lipid contents of microalgae and *A. franciscana* were recorded. Then, microalgae were exposed to N-, S-, P-deprivation and high salt stress for 5 days of incubation.

Key words: *Artemia*, cyanobacteria, *Nannochloropsis*, *franciscana*

INTRODUCTION

Artemia (brine shrimp) is used as a live-feed stuff for seed fish in fish hatcheries and aquarium fisheries. Nutritional properties of *Artemia* are in close relationship with the nutritional facts of the microalgae it is fed by. In this study, 20 different microalgae and cyanobacteria (indigenous strains from *Dunaliella*, *Isochrysis*, *Phaeodactylum*, *Tetraselmis*, *Nannochloropsis*, *Spirulina*, *Synechocystis*, *Synechococcus*, *Chlamydomonas*, *Chlorella*, and *Scenedesmus* genus) were supplied to *A. franciscana* as food source and growth characteristics of *A. franciscana* and were followed during 10 days of growth. Seven microalgae strains were selected for *Artemia* breeding and dry weight, total protein, starch and lipid contents of microalgae and *A. franciscana* were recorded. Then, microalgae were exposed to N-, S-, P-deprivation and high salt stress for 5 days of incubation. Total lipid, protein and carbohydrate contents of those strains were recorded and 5-days stress exposed microalgae were supplied to *A. franciscana* as only food source.

Lastly total lipid, protein and carbohydrate content of *A. franciscana* was followed during 10 days of growth. In most cases, feeding *A. franciscana* with *D. tertiolecta* was superior to other strains studied. Portunid crabs stand out as highly valued resources for fisheries and aquaculture because of their export potential and high nutritional value. Due to their size, meat content and unique flavor, their products are highly priced in domestic and international markets. Since global portunid product demands exceed expectations each year, world fisheries captures have grown steadily, surpassing 1 million tons by 2016, leading to the local overexploitation of some species. At the same time, unsatisfied market demands have been increasingly sustained by restocking and aquaculture production in excess of 0.38 million tons by 2016, 96% of which were produced in East Asia.

At present, portunid aquaculture is restricted to meat production of *Scylla serrata*, *Portunus pelagicus*, *Portunus trituberculatus*, *Portunus sanguinolentus* and *Charybdis feriata* and to restocking of *Callinectes sapidus*. Still, the culture potential for many other large-sized species from the taxon is virtually unexplored. Indeed, reported production out of the west coast of Asia is still negligible, representing both a challenge and an opportunity for the industry sector elsewhere. The southern surf crab *Ovalipes trimaculatus* (de Haan, 1833), one of the species with high potential for aquaculture, is widely distributed in coastal areas of the South Atlantic, Indian and Pacific Oceans, being present along the mid-latitude (25°–45° S) Argentinean coast, where populations have been targeted by artisanal fisheries over the last 10 years providing products with good acceptance in the local shellfish markets.

Although several studies have been conducted on the structure of its populations, reproduction, growth and some behavioral and anatomical aspects information available on the biology of its early life stages is still scarce and insufficient to allow encouraging their breeding in aquaculture facilities. Larval stages of decapod crustaceans may be lecithotrophic or planktotrophic depending on the reproductive strategy of each species. While the former cover their food requirements by consuming abundant yolk reserves stored in the oocytes, the later start feeding on different plankton components soon after hatching. Thus, breeding planktotrophic decapod larvae requires assessing the quality and frequency of food consumption to optimize survival, growth and physiological condition. However, this involves the simultaneous maintenance of larvae and auxiliary food cultures, representing the bottleneck for the aquaculture of many decapods, including portunid crabs.

Research on dietary quantity and quality requirements has contributed to minimize mortality and to enhance growth and fitness of zoeae from several portunids including *Scylla serrata*, *Portunus sanguinolentus*, *P. pelagicus* and *Callinectes sapidus*. Among other live feeds, brine shrimps (*Artemia* spp.) are widely used due to their food carrying capacity and good acceptance. At low temperatures (i.e., 12°), like those experienced by *O. trimaculatus* during the reproductive season, *Artemia persimilis*, a brine shrimp species native from Argentina and Chile, shows higher survival rates compared to its native congener *A. franciscana*, probably resulting from its adaptation to Patagonian climate conditions. Its cysts have nutritional properties comparable to those of other commercial species traded in international markets, displaying high hatching efficiencies and producing small-sized nauplii with Extended Abstract International Conference on Biotechnology, Biomarkers & Systems Biology March 04-05, 2019 | Amsterdam, Netherlands Volume 4 Issue 5 elevated fatty acid unsaturation, highly desirable for use as live food in aquaculture. Still, up to date the species has been rarely used to feed larval stages of fishes or marine invertebrates

REFERENCES

- [1] Sreelakshmi, K. R., Manjusha, L., Vartak, V. R. et al. Variation in proximate composition and fatty acid profiles of mud crab meat with regard to sex and body parts. *Indian J. Fish* **2016**; 63(2), p. 147–150
- [2] Aguilar, R., Johnson, E. G., Hines, A. et al. Importance of blue crab life history for stock enhancement and spatial management of the fishery in Chesapeake Bay. *Rev. Fish. Sci.* **2008**; 16(1–3), p. 117–124
- [3] Jenkins, G. I., Michael, R., Tweedley, J. R. Identifying future restocking options for Blue Swimmer Crabs (*Portunus armatus*). In Recreational Fishing Initiatives Fund: Project **2015/09**, p. 30.
- [4] Feng, X., Johnson, E. G., Williams, E. P. & Place, A. R. Successful identification and discrimination of hatchery-reared blue crabs (*Callinectes sapidus*) released into the Chesapeake Bay using a genetic tag. *J. Shellfish Res.* **2017** 36(1), p. 277–282
- [5] Spivak, E. D. Los crustáceos decápodos del Atlántico sudoccidental (25°–55°S): distribución y ciclos de vida. *Investig. Mar.* **1997**; 25, 69–91
- [6] Dima, J. B., Barón, P. J. & Zaritzky, N. E. Mathematical modelling of the heat transfer process and protein denaturation during the thermal treatment of Patagonian marine crabs. *J. Food Eng.* **2012**; 113(4), p. 623–634