



Scholars Research Library

Archives of Applied Science Research, 2012, 4 (3):1411-1417
(<http://scholarsresearchlibrary.com/archive.html>)



Synthesis, Characterization and Antimicrobial Studies of (N¹E, N⁶E)-N¹, N⁶- bis ((6-bromo-2-hydroxyquinolin-3-yl) methylene) adipohydrazide

K. Siddappa* and Mallikarjun Kote

Department of Post Graduate Studies and Research in Chemistry, Gulbarga University,
Gulbarga-585106, Karnataka, India

ABSTRACT

The Cu(II), Co(II), Ni(II), Zn(II), Cd(II), Hg(II), and Mn(II) complexes of Schiff base (N¹E, N⁶E)-N¹, N⁶-bis((6-bromo-2-hydroxyquinolin-3-yl)methylene)adipohydrazide has been synthesized and characterized by various spectral techniques. The complexes were prepared by reacting the ligand and metal chloride of Cu(II), Co(II), Ni(II), Zn(II), Cd(II), Hg(II), and Mn(II) in ethanol to get a series of mononuclear complexes. The complexes were characterized by CHN analysis, conductivity measurements, magnetic susceptibility, IR, NMR, ESR, UV-Vis and X-ray powder diffraction studies. By these spectral studies it is found that Cu(II), Co(II), Ni(II), Zn(II), Cd(II) Hg(II) and Mn(II) complexes have exhibited octahedral geometry. The ligand and its metal complexes have been screened for their antimicrobial activities. The prepared ligand shows low activity and its metal complexes shows moderate to good activity.

Key words: Quinoline, Schiff base, spectral studies, metal complexes and antimicrobial activity

INTRODUCTION

The synthesis of quinoline and its derivatives have attracted considerable attention of organic and medicinal chemists for many years.^[1-4] The structural core of quinoline is frequently associated with medicinal applications, such as anti-cancer,^[5] antimicrobial,^[6] integrase inhibitors,^[7] HIV protease inhibitors,^[8] antileishmanial activity,^[9] quinoline derivatives as a source of valuable drug candidates and useful intermediates in organic chemistry^[10] thus, the synthesis of this heterocyclic nucleus is of much used in pharmacopeia in treatment of malaria^[11] and more recently tumors^[12]. Our work is concerned with complexation in aqueous- organic solvents attract study researchers participation of the organic component. As a continuation of studies concerning the effect of acid hydrazides^[13-15], we examined in this study the complexing properties of adipic acid hydrazide ion aqueous-ethanol solutions. The main goals of this study were describing the complexation of Cu(II), Co(II), Ni(II), Zn(II), Cd(II), Hg(II) and Mn(II) with ligand and revealing the effects of the central metal-ion.

MATERIALS AND METHODS


Characterization Techniques

Infrared spectra of the ligand and its metal complexes in KBr pellets were recorded in the spectral range 3800 – 350 cm^{-1} with Perkin Elmer Spectrum One FT-IR Spectrometer. $^1\text{H-NMR}$ Spectra were recorded on AMX-4000 NMR Spectrometer, using TMS as internal standard and DMSO as a solvent. The Electronic Spectra were recorded on an Elico-SL-164 Double BEAM UV-Visible Spectrophotometer in the range of 200–1200nm in DMF. X-ray powder diffraction was recorded at department of physics Gulbarga university, Gulbarga. Magnetic susceptibility measurements were recorded on Guoy balance method. The Conductance measurements were made on an ELICO-CM-82-conductivity bridge.

Preparation of Ligand

(N^1E, N^6E)- N^1, N^6 -bis ((6-bromo-2-hydroxyquinolin-3-yl) methylene) adipohydrazide:

The Schiff base ligand were prepared by condensation of adipic dihydrazide (1 mole, 1.74g) and 6-bromo-2-hydroxyquinoline-3-carbaldehyde (2mole, 2.52g) in ethanol (30ml) was refluxed on water bath for 5-6 hours in presence of few drops of acetic acid. The reaction mixture was cooled to room temperature, the separated Schiff base were filtered and washed with hot alcohol and recrystallized from alcohol to get a pure sample. The purity of Schiff base was checked by TLC. Yield: 70%, MP; 275°C, Mol. Wt = 642.30.



SCHEME

Preparation of Metal complexes

A solution of 0.01mole of metal chloride in ethanol was mixed with the ethanolic solution of 0.01 mole of ligand and refluxed for 3-4 hours on water both to get clear solution. 0.5gm of excess sodium acetate was added to the reaction mixture to adjust the pH 7-8 of the solution. The reaction mixture was further, refluxed for 2 hours more.

The resulting mixture was decomposed by pouring into a 100ml of distilled water with constant stirring. The suspended solid complex was allowed to settle and collected by filtration, washed with sufficient quantity of distilled water and then with little hot ethanol to apparent dryness and dried in a vacuum over anhydrous calcium chloride in a desiccators. (Yield, 55-70%)

RESULTS AND DISCUSSION

Nature and Stoichiometry

All the complexes are colored in nature and were sparingly soluble in common organic solvents but soluble in DMF, DMSO and acetonitrile. The analytical data indicates that the complexes are agree well with 1:1 metal to ligand stiochiometry for Cu(II), Co(II), Ni(II) Zn(II), Cd(II), Hg(II) and Mn(II). The stoichiometry of all the complexes was confirmed by spectrophotometric method. The observed molar conductance values measures in DMF solution fall in the range 12–20 $\text{Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$ (Table 1).These observed values of the molar conductance are well within the expected range for non-electrolytic nature ^[16]. The physical characterization data of all the compounds has been summarized in Table 1.

Table 1. Analytical, magnetic susceptibility, molar conductance of the ligand and its metal complexes

Ligand / Complex	Mol. Wt.	M.P (°C)	Yield (%)	Found /Calculated (%)				μ_{eff} B.M.	Molar conductance $\text{Ohm}^{-1}\text{cm}^2\text{mol}^{-1}$
				M	C	H	N		
$\text{C}_{26}\text{H}_{22}\text{Br}_2\text{N}_6\text{O}_4$	642.30	294	70	-	48.60 (48.62)	3.40 (3.45)	13.10 (13.08)	-	-
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{CuN}_6\text{O}_4$	703.83	320	65	9.05 (9.03)	44.35 (44.37)	2.80 (2.86)	11.90 (11.94)	1.92	18
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{CoN}_6\text{O}_4$	699.22	316	62	8.48 (8.43)	44.60 (44.66)	2.82 (2.88)	12.05 (12.02)	4.74	17
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{NiN}_6\text{O}_4$	698.98	325	62	8.42 (8.40)	44.66 (44.68)	2.80 (2.88)	12.07 (12.02)	2.92	18
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{MnN}_6\text{O}_4$	705.69	332	60	9.34 (9.27)	44.30 (44.25)	2.85 (2.86)	12.00 (11.91)	5.75	17
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{CdN}_6\text{O}_4$	752.69	344	65	14.92 (14.93)	41.55 (41.49)	2.62 (2.68)	11.20 (11.17)	Diamagnetic	18
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{HgN}_6\text{O}_4$	840.87	358	68	23.92 (23.85)	37.20 (37.14)	2.49 (2.40)	9.91 (9.99)	Diamagnetic	12
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{ZnN}_6\text{O}_4$	695.22	367	70	7.85 (7.90)	44.85 (44.92)	2.95 (2.90)	12.12 (12.09)	Diamagnetic	16

Table 2. IR Spectral data of the ligand and its metal complexes (cm^{-1})

Ligand / complex	ν_{OH}	ν_{NH}	$\nu_{\text{C=O}}$	$\nu_{\text{C=N}}$	$\nu_{\text{C-O}}$	$\nu_{\text{M-O}}$	$\nu_{\text{M-N}}$
$\text{C}_{26}\text{H}_{22}\text{Br}_2\text{N}_6\text{O}_4$	3430	3172	1660	1596	1201	-	-
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{CuN}_6\text{O}_4$	-	3184	1615	1562	1236	550	472
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{CoN}_6\text{O}_4$	-	3198	1626	1565	1262	562	492
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{NiN}_6\text{O}_4$	-	3186	1648	1572	1265	564	480
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{N}_6\text{O}_4\text{Zn}$	-	3196	1642	1578	1264	558	462
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{CdN}_6\text{O}_4$	-	3194	1642	1562	1230	554	464
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{HgN}_6\text{O}_4$	-	3195	1645	1558	1228	556	470
$\text{C}_{26}\text{H}_{20}\text{Br}_2\text{MnN}_6\text{O}_4$	-	3194	1652	1560	1262	554	478

Infrared Spectra

The significant IR bands for the ligand ($\text{N}^1\text{E}, \text{N}^6\text{E}$)- N^1, N^6 -bis((6-bromo-2-hydroxyquinolin-3-yl)methylene)adipohydrazide as well as for its metal complexes and their tentative assignments are compiled and represented Table 2. The broad band observed at 3430cm^{-1} in the IR spectra of the ligand assigned to ν_{OH} , which were found to have disappeared in all their respective complexes, there by indicating the involvement of phenolic oxygen is bonding with metal ions through deprotonation. The band ν_{NH} observed at 3172cm^{-1} in ligand and complexes ν_{NH} observed at $3172\text{-}3198\text{cm}^{-1}$ respectively. A strong sharp band observed at 1660cm^{-1} is assigned to adipic hydrazide ring $\nu_{\text{C=O}}$, which was shifted to $5\text{-}48\text{cm}^{-1}$ in all complexes, indicates the involvement of adipic hydrazide ring carboxyl in complexation with metal ion, the band at 1596cm^{-1} is assigned to the azomethine $\nu_{\text{C=N}}$ group, lowering of $\nu_{\text{C=N}}$ $1\text{-}38\text{cm}^{-1}$ in the complexes as compared to its ligand, is due to reduction

of double bond character carbon-nitrogen bond of the azomethine group^[17]. The band observed at 1201 cm⁻¹, of the ligand is attributed to phenolic $\nu_{(C-O)}$ in view of previous observations^[18]. This band is shifted to higher frequency and is found in the region 1201-1265 cm⁻¹ (1-64 cm⁻¹) for the complexes. Thus, the low frequency skeletal vibrations due to $\nu_{(M-O)}$ and $\nu_{(M-N)}$ stretching provide direct evidence for complexation. In the present investigation the bands in the 550-564 cm⁻¹ region for $\nu_{(M-O)}$ and 470 -492 cm⁻¹ region for $\nu_{(M-N)}$ vibration respectively.

Magnetic movement and electronic spectra of the complexes

Copper(II) complex

The complexes having temperature independent magnetic moments in the range of 1.74-2.2 B.M. agreeable to the spin only value. These complexes are mononuclear having no major spin-spin interactions; however no compromise can be set between the magnitude of the distortion and other variable factors. The Cu(II) complexes with this behavior indicate marked spin-exchange and strong coupling interaction between Cu(II) atoms. As the spin only value for a single unpaired electron is 1.72 B.M. The slight excess over this value for the Cu(II) complexes can be attributed to spin orbital coupling. The observed magnetic moment values for the present Cu(II) complexes is 1.92 B.M.^[19]. This value is expected for one electron system and is in the normal range of Cu(II) complex system. As the spin only values for a single unpaired electron is 1.72 B.M. The slight excess over this value found for the Cu(II) complex can be attributed to spin orbital coupling. In view of the above observations the magnetic moment values for the present Cu(II) complexes, suggest the distorted octahedral geometry around Cu(II) ions.

Electronic spectra of Cu(II) complexes exhibit a broad, low intensity shoulder band. The 2E_g and ${}^2T_{2g}$ states of octahedral Cu(II) ion (d^9) split under the influence of the tetragonal distortion and the distortion can be such as to cause the three transition viz., ${}^2B_{1g} \rightarrow {}^2B_{2g}$, ${}^2B_{1g} \rightarrow {}^2E_g$ and ${}^2B_{1g} \rightarrow {}^2A_{1g}$, to remain unresolved in the spectra. It is concluded that all three transition lie within the single broad band in the region 25400 – 31000 cm⁻¹, have been assigned to charge transfer bands from ligand to metal. The observation favor distorted octahedral geometry around Cu(II) complexes. A broad asymmetric band in the region 13495-16605 cm⁻¹, the symmetry being on the lower energy side^[20]. The broadness of the band may be due to dynamic John-Teller distortion. The band maxima observed for the present Cu(II) complexes is at 12099-16700 cm⁻¹, 12305-16435 cm⁻¹ and 12400 – 16360 cm⁻¹ suggest that the complex displays coordination number six. In addition, we have observed a high intensity band around 15165cm⁻¹ that can be attributed to the ligand-metal charge transfer transition.

Cobalt(II) complex:

In octahedral Co(II) complexes the ground state is ${}^4T_{1g}$ and a large orbital contribution to the magnetic moment values for various Co(II) complexes are in the range 4.11-4.71 B.M. and 4.6-5.18 B.M. for tetrahedral and octahedral complexes respectively. In present investigation the observed magnetic moment values is in the range 4.74 B.M.^[21], which indicates octahedral geometry for these Co(II) complexes. This is due to partial quenching of orbital contribution to the magnetic moment. Electronic spectra of Co(II) complex exhibits three bands in the region 9624-10350cm⁻¹, 16504-18385⁻¹ and 24844-26954 cm⁻¹ due to the transition. ${}^4T_{1g}(F) \rightarrow {}^4T_{2g}(F)$ $\nu_1 = 9624-10350\text{cm}^{-1}$, ${}^4T_{1g}(F) \rightarrow {}^4A_{2g}(F)$ $\nu_2 = 16504-18385\text{cm}^{-1}$ and ${}^4T_{1g}(F) \rightarrow {}^4T_{1g}(P)$ $\nu_3 = 24844-26954\text{cm}^{-1}$. These transitions suggest octahedral geometry. The region at 25650-26870 cm⁻¹ refers to the charge transfer band. These transitions suggest octahedral geometry for the Co(II) complexes. These assignments are in good agreement with the reported values^[22].

Nickel(II) complex:

Ni(II) complexes exhibits the magnetic moment value of 2.92 which are well within the range of expected value for octahedral geometry around the central metal ion^[22]. Electronic spectra of Ni(II) complexes in octahedral coordination is ${}^3A_{2g}$, the Ni(II) complexes show three transition in an octahedral field, viz., ${}^3A_{2g}(F) \rightarrow {}^3T_{2g}(F)$; $\nu_1 = 7475- 8200\text{cm}^{-1}$, ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(F)$; $\nu_2 = 16504-17100\text{cm}^{-1}$ ${}^3A_{2g}(F) \rightarrow {}^3T_{1g}(P)$; $\nu_3 = 24844 -26000\text{cm}^{-1}$. The observed transition bands lie well within in the range of reported values. These values indicate considerable covalent character and support the octahedral geometry for the Ni(II) complex^[23].

Manganese(II) complex

The observed magnetic moment values for Mn(II) complex of the ligand is in the range 5.75 B.M. The electronic spectra of Mn(II) complex shows three transition, which corresponds to the following transitions, ${}^6A_{1g} \rightarrow {}^4T_{1g}(4G)$; $\nu_1 = 15637-16500\text{cm}^{-1}$, ${}^6A_{1g} \rightarrow {}^4E_g(4D)$; $\nu_2 = 17525-18600\text{cm}^{-1}$, ${}^6A_{1g} \rightarrow {}^4T_{1g}(4D)$; $\nu_3 = 21743-22700\text{cm}^{-1}$ These values

indicate considerable covalent character and support the octahedral geometry^[24]. Hence under the present study octahedral geometry is suggested for Mn(II) complex.

Table 3. Electronic spectral data of ligand field parameters of Cu(II), Co(II), Ni(II) and Mn(II) metal complexes

Complexes	$\nu_1(\text{cm}^{-1})$	$\nu_2(\text{cm}^{-1})$	$\nu_3(\text{cm}^{-1})$	$Dq(\text{cm}^{-1})$	B^1	β	$\beta\%$	ν_2/ν_1	ν_3/ν_2	LFSE k cal mol ⁻¹
C ₂₆ H ₂₀ Br ₂ CuN ₆ O ₄	12099 – 167 00			1428	-	-	-	-	-	24.56
C ₂₆ H ₂₀ Br ₂ CoN ₆ O ₄	9624	14618	20743	931	824	0.94	15.06	1.54	1.47	14.34
C ₂₆ H ₂₀ Br ₂ NiN ₆ O ₄	7475	16504	24844	911	836	0.85	19.84	1.23	1.54	32.76
C ₂₆ H ₂₀ Br ₂ MnN ₆ O ₄	15637	17525	21743	934	868	0.77	22.36	1.15	1.26	14.48

¹H NMR Spectra:

Spectrum of ¹H NMR is DMSO-d₆ solvent used. In ligand (N¹E, N⁶E)-N¹, N⁶-bis((6-bromo-2-hydroxyquinolin-3-yl)methylene)adipohydrazide showed the sharp peak at δ 9.6 (S, 2H) due to OH at 2-position of phenyl ring of 6-bromo-2-hydroxy quinoline moiety has resonated, but in the case of Zn(II) complex which has been disappeared indicating the involvement of phenolic oxygen in the coordination *via*, deprotonation^[25,26]. A single large peak showed at δ 3.3 (S, 3H, CH₃) due to protons of the azomethine group in ligand but in case of Zn(II) complex the peak observed at δ 2.5 (S, 3H, CH₃)^[27]. The Twelve aromatic protons due to quinoline and phenyl rings have resonated in region δ 7.0 - 7.9 (m, 12H, Ar-H) as a multiplet, in Zn (II) complex the twelve aromatic protons have been observed in the region δ 7.0-8.6 (m, 12H, Ar-H) as a multiplet. The Zn(II) complex suggests coordination of the phenolic OH with metal ion.

ESR spectrum:

The ESR spectrum of the powder Cu(II) complex was recorded at room temperature using DPPH as a standard showed a broadened feature without hyperfine splitting due to the dipolar interaction from the ESR spectrum of a set of magnetic parameter $g_{\parallel} = 2.0533$, $g_{\perp} = 2.0264$. The observed ESR spectrum is characteristic of distorted octahedral geometry g value averaged to over all directions and G which is measure of extent of exchange interaction between metal ion have been calculated. In present case the value of G was found to be 4.0266. according to Hathway, If G value if grater then 4, the spin exchange interaction is negligible where as G value is less than 4 indicate considerable interaction between metal ions in solid complex clearly indicate that Cu(II) ion in the complex is mononuclear nature of the complex.

Table 4: ESR data of the Cu(II) complex

Complex	g_{\parallel}	g_{\perp}	g_{av}	g_{iso}	G
C ₂₆ H ₂₀ Br ₂ CuN ₆ O ₄	2.0533	2.0264	2.03976	2.2984	4.0266

Table 4. X- ray powder diffraction data of Cu(II) complex

2θ	θ	$\sin\theta$	$\text{Sin}^2\theta$	$h^2+k^2+l^2$ (a)	$h^2+k^2+l^2$ (b)	h k l	d-spacing		Relative intensity (%)	a (Å°)
							Cal.	Abs		
10.78	5.39	0.0921	0.0083	1	1	1 0 0	8.356	8.342	95.84	8.3556
15.65	7.82	0.1363	0.0186	2.174	2	1 1 0	5.651	5.665	44.61	8.0031
15.97	7.98	0.1391	0.0191	2.271	2	1 1 0	5.542	5.536	54.63	8.8414
16.16	8.08	0.1407	0.0193	2.313	2	1 1 0	5.494	5.483	57.25	8.7615
16.81	8.40	0.1462	0.0219	2.509	2	1 1 0	5.275	5.277	47.55	8.4632
20.78	10.39	0.1891	0.0356	4.201	4	2 0 0	4.073	4.392	51.84	8.1538
20.33	10.16	0.1769	0.0317	3.645	4	2 0 0	4.374	4.368	52.32	8.7499
21.60	10.30	0.1877	0.0355	4.118	4	2 0 0	4.117	4.114	55.61	8.2347
21.77	10.88	0.1884	0.0354	4.174	4	2 0 0	4.088	4.085	56.76	8.1751
22.25	11.12	0.1925	0.0376	4.369	4	2 0 0	3.992	3.994	61.73	8.0009
22.38	11.14	0.1943	0.0375	4.421	4	2 0 0	3.967	3.966	59.74	8.0195

X-ray powder diffraction studies:

X-ray powder diffraction pattern for Cu(II) complex has characterized with a view to find the type of crystal system the XRD data given the table the diffractogram of Cu(II) complex consists of eleven reflections in the range of 10-30 (2θ value) with maxima at $2\theta = 10.78^\circ$ The interplanar spacing(d) has been calculated from the position of intense peaks using Bragg's equation $n\lambda = 2d\sin\theta$, $\lambda = 1.5406 \text{ \AA}$ The observed and calculated values of d are quite consistent (Table 4). The unit cell calculations have been carried out for the cubic system, the set of $h^2 + k^2 + l^2$

values of the complex were found to be 1, 2 and 4 which corresponds to the planes and absence of forbidden number its confirms the cubic symmetry from the above results the unit cell constants for cubic system were found to be $a=b=c = 8.3556 \text{ \AA}$ for the Cu(II) complex of the ligand the complex showed broad peak indicates amorphous nature^[28-29].

Antibacterial activity:

The antibacterial activity results revealed that the ligand and its complexes shown weak to good activity (Table 5). The ligand and its Cu(II), Zn(II), Ni(II) complexes shows weakly active with the zone of inhibition 10-14 mm against the both organisms when compared to the standard drug streptomycin. The Mn(II), Co(II), Cd(II) shows moderate activity as compared to its ligand with zone of inhibition 15-17. The Hg(II) is a good activity with zone of inhibition 18-20 when compared to the standard drug streptomycin^[30-31].

Antifungal Activity:


The antifungal activity, results revealed that the ligand and its Cu(II), Co(II), Ni(II), Zn(II), Cd(II), Hg(II), and Mn(II) complexes have exhibited weak to good activity (Table 5). The ligand and its Co(II) complex shows weak activity with zone of inhibition, the Cu(II), Ni(II), Mn(II), Zn(II), Cd(II), Hg(II) shows moderate activity as compared to its ligand with the zone of inhibition 15-17mm, to good activity with the zone of inhibition of 18-20mm when compared to the standard drug chlotrimazole^[32-34].

Table 5. Antimicrobial activity of the ligand and its metal complexes

Sl. No.	Compound	Antibacterial Activity Zone of inhibition (in mm)		Antifungal Activity Zone of inhibition (in mm)	
		<i>E.Coli</i>	<i>S.aureus</i>	<i>A.niger</i>	<i>A.flavus</i>
1.	C ₂₆ H ₂₂ Br ₂ N ₆ O ₄	08	09	09	10
2.	C ₂₆ H ₂₀ Br ₂ CuN ₆ O ₄	10	12	15	13
3.	C ₂₆ H ₂₀ Br ₂ CoN ₆ O ₄	15	114	15	14
4.	C ₂₆ H ₂₀ Br ₂ NiN ₆ O ₄	13	14	17	16
5.	C ₂₆ H ₂₀ Br ₂ MnN ₆ O ₄	16	13	19	20
6.	C ₂₆ H ₂₀ Br ₂ N ₆ O ₄ Zn	12	14	17	15
7.	C ₂₆ H ₂₀ Br ₂ CdN ₆ O ₄	16	14	18	16
8.	C ₂₆ H ₂₀ Br ₂ HgN ₆ O ₄	16	18	18	19
9.	Streptomycin	24	23	--	--
10.	Chlotrimazole	--	--	25	26
11.	DMF (Control)	0	0	0	0
12.	Bore size	08	08	08	08

CONCLUSION

The elemental analysis, magnetic susceptibility, electronic spectra, IR, ¹H NMR, ESR spectra and X-ray data observations projects the following structures for these complexes where in Cu(II), Co(II), Ni(II), Zn(II), Cd(II) Hg(II) and Mn(II) are exhibit six coordinated octahedral geometry.



Where M = Cu(II), Co(II), Ni(II), Zn(II), Cd(II), Hg(II) and Mn(II),

Acknowledgement

The authors are thankful to the Chairman, Department of Chemistry, Gulbarga University, Gulbarga for providing laboratory facilities. IISc Bangalore for providing ¹HNMR, ESR and XRD are providing for Department of Physics Gulbarga University, Gulbarga.

REFERENCES

- [1] Balasubramanian, M., Keay, A.R., Katritzky, Rees, C.W., Scriven, E.F.V., Eds., **1996**, *Comprehensive Heterocyclic Chemistry II*, Vol. 5, Pergamon, Oxford.
- [2] De, D., Byers, L.D., Krogstad, D.J., **1997**, *J. Heterocycl. Chem.* 34, 315.
- [3] Gilchrist, T.; *J. Chem. Soc., Perkin Trans.* **1**, 2001, 2491.
- [4] Kouznetsov, V., Mendez, L., Gomes, C., **2005**, *Curr. Org. Chem.* 9, 141.
- [5] Elderfield, R.C., Le Von, E.F., **1960**, *J. Org. Chem.* 25, 1576.
- [6] Musiol, R., Jampilek, J., Buchta, V., Silva, L., Niedbala, H., Podeszwa, B., Palka, A., Majerz-Maniecka, K., Oleksyn, B., Polanski., **2006**, *J. Bioorg. Med. Chem.* 14, 3592.
- [7] Bénard, C., Zouhiri, F., Normand-Bayle, M., Danet, M., Desmaële, D., Leh, H., Mouscadet, J.F., Mbemba, G., Thomas, C.M., Bonnenfant, S., Le Bret, M., Angelo., **2004**, *J. Bioorg. Med. Chem. Lett.*, 14, 2473.
- [8] Garrouste, P., Pawlowski, M., Tonnaire, T., Sicsic, S., Dumy, P., De Rosny, E., Reboud-Ravaux, M., Fulcrand, P., Martinez J. **1998**, *Eur. J. Med. Chem.* 33, 423.
- [9] Desrivot, J., Herrenknecht, C., Ponchel, G., Garbi, N., Prina, E., Fournet, A., Bories, C., Figadère, B., Hocquemiller, R., Loiseau P.M., **2007**, *Biomed. Pharmacother.* 61, 441.
- [10] Geey, J.A De., Vam Sumare, C.F., **1966**, *Arch Int. Physiological Biochem.*, **74**, 512.
- [11] Fathi, R., Zhu Qus., 2004, *Appl Publ.*, 43,
- [12] Sanket, P., Chaudhari, Nandini, R., **2007**, *Ind. J. Het. Chem.* 17, 149.
- [13] Okuyama, E., Nishimura, S., Ohmori S., Ozaki Sutake, M., Yamazaki, M., **1993**, *Chem. Pharm. Bull.*, 41, 926.
- [14] Hudson, J.B., Eds, **1990**, "In Antiviral Compounds from plants", CRC Press, Boca Raton, Florida.
- [15] Hankare, P.P., Naravane, S.R., Bhuse, V.M., Debkar, S.D. Jagtap A.H., 2004, *Ind. J. Chem.*, 43A, 1464.
- [16] Geary, W.J., *Coord.* **1972**, *Chem. Rev.*, 1, 81.
- [17] Elzahany, E.A., Hegab, K.H., Khalil, S.K.H., Youssef, N.S., **2008**, *Austrelian J. Basic and Apl. Sci.* (2), 2220
- [18] Siddappa, K., Shikkargol, R.K., Angadi, S.D., **2009**, *Proc Indian Natn Sci. Acad* 75(2), 73.
- [19] Siddappa, K., Tukaram Reddy, P., Chandrakant Reddy, P., Mallikarjun, M., Mahesh, T., Mallikarjun K., **2008**. *International J. Pure & Appl. Chem.* 3(2), 87.
- [20] Prasad, R.N., Jain A., **2007**, *J. Indian Chem. Soc.*, 81, 319.
- [21] Mohamed, G.G., et.al., **2006**, *Turk J. Chem.*, 60, 366.
- [22] Mishra, A.P., Krishna, K., **2009**, *J. Indian Chem. Soc.*, **86**, 1150.
- [23] Chitra, G., **2007**, *J. Ind. Coun. Chem.*, 24(1), 16.
- [24] Chandra, S., Gupta, L.K., **2005**, *Ind. J. Chem. Soc.*, **82**, 454.
- [25] Eman, A., Khaled, H.H., Saffaa Khalil., Nabil S., Youssef., **2008**, *AJBAS* 2(2), 213.
- [26] Mashelkar, U.C., and Audi A.A., **2006**, *Ind. J. Chem.*, 4513, 1464.
- [27] Zhang, Y.H., Wang Q.L., Yang G.M., **2006**, *Trans. Met. Chem.*, 31, 856.
- [28] Siddappa, K., Tukaram Reddy., Chandrakant Reddy, P., Mallikarjun, M., Mallikarjun, K., **2008**, *Mat. Sci. Res. India.*, 5(1), 131.
- [29] Woolfson, M.M. **1980**, "An Introduction to X-ray Crystallography" 2nd edition Cambridge, Cambridge, University Press, 125.
- [30] Siddappa, K., Tukaram Reddy., Chandrakant Reddy, P. Mallikarjun, M., Mahesh Tambe, Mallikarjun. K., *Intr.* **2008**, *J. of Pure and Appl. Chem.*, 3(2), 87.
- [31] Siddappa, K., Tukaram Reddy., Mallikarjun, M., Reddy, CV., **2008**, *E-J. Chem.* 5(1), 155.
- [32] Chakrawarti, P.B., **2001**, *J Indian Chem Soc*, 78, 273.
- [33] Siddappa, K., Mallikarjun, K., Tukaram Reddy., Chandrakant Reddy, P., Mallikarjun, M., Mahesh Tambe., **2009**, *E-J. Chem.* 6(3), 615.
- [34] Bhattacharjee, C.R., Paul, S.B., Abhijit Nath., Choudhury, P.P.N., Sudip Choudhury., **2009**, *Materials*, 2, 345