Available online at www.scholarsresearchlibrary.com



Scholars Research Library

Archives of Applied Science Research, 2013, 5 (6):239-247 (http://scholarsresearchlibrary.com/archive.html)



Theoretical study of the effects of solvents on energy components of -acceptor-1,3,5-trinitrobenzene

K. K. Srivastava^a, Shubha Srivastava^b, Md. Tanweer Alam^a and Rituraj^a

^aDepartment of Chemistry, Vinoba Bhave University, Hazaribagh ^bDepartment of Chemistry, KBW College, Hazaribagh

ABSTRACT

Quantum mechanical calculations of different energies components of 1,3,5-trinitrobenzene (TNB) in ground state were carried out by DFT method, in isolated state and in various solvents to study the effects of solvents on various energy components. The solvation energy, chemical potential, hardness, electrophilicity of picric acid were calculated with the help of computed HOMO-LUMO gap of picric acid in different solvents in ground state. The plots of energy components and thermodynamic parameters against the dielectric constant of the corresponding solvents were found to be polynomial of higher order. The 3D plot of HOMO-LUMO of TNB and dielectric constants of various solvents in ground state reveals that LUMO of TNB is more affected than that of HOMO by change in the dielectric constant of the solvent

Keyword: DFT, Ground state, TNB, energy components, HOMO, LUMO, solvation energy, dielectric constants.

INTRODUCTION

Physical and chemical property of a molecule depends on the structure and the various kinds of energies of the molecule. Chemical reaction of a molecule in solution is affected by the nature of the solvent; solvent affects not only the energies of HOMO and LUMO of the molecule, but also other kinds of energies. Energy of a molecule may be considered to have various energy components such as reaction field energy, total zero-electron terms, Nuclear-nuclear, Nuclear-solvent, total one-electron terms, Electron-nuclear, Electron-solvent, Kinetic, total two-electron terms, Electronic energy, total quantum mech. energy, Gas phase energy, Solution phase energy, total solute energy, total solute energy, total solute energy, total solute energy, total Gibbs free energy, and zero point energy. TNB is a pale green yellow crystalline, explosive solid[1-2], It is used in preparation of Charge transfer complexes[3-13] of various utilities. Charge-transfer interactions between aromatic amines and trinitrobenzene have also been utilized for the distinction and determination of some primary, secondary, and tertiary aromatic amines [14]. Keeping in view the utility of TNB various kinds of energies of TNB in the ground state in gaseous phase and in different kinds of solvents have been theoretically calculated in this paper.

Computational methods

The initial structure of TNB was built with Chem-Draw ultra8.0 and the structure was optimized on Chem3D ultra 8.0. The structure was exported to Maestro 9.3 of Schrodinger 2012 version. The optimization of the structure was done on the Jaguar panel of the Maestro 9. The DFT-BPLY-3 method of theory was chosen. 6-31g ^{##} basis set was

selected and 255 basis functions were created for calculation. The molecule was assigned net zero charge and singlet multiplicity. In the solvent menu of the jaguar panel PBF solver was used for optimization of the structure in both the gaseous and solution phase. The optimization gaseous state and in the different solutions were done inground state of the molecule.

Geometry optimization

for perform a geometry optimization one needs to guess at the geometry and the direction in which to search, a set of co-ordinates to optimize, and some criteria for when to optimization is complete. The search direction is obtained from the gradient of the energy and the initial Hessian. An initial Hessian(second derivative matrix or force constant matrix) and the gradient are used to define search direction that should result in lowering of energy. The choice if co-ordinate systems have a substantial impact on the convergence of the optimization. The ideal set of Co-ordinate is one in which the energy change along each co-ordinate is maximized, and the coupling between co-ordinates is minimized. Jaguar chooses the coordinate system by default. It has two options Cartesian and z-matrix that produces an efficient optimization requires an understanding of the coupling between simple internal co-ordinates.

For optimization to minimum energy structures, the convergence criterion for SCF calculation is chosen to assure accurate analyses gradients. For these jobs, a wave function is considered converged when the root mean square (RMS) change in density matrix element is less than the RMS density matrix element change criterion, whose default value is 5.0×10^{-6} . The geometry is considered to have converged when the energy of successive geometries and the elements of analyze gradients of the energy and the displacement has met convergence criteria. For optimization in solution, the default criteria are multiplied by a factor of three, and a higher priority is given to the energy convergence criterion. Thus if the energy change criterion is met before the gradient and displacement criteria have been met, the geometry is considered converged. The optimized geometry may not have a local minimization energy i,e it may have reside on a saddle. To know whether it is global minimization we look for the value of vibrational frequencies. If all the vibrational frequencies are real (i,e+ve) then it is local minimum.

Performing a solvation calculation

It involves several iterations in which the wave functions for the molecule in the gas phase are calculated. The program ch performs electrostatic potential fitting, which represents the wave function as a set of point charges on the atomic centers. The interactions between the molecule and the solvent are evaluated by Jaguar's Poisson-Boltzmann solver[12-13], which fits the field produced by the solvent dielectric continuum to anotherset of point charges. These charges are passed back to scf, which performs a new calculation of the wave function for the molecule in the field produced by the solvent point charges. Electrostaticpotential fitting is performed on the new wave function, the solvent-molecule interactionsare reevaluated by the Poisson-Boltzmann solver, and so on, until the solvation free energy for the molecule converges.

For solvation calculations on neutral systems in water the program pre evaluates the Lewis dot structure for the molecule or system and assigns atomic van der Waals radii accordingly. Thesevan der Waals radii are used to form the boundary between the solvent dielectric continuum and the solute molecule. The Lewis dot structure and van der Waals radii information both appear in the output from the program pre. The radii are listed under the heading "vdw2" in the table of atomic information below the listing of non-default options. After the pre output, the usual output appears for the first, gas-phase calculation, except that the energy breakdown for the scf output also describes the electron-nuclear and kinetic contributions to the total one-electron terms in the energy, as well as the virial ratio – V/T, where V is the potential energy and T is the kinetic energy. This ratio should be -2 if the calculation satisfies the virial theorem. After the first scf output, the output from the first run of the program ch appears. Since performing a solvation calculation enables electrostatic potential fitting to atomic centers, the usual output for that option is included every time output from the program ch appears in the output file. The post program writes out the necessary input files for the Poisson-Boltzmann solver; this step is noted in the output file. The next output section comes from the Poisson-Boltzmann solver. The output includes information on the area (in Å2) of the molecular surface formed from the intersection of spheres with the van der Waals radii centered on the various atoms; the reaction field energy in kT (where T = 298 K), which is the energy of the interaction of the atom-centered charges with the solvent; the solvent-accessible surface area (in Å2), which reflects the surface formed from the points whose closest distance from the molecular surface is equal to the probe radius of the solvent; and the cavity energy in kT, which is computed to be the solvation energy of a nonpolar solute whose size and shape are the same as those of the actual solute molecule. The output from the program solv follows the Poisson-Boltzmann solver results,

K. K. Srivastava et al

giving the number of point charges provided by the solver to model the solvent, the sum of the surface charges, the nuclear repulsion energy already calculated by Jaguar, the nuclear-point charge energy representing the energy of interaction between the molecule's nuclei and the solvent point charges, and the point-charge repulsion energy, which is calculated but not used because it is irrelevant to the desired solvation results. After this output, the output for the second solvation iteration begins. The output from scf comes first, giving the results for the molecule-andsolvent-point-charges system. Total quantum mech. energy corresponds to the final energy from the scf energy table for that iteration, and includes the entire energies for the molecule-solvent interactions. The output next includes the gas phase and the solution phase energies for the molecule, since these terms are, of course, necessary for solvation energy calculations. The first solution phase energy component is the total solute energy, which includes the nuclear-nuclear, electronnuclear, kinetic, and two-electron terms, but no terms involving the solvent directly. The second component of the solution phase energy is the total solvent energy, which is computed as half of the total of the nuclear-solvent and electron-solvent terms, since some of its effect has already changed the solute energy. Third, a solute cavity term, which computes the solvation energy of a nonpolar solute of identical size and shape to the actual solute molecule, as described in reference [12], is included. This is only done for water as solvent. The last solution phase energy component (shown only if it is nonzero) is term (T), the first shell correction factor, which depends on the functional groups in the molecule, with atoms near the surface contributing most heavily. Finally, the list ends with the reorganization energy and the solvation energy. The reorganization energy is the difference between the total solute energy and the gas phase energy, and does not explicitly contain solvent terms. The final solvation energy is calculated as the solution phase energy described above minus the gas phase energy. The solvation energy is listed in Hartrees and in kcal/mol,

Chemical potential (µ)[14]

HOMO as ionization energy(IE) and LUMO as electron affinity (EA) have been used for calculating the electronic chemical potential (m) which is half of the energy of HOMO and LUMO

 $\mu = (E_{HOMO} + E_{LUMO})/2$

Hardness (η) [15]

The hardness (h) as half of the gap energy of HOMO and LUMO has been calculated using the following equation:

 $Gap = E_{HOMO} - E_{LUMO}$

 $\eta = Gap/2$

Electrophilicity (*w*)[16]

The electrophilicity (ω) has been calculated using equation

 $\omega = \mu^2/2 \eta$

Reaction field energy (in KT)

This gives us the energy of the interactions of atom centered charges with the solvent; Solvent accessible surface area (SASA in A^{0^2}) reflects the surface formed form the points whose closest distance from the molecular surface is equal to the probe radius of the solvent.

Cavity energy (in KT)

This is solvation energy of a non-polar solute whose size and shape are the same as those of actual solute molecule.

Quantum mechanical energy

This term corresponds to the entire energies for the molecule solvent interaction and is equal to the sum of total zero electron terms and electronic energy.

Reorganisation energy

This is the difference between the total solute energy and the gas phase energy, and does not explicitly contain solvent terms.

K. K. Srivastava et al

RESULTS AND DISCUSSION

Solvent parameters:

Table-1 summarizes the solvent parameters such as dielectric constants, molecular weight, density and polarity of the solvents used for the present theoretical study by Poisson-Boltzmann solver. In table-3 the energy components calculated by DFT method on Jaguar panel of the Maestro 9.3 with 6-31g^{##} basis set utilizing 255 basis functions for 1,3,5-trinitrobenzene in the ground state have been incorporated. The pictures of HOMO and LUMO of TNB in gaseous state and in various solvents have been shown in fig.1

An electron acceptor represents the ability to obtain an electron in the LUMO and HOMO represents the ability to donate electron.

The (E_{HOMO} - E_{LUMO}) gap is an important scale of stability [17] and compounds with large (E_{HOMO} - E_{LUMO}) gap value tend to have higher stability. The perusal of the table-2 indicates the stability of TNB increases in the solvents in the ground state in the order;

methanol>acetonitrile=cyclohexane>dmf>carbontetrachloride> benzene> dichlormethane >THF> water

The plot of the energy gap between HOMO and LUMO versus dielectric constant of solvents in ground state have been shown in the fig 2.

The dependence of the energy gap (y) on dielectric constant (x) in ground state follows the equation $y = -4E-08x^5 + 6E-06x^4 - 0.0004x^3 + 0.0078x^2 - 0.0591x - 5.2127$, (R² = 0.9943).

The TNB molecule has been found to be stabilized in the ground state in

methanol>acetonitrile=cyclohexane>dmf>carbontetrachloride> benzene> dichlormethane>THF> water.

Therefore, if it is desired to stabilize TNB in the ground state then out of ten solvents studied methanol is the best.

The 3D plots of HOMO-LUMO and dielectric constant shown in figure 7. reveals that LUMO of picric acid is more effected than HOMO in both the ground state by change in the dielectric constant of the solvent.

The chemical potentials(μ) of TNB in the ground state increases in the order;

Water> methanol> acetonitrile> dmf> dichloromethane> THF> chloroform> benzene> carbontetrachloride> cyclohexane.

The plot of the chemical potential versus dielectric constant of solvents in ground state have been shown in the fig3.

The dependence of the chemical potential (y) on dielectric constant(x) follows the equation $y = 7E-06x^3 - 0.0009x^2 + 0.0303x - 6.2436$ (R² = 0.9919)

The TNB molecule has been found to possess higher chemical potential in the ground state in benzene, chloroform, gas, methanol, water in the order;

Water> methanol> acetonitrile> dmf> dichloromethane> THF> chloroform> benzene> carbontetrachloride> cyclohexane.

Therefore, if it is desired to have highest chemical potential, TNB in the ground state, then out of ten solvents studied water is the best.

The hardness(η) of TNB increases in the ground state in the following order;

Chloroform > water > THF = dichloromethane > benzene > carbon tetrachloride > dmf = aceton itrile > methanol = cyclohexane = c

Scholars Research Library

The plot of hardness versus dielectric constant of solvents in the ground state have been shown in the fig4.

The dependence of hardness (y) on dielectric constant(x) follows $y = 2E-08x^5 - 3E-06x^4 + 0.0002x^3 - 0.0039x^2 + 0.0296x + 2.6064$, $R^2 = 0.9943$

The TNB molecule has been found to be hardest in chloroform in the ground state. Therefore, if it is desired to increased hardness of TNB to largest extent in the ground state then out of ten solvents studied chloroformis the best.

Solvents	M.W	Density	Dielectric constant	Probe radius	
	g/mol	g/ml		A ^o	
1.Acetonitrile	37.5	0.777	37.5	2.19	
2.Benzene	78.12	0.87865	2.284	2.6	
3.Carbontetrachloride	153.82	1.594	2.238	2.67	
4.Chloroform	119.38	1.4832	4.806	2.52	
5.Cyclohexane	84.16	0.77855	2.023	2.78	
6.Dichloromethane	84.93	1.3266	8.93	2.33	
7.DMF	73.09	0.944	36.7	2.49	
8.methanol	32.04	0.7914	33.62	2	
9.THF	72.11	0.8892	7.6	2.52	
10.Water	18.02	0.99823	80.37	1.4	

Table1. Physical parameters of various solvents

Table 2 Values of HOMO-LUMO energy, µ, n.	ω of TNR calculated by DFT -R3LVP/6-31G- level
Table 2. Values of HOMO-LOMO chergy, μ , η ,	to of 110 calculated by DF1 -D5E1170-510- level

Solvents	HOMO	LUMO	Gap	µ=Ehomo+Elumo/2	η=(Lumo-Homo)/2	Electrophilicity
HOMO, LUMO energy in eV						$\omega = \mu^2/2 \eta$
acetonitrile	-8.630	-3.327	-5.303	-5.979	2.652	0.064
benzene	-8.835	-3.523	-5.311	-6.179	2.656	0.068
carbontetrachloride	-8.837	-3.527	-5.310	-6.182	2.655	0.069
chloroform	-8.780	-3.425	-5.355	-6.102	2.678	0.067
cyclohexane	-8.847	-3.544	-5.303	-6.195	2.651	0.069
dichlormethane	-8.710	-3.373	-5.337	-6.042	2.668	0.066
dmf	-8.635	-3.332	-5.304	-5.983	2.652	0.064
methanol	-8.629	-3.327	-5.302	-5.978	2.651	0.064
THF	-8.732	-3.387	-5.345	-6.060	2.673	0.066
Water	-8.631	-3.281	-5.349	-5.956	2.675	0.064

The electrophilicity (a) of TNB increases in ground state in the following order: *Carbontetrachloride=cyclohexane>benzene> chloroform>THF=dichloromethane> water=dmf=acetonitrile.*

The plot of electrophilicity (y) versus dielectric constant(x) of solvents in ground state have been shown in the fig5.

The dependence of the electrophilicity on dielectric constant follows the $y = 4E-11x^6 - 6E-09x^5 + 3E-07x^4 - 8E-06x^3 + 9E-05x^2 - 0.0009x + 0.0701$, ($R^2 = 1.0002$) in the ground.

The TNB molecule has been found to possess high electrophilicity in the ground in carbontetrachloride and cyclohexane. Therefore, if it is desired to increase electrophilicity of TNB to larger extent in the ground state, then out of ten solvents studied carbontetrachloride and cyclohexane are the best.

The Solvation energy of TNB in the ground state are in the following order; methanol>acetonitrile> dmf> dichloromethane> THF>chloroform> benzene> carbontetrachloride> water>cyclohexane.

The plot of the solvation energy versus dielectric constant of solvents in ground state have been shown in the fig.6.

The dependence of the solvation energy(y) on dielectric constant(x) follows $y = -1E-06x^5 + 0.0002x^4 - 0.014x^3 + 0.3368x^2 - 3.4151x + 0.1895$ R² = 0.999

1,3,5-trinitrobenzene is most highly solvated in the ground state in methanol than other studied solvents.

Correct distants											
Ground state			D		C [1] C	G 11	D' 11	1 6	a 1	THE	XX 7 /
Energy components, in ev	Gas-phase	Acetonitrile	Benzene	Carbon	Chloroform	Cyclonexane	Dichloro	dmf	methanol	IHF	water
				tetrachloride			methane				
(A)Total zero electon terms		26186.21	26221.75	26222.54	26208.57	26224.84	26201.42	26186.98	26186.01	26203.12	26187.67
(B)Nuclear-nuclear	26243.02	26241.33	26245.65	26245.84	26247.45	26245.58	26248.4	26241.51	26241.22	26248.11	26241.93
(C)Nuclear-solvent		-55.1226	-23.9049	-23.3075	-38.8804	-20.7432	-46.9824	-54.5319	-55.2101	-44.9861	-54.2619
(E)Total one electron terms	-83481.5	-83405.6	-83455.5	-83456.6	-83439.4	-83459.5	-83430.6	-83406.6	-83405.3	-83432.5	-83413.4
(F)Electron-nuclear		-106271	-106293	-106293	-106290	-106294	-106289	-106272	-106271	-106289	-106280
(G)Electron-solvent		53.64956	23.32805	22.74615	37.89271	20.24802	45.75774	53.07602	53.73365	43.82044	53.37672
(H)Kinetic		22812.18	22813.83	22813.88	22813.19	22813.98	22812.88	22812.23	22812.16	22812.99	22813.22
(I)Total two electron terms	34224.98	34204.55	34219.72	34220.09	34216.42	34220.7	34214.56	34204.85	34204.48	34214.88	34211.46
(L)Electronic energy (E+I)	-49256.5	-49201	-49235.8	-49236.5	-49223	-49238.8	-49216	-49201.8	-49200.8	-49217.7	-49202
(N)Total quantum mechanical energy(A+L)	-23013.5	-23014.8	-23014	-23014	-23014.4	-23013.9	-23014.6	-23014.8	-23014.8	-23014.5	-23014.3
(O)Gas phase energy		-23013.5	-23013.5	-23013.5	-23013.5	-23013.5	-23013.5	-23013.5	-23013.5	-23013.5	-23013.5
(P)Solution phase energy(Q+R+S)		-23014.1	-23013.7	-23013.7	-23013.9	-23013.7	-23014	-23014.1	-23014.1	-23014	-23013.7
(Q)Total solute energy(N-C-G)		-23013.3	-23013.4	-23013.4	-23013.4	-23013.4	-23013.4	-23013.3	-23013.3	-23013.4	-23013.4
(R)Total solvent energyC/2+G/2)		-0.73651	-0.28843	-0.28065	-0.49382	-0.24758	-0.61233	-0.72795	-0.73824	-0.58285	-0.44261
(S)Solute cavity energy	0	0	0	0	0	0	0	0	0	0	0.136838
(U)Reorganization energy(Q-O)		0.125323	0.023318	0.022234	0.061947	0.01754	0.090569	0.12348	0.125239	0.083756	0.059226
(V)solvation energy(P-O) kcal/mol		-14.090	-6.112	-5.957	-9.956	-5.303	-12.028	-13.935	-14.132	-11.506	-5.684

Table 3. Values of energy components of TNB in gaseous state and various solvents in ground state calculated by DFT -B3LYP/6-31G- level



Figure 1. Picture of HOMO-LUMO in the ground state in gaseous and ten various solvents

Figure 2. Effect of dielectric contant on the HOMO-LUMO gap of TNB in the GS



Scholars Research Library





100

80

60

dielectric constant

Scholars Research Library

40

20

0.000

-10.000

-20.000

Figure 7. Figure 3D plot of HOMO-LUMO-dielectric constant to study the effect on HOMO and LUMO of TNT with the change in the dielectric constant of solvents in ground state



CONCLUSION

The present study on solvent effect on the energy components of 1,3,5-trinitrobenzenein ground state by ten different solvents has lead us to conclude it is highly solvated in methanol while lowest in cyclohexane. It has been found that TNB is most hard, electrophilic and chemical potential in chloroform, carbontetrachloride (also in cyclohexane and benzene), and water respectively while least in methanol (also in cyclohexane) ,water(DMF and acetonitrile) and cyclohexane respectively.

Acknowledgment

We are thankful to Will Richard, Raghu Rangaswamy, Vinod Dewarjee for providing the Shrodinger2012 software.

REFERENCES

[1] G. M. Murray, B. M. Arnold and D. S. Lawrence, US Pat. Appl.,2001077664, 2001.

[2]Sarah J. Toal and William C. Trogler, Journal of Materials Chemistry 2006 DOI: 10.1039/b517953j

[3] Ramadan RM, El-Atrash AM, Ibrahim AMA, Etaiw SEH (1991). Thermochim. Acta, 178, 331-338.

[4] Foster, R. & Fyfe, C. A. (1966) J. Chem. Soc. B 926-929.

[5] M. T. Sung and J. A. Parker, Proc. Nat. Acad. Sci. USA Vol. 69, No. 5, pp. 1196-1200, May 1972.

[6]P.Pfeiffer, "Organische Molekulverbindungen," 2nd edition, Ferdinand Enke, Stuttgart ,**1927**

[7] Allison, A. C., Andnash, T. Nature, 197:758-63, 1963.

[8] A.M.Hindawey, A.M.G. Nassar, R, M.Issa and Y.M.Issa. *Indian Journal of Chemistry*, Vol 19A, July 1980.615-619.

[9] H.A. Dessouki, M. Gaber, R.M.Issa and E.E.Hossani, *J.Indian Chem.Soc.* volLXV, April 1988, pp. 258-262,
[10]Moamen S. Refat, *Journal of Molecular Structure* 985 2011, 380–390

[11]J. P. Sharma and R.D. Tiwari (1972), *Microchem. J.*, 17, 2, 151-159.

[12]Tannor, D. J.; Marten, B.; Murphy, R.; Friesner, R. A.; Sitkoff, D.; Nicholls, A.; Ringnalda, M.; Goddard, W. A., III; Honig, B. J. Am. Chem. Soc. **1994**, *116*, 11875

[13]Marten, B.; Kim, K.; Cortis, C.; Friesner, R. A.; Murphy, R. B.; Ringnalda, M. N.; Sitkoff, D.; Honig, B. J. Phys. Chem. **1996**, 100, 11775.

[14]Parr, R. G.; Pearson, R. G. J. Am. Chem. Soc., 1983, 105, 7512.

[15] Pearson, RG 1991, Density functional theory-electronegativity and hardness chemtracts- Inorg.chem. 3:317-333
[16] Parr, R.G.; Yang, W. Chemical potential derivatives. In *Density-Functional Theory of Atoms and Molecules*, 1st ed.; Oxford University Press: New York, NY, USA, **1989**; pp. 87–95.

[17]Johnson, E. R.; Keinan, S.; Mori-Sanchez, P.; Contreras-Garcia, J.; Cohen, A. J.; Wang, W. J. Am. Chem. Soc. 2010, 132, 6498.